



Supplement of

Decline in Etesian winds after large volcanic eruptions in the last millennium

Stergios Misios et al.

Correspondence to: Stergios Misios (smisios@noa.gr)

The copyright of individual parts of the supplement might differ from the article licence.

Supplementary Material: Decline of Etesian winds after large volcanic eruptions in the last millennium

Stergios Misios^{1,2}, Ioannis Logothetis³, Mads F. Knudsen^{2,4}, Christoffer Karoff^{2,4}, Vassilis Amiridis¹, Kleareti Tourpali³

¹ Institute for Astronomy, Astrophysics, Space Applications, and Remote Sensing, National Observatory of Athens, Athens, Greece

² Department of Geoscience, Aarhus University, Aarhus, Denmark

³ Aristotle University of Thessaloniki, Thessaloniki, Greece

⁴ iCLIMATE Interdisciplinary Centre for Climate Change, Aarhus University, Aarhus, Denmark

Correspondence to: Stergios Misios (smisios@noa.gr)




Figure S1 Similar to Figure 3 but for all selected strong eruptions from 850 to 2005 listed in Table 1.




Figure S2 Similar to Figure 5 but for all selected strong eruptions from 850 to 2005 listed in Table 1.




Figure S3 Similar to Figure 5 but for surface temperature anomalies only (K) after the zonal mean response is subtracted.




Figure S4 Similar to Figure 6 but for all selected strong eruptions from 850 to 2005 listed in Table 1.