



# Influence of mid-latitude sea surface temperature fronts on the atmospheric water cycle and storm track activity

Fumiaki Ogawa<sup>1,2,3,4</sup> and Thomas Spengler<sup>3,4</sup>

<sup>1</sup>Research Center for Advanced Science and Technology, University of Tokyo, Meguro, Japan

<sup>2</sup>Faculty of Science, Hokkaido University, Sapporo, Japan

<sup>3</sup>Geophysical Institute, University of Bergen, Bergen, Norway

<sup>4</sup>Bjerknes Centre for Climate Research, Bergen, Norway

**Correspondence:** Fumiaki Ogawa (fumiaki.ogawa@atmos.rcast.u-tokyo.ac.jp)

Received: 12 March 2024 – Discussion started: 14 March 2024

Revised: 15 June 2024 – Accepted: 20 June 2024 – Published: 15 August 2024

**Abstract.** The climatological mean enhancement of the turbulent air–sea heat exchange along mid-latitude sea surface temperature (SST) fronts that anchor mid-latitude storm tracks suggests its crucial role in the atmospheric water cycle and storm tracks through the intensification of atmospheric cyclones and their associated precipitation. We investigate the sensitivity of the atmospheric water cycle to the SST front through a set of aqua-planet experiments. Varying the latitude of a zonally symmetric mid-latitude SST front, the mid-latitude atmospheric water cycle responds through distinct changes in surface latent heat fluxes, precipitation, and atmospheric moisture fluxes, whereas the tropical latitudes remain largely unchanged. As storm tracks are self-maintained through the diabatic generation of eddy available potential energy, the position of the storm track is diabatically anchored at the SST front. While the position of the SST front determines the position of the eddy moisture convergence and thus the diabatic heating that energises the storm track, the underlying SST determines the general strength of the water cycle and thus the intensity of the storm track. The strong connection identified between the eddy moisture flux and the SST front implies a diabatic pathway of latent heating to anchor the storm track along SST fronts.

## 1 Introduction

The climatological mean vertically integrated global atmospheric water budget obeys the balance

$$C = P - E, \quad (1)$$

with evaporation ( $E$ ), precipitation ( $P$ ), and convergence ( $C$ ) of the vertically integrated moisture flux ( $F$ ) (e.g., Oki et al., 2004; Trenberth et al., 2007; Schneider et al., 2010; Seager et al., 2010; Trenberth, 2011; Hartmann, 2016; Dey and Döös, 2019). Given that  $P$  and  $E$  in the extra-tropics are maximised, respectively, along and on the equatorward flank of mid-latitude sea surface temperature (SST) fronts associated with the confluent regions of warm and cool oceanic western boundary currents (Zolina and Gulev, 2003; Nakamura et al., 2008; Tilinina et al., 2018; Ogawa and Spengler, 2019), the global atmospheric water cycle is most likely highly sensitive to the position and characteristics of SST fronts. While the response of the mid-latitude atmospheric circulation to SST fronts has been well studied (e.g., Nakamura et al., 2004, 2008; Sampe et al., 2010; Ogawa et al., 2012), the response of the water cycle and related implications for the position and strength of the storm track are largely unknown. Hence, we investigate the sensitivity of the global atmospheric water cycle and storm track to the position and strength of SST fronts.

With  $P < E$  in the subtropical latitudes and  $P > E$  in both the equatorial and the mid-latitudes,  $F$  needs to transport moisture from the subtropics to both the inner tropics and mid-latitudes (Bryan and Oort, 1984).  $F$  can be separated into contributions from the time-mean flow ( $F_m$ ) and

eddies ( $F_e$ ) (Hartmann, 2016), where the relative contributions of  $F_m$  and  $F_e$  to  $F$  vary with latitude. While  $F$  and  $C$  in the tropics are mainly determined by  $F_m$ , due to the Eulerian mean characteristics of the Hadley cell,  $F$  and  $C$  are dominated by  $F_e$  in the mid-latitudes due to the prevalence of atmospheric transient eddies (Peixoto and Oort, 1992; Newman et al., 2012). These transient eddies, associated with the mid-latitude storm tracks, play a crucial role in the hydrologic cycle in atmosphere–ocean coupled climate simulations (Dey and Döös, 2019), where most of the transient moisture transport occurs in warm conveyor belts associated with these synoptic eddies (Zhu and Newell, 1998; Neiman et al., 2008; Madonna et al., 2014).

Given that storm tracks in the mid-latitudes tend to be anchored to mid-latitude SST fronts (Nakamura et al., 2004, 2008; Sampe et al., 2010; Ogawa et al., 2012; Deremble et al., 2012), the position and strength of SST fronts most likely have significant ramifications for  $F_e$  and  $C$ , contributing to the self-maintenance of storm tracks through the latent heat release associated with  $P$ , which is maximised along the SST fronts due to the collocated storm track activity (Hoskins and Valdes, 1990; Nakamura et al., 2008; Papritz and Spengler, 2015). Furthermore, eddy meridional surface wind yields a maximum in  $E$  on the equatorward flank of mid-latitude SST fronts (Zolina and Gulev, 2003; Tilinina et al., 2018; Ogawa and Spengler, 2019). Despite these strong constraints of the eddy-driven atmospheric water cycle to mid-latitudes SST fronts, the sensitivity of the water cycle to the position and strength of SST fronts has not been investigated systematically.

Prescribing zonally symmetric mid-latitude SST fronts in idealised aqua-planet simulations highlights a strong sensitivity of the zonal mean atmospheric circulation as well as the latitude of the storm track to the position of SST fronts (Ogawa et al., 2012, 2016). As both low-level atmospheric baroclinicity associated with the SST front (Nakamura et al., 2008; Sampe et al., 2010; Hotta and Nakamura, 2011) and latent heat release in mid-latitude storms (Hoskins and Valdes, 1990; Deremble et al., 2012; Papritz and Spengler, 2015) have been argued to maintain the location of the storm track, it is important to further clarify the constraints of SST fronts on the atmospheric water cycle and their relevance to the position and intensity of storm tracks. We quantify this sensitivity of the mean meridional atmospheric water cycle and storm track intensity to the existence and latitude of mid-latitude SST fronts using a set of aqua-planet atmospheric general circulation model (AGCM) experiments.

## 2 Aqua-planet experiments

We conduct a set of aqua-planet AGCM experiments with prescribed zonally symmetric SSTs featuring a mid-latitude SST front. We use the AGCM of the Earth Simulator (Ohfuchi et al., 2004; Enomoto et al., 2008; Kuwano-Yoshida

et al., 2010, AFES;) with a triangular truncation of T79, corresponding to  $\approx 1.5^\circ$  grid spacing in longitude and latitude. Though the resolution is insufficient to resolve mesoscale features of the SST, the resolution is sufficient to realistically represent both the climatological mean state and annular mode variability (Nakamura et al., 2008; Sampe et al., 2010, 2013; Ogawa et al., 2012, 2015, 2016). The model has 56  $\sigma$  levels in the vertical that reach up to 0.09 hPa, which is well above the stratopause. The model was integrated for 120 months using the perpetual austral winter solstice condition, where we disregard the first 6 months as spin-up.


The convection scheme employed in this model is Emanuel (1991), which calculates sub-grid-scale convection processes and the associated condensation, precipitation, and latent heating. The large-scale condensation scheme in this model follows the framework of Le Trent and Li (1991), which represents cloud-related condensation processes other than cumulus convection.

The default SST profile is based on the observed SST averaged over the south Indian Ocean ( $60\text{--}80^\circ\text{E}$ ) using the Optimum Interpolation SST (OISST) data (Reynolds et al., 2007). The observed climatological mean south Indian Ocean SST for austral winter (June–July–August) was assigned to the Southern Hemisphere, and the corresponding austral summertime profile (December–January–February) was assigned to the Northern Hemisphere (solid black line in Fig. 1). Henceforth, we will refer to the Southern Hemisphere and Northern Hemisphere as the winter hemisphere and summer hemisphere, respectively. The SST is identical to the one used in Ogawa et al. (2016).

For the sensitivity experiments, the magnitude of the SST gradient was kept identical, while the latitude of the SST fronts in both hemispheres was shifted to either  $35$  or  $55^\circ$  (red and blue lines in Fig. 1b). We refer to the experiments as F35, F45, and F55. In addition, we performed an experiment using a non-front reference (REF) SST profile, where the mid-latitude SST front is smoothed while maintaining the same SST gradient poleward of the SST front (grey line in Fig. 1). More details about the model setup can be found in Ogawa et al. (2016). The REF experiment is identical to the NF experiment in Ogawa et al. (2016), though it is referred to as REF because of the presence of a weak local maximum of the subtropical SST gradient around  $33^\circ$  in both hemispheres.

For all experiments, the SST equatorward of  $25^\circ$  is unchanged (Fig. 1a) to avoid a direct impact on the Hadley cell (Höjgård-Olsen et al., 2020), which would yield a remote influence on the mid-latitude jets (Watt-Meyer and Frierson, 2019). The experiments in this study are not meant to reproduce the observed atmospheric circulation but to clarify the impact of the mid-latitude SST front on the atmospheric circulation and the atmospheric water cycle.

In addition, we performed experiments where we uniformly increased or decreased the SST by 5 K for the F45 and REF setup. We refer to these as F45+5 and F45–5 for simulations with the SST front at  $45^\circ$  and a 5 K higher and lower



**Figure 1.** (a) Meridional profiles of the prescribed zonally symmetric sea surface temperature [°C] and (b) their meridional gradient [K 10<sup>-2</sup> km<sup>-1</sup>]. Colours (red, black, blue) indicate the latitude of the SST front (35, 45, and 55°) as indicated by the triangles. The grey line corresponds to the non-front reference (REF) experiment.

SST, respectively. Similarly, we refer to REF+5 and REF-5 for simulations with the reference state without an SST front and a 5 K higher and lower SST, respectively. The SST gradients for these experiments are identical to F45 and REF (see Fig. 1b), while the SST is uniformly higher or lower, respectively (not shown).

We use 6-hourly instantaneous fields of surface pressure ( $p_s$ ), specific humidity ( $q$ ), evaporation ( $E$ ), convective precipitation ( $P_c$ ), large-scale precipitation ( $P_l$ ), zonal ( $u$ ) and meridional wind ( $v$ ), vertical pressure ( $\omega$ ), and sigma velocity ( $\dot{\sigma}$ ), as well as 6-hourly integrated values of changes in temperature ( $Q$ ) and specific humidity due to parameterisations ( $\dot{q} = \dot{q}_c + \dot{q}_m + \dot{q}_b$ ), which is split up in tendencies from convection parameterisation ( $\dot{q}_c$ ), microphysics parameterisation ( $\dot{q}_m$ ), and boundary layer parameterisation ( $\dot{q}_b$ ). We use data at all available  $\sigma$  levels.

### 3 Diagnosing the atmospheric water cycle

We use the tendency equation for specific humidity (Eq. 2) and continuity in sigma coordinates (Eq. 4),

$$\frac{\partial q}{\partial t} + \frac{u}{a \cos \phi} \frac{\partial q}{\partial \lambda} + \frac{v}{a} \frac{\partial q}{\partial \phi} + \dot{\sigma} \frac{\partial q}{\partial \sigma} = \dot{q}, \quad (2)$$

$$\begin{aligned} \frac{\partial p_s}{\partial t} + \frac{1}{a \cos \phi} \frac{\partial (p_s u)}{\partial \lambda} + \frac{1}{a \cos \phi} \frac{\partial (p_s v \cos \phi)}{\partial \phi} \\ + \frac{\partial (p_s \dot{\sigma})}{\partial \sigma} = 0, \end{aligned} \quad (4)$$

to derive the flux form for specific humidity:

$$\begin{aligned} \frac{\partial (p_s q)}{\partial t} + \frac{1}{a \cos \phi} \frac{\partial (p_s u q)}{\partial \lambda} + \frac{1}{a \cos \phi} \frac{\partial (p_s v q \cos \phi)}{\partial \phi} \\ + \frac{\partial (p_s \dot{\sigma} q)}{\partial \sigma} = p_s \dot{q}, \end{aligned} \quad (5)$$

where the wind vector in sigma coordinates is  $\mathbf{v} = (u, v, \dot{\sigma})$ ,  $q$  is specific humidity,  $p_s$  is surface pressure,  $a$  is the radius of Earth, and  $\phi$  and  $\lambda$  are the latitude and longitude, respectively.

Due to the zonal symmetry of the experiments, we use Eq. (5) to derive the atmospheric water cycle for the zonally and time-averaged climatological mean state:

$$\begin{aligned} \nabla \cdot \mathbf{F} &= \frac{1}{a g \cos \phi} \frac{\partial \left( \overline{[p_s v q]} \cos \phi \right)}{\partial \phi} + \frac{1}{g} \frac{\partial \overline{[p_s \dot{\sigma} q]}}{\partial \sigma} \\ &= \overline{[p_s \dot{q}]}, \end{aligned} \quad (6)$$

where  $\frac{\partial (p_s q)}{\partial t} = 0$ ,  $g$  is the gravitational constant,  $\mathbf{F} = \frac{1}{g} \left( 0, \overline{[p_s v q]}, \overline{[p_s \dot{\sigma} q]} \right)$  is the moisture flux, and the squared brackets  $[f]$  and the bar  $\bar{f}$  denote the zonal average and time mean, respectively. Given the dominant contribution of the eddy moisture fluxes in the extra-tropics (e.g., Peixoto and Oort, 1992; Newman et al., 2012), we separate the variables ( $f = \bar{f} + f'$ ) into a time mean ( $\bar{f}$ ) and eddy ( $f'$ ) component yielding

$$\nabla \cdot \mathbf{F} \approx \nabla \cdot (\mathbf{F}_m + \mathbf{F}_e), \quad (7)$$

with  $\mathbf{F}_m = \frac{1}{g} \left( 0, \overline{[p_s \bar{v} \bar{q}]}, \overline{[p_s \bar{\sigma} \bar{q}]} \right)$  and  $\mathbf{F}_e = \frac{1}{g} \left( 0, \overline{[p_s v' q']}, \overline{[p_s \dot{\sigma}' q']} \right)$  being the mean and eddy components, where we excluded terms proportional to  $\bar{q} p'_s v'$ ,  $\bar{v} p'_s q'$ ,  $\bar{q} p'_s \dot{\sigma}'$ ,  $\bar{\sigma} p'_s q'$ ,  $\dot{\sigma}' p'_s q'$ , and  $v' p'_s q'$ , as they turned out to be negligible (not shown).

Integrating Eq. (6) vertically and letting the fluxes at the surface be precipitation ( $P$ ) and evaporation ( $E$ ) yield the atmospheric water cycle

$$C = P - E = P_l + P_c - E, \quad (8)$$

where  $P_l$  and  $P_c$  are large-scale and convective precipitation, respectively, and

$$\begin{aligned} C \approx C_m + C_e &= - \int_0^1 \nabla \cdot (\mathbf{F}_m + \mathbf{F}_e) d\sigma \\ &= - \frac{1}{a g \cos \phi} \frac{\partial}{\partial \phi} \int_0^1 \left( \overline{[p_s \bar{v} \bar{q}]} \cos \phi + \overline{[p_s v' q']} \cos \phi \right) d\sigma \end{aligned} \quad (9)$$

is the climatological mean vertically integrated atmospheric moisture flux convergence, where we used  $\bar{\sigma} = 0$  at  $\sigma = 0/1$ . The first and second terms in Eq. (9) indicate the moisture flux convergence by the mean ( $C_m$ ) and eddy ( $C_e$ ) flux, respectively.

**4 Energy framework for the storm track**

The climatological mean energy cycle can be diagnosed using the Lorenz framework (Lorenz, 1955; Oort and Peixoto, 1974; Peixoto and Oort, 1974), where the contribution of latent heating can be diagnosed as the diabatic eddy energy generation. Following Peixoto and Oort (1974), we use the climatological mean and zonally averaged atmospheric energy as well as its conversion terms defined on the two-dimensional latitude–pressure domain:

$$APE_Z = \frac{c_p}{2} \gamma [\overline{T}]^{\#2}, \tag{10}$$

$$APE_E = \frac{c_p}{2} \gamma [\overline{T'^2} + \overline{T^{*2}}], \tag{11}$$

$$KIN_Z = \frac{1}{2} ([\overline{u}]^2 + [\overline{v}]^2), \tag{12}$$

$$KIN_E = \frac{1}{2} [\overline{u'^2} + \overline{v'^2}] + \frac{1}{2} [\overline{u^{*2}} + \overline{v^{*2}}], \tag{13}$$

$$CE = -\frac{R}{p} [\overline{\omega'T'} + \overline{\omega^*T^*}], \tag{14}$$

$$CZ = \frac{R}{p} [\overline{\omega}]^{\#} [\overline{T}]^{\#}, \tag{15}$$

$$CA = -c_p \gamma [\overline{v'T'} + \overline{v^*T^*}] \frac{\partial [\overline{T}]}{a \partial \phi} - c_p p^{-\kappa} [\overline{\omega'T'} + \overline{\omega^*T^*}] \frac{\partial (\gamma p^{\kappa} [\overline{T}]^{\#})}{\partial p}, \tag{16}$$

$$CK = [\overline{v'u'} + \overline{v^*u^*}] \cos \phi \frac{\partial ([\overline{u}] / \cos \phi)}{a \partial \phi} + [\overline{v'^2} + \overline{v^{*2}}] \frac{\partial [\overline{v}]}{a \partial \phi} + [\overline{\omega'u'} + \overline{\omega^*u^*}] \frac{\partial ([\overline{u}])}{\partial p} + [\overline{\omega'v'} + \overline{\omega^*v^*}] \frac{\partial ([\overline{v}])}{\partial p} - [\overline{v}] [\overline{u'^2} + \overline{u^{*2}}] \frac{\tan \phi}{a}, \tag{18}$$

$$GE = \gamma [\overline{Q'T'} + \overline{Q^*T^*}], \tag{19}$$

$$GZ = \gamma [\overline{Q}]^{\#} [\overline{T}]^{\#}, \tag{20}$$

with  $\gamma = -\left(\frac{R\theta}{T c_p p}\right) \left(\frac{\partial \Theta}{\partial p}\right)^{-1}$ ,  $\theta = T(1000/p)^{\kappa}$ , and  $\kappa = \frac{R}{c_p}$ , where  $R$  is the gas constant,  $c_p$  is the specific heat at constant

pressure,  $\Theta$  is the global horizontal average of  $\theta$ ,  $Q$  is the diabatic heating rate, and superscripts \* and # denote anomalies from zonal and meridional averages, respectively. APE and KIN denote the available potential energy and kinetic energy, respectively. Eddy and mean-flow components in APE and KIN are denoted by the subscripts  $E$  and  $Z$ , respectively. CA is the energy conversion from  $APE_Z$  to  $APE_E$ . CE is the energy conversion from  $APE_E$  to  $KIN_E$ . CK is the energy conversion from  $KIN_E$  to  $KIN_Z$ . CZ is the energy conversion from  $KIN_Z$  to  $APE_Z$ . GZ and GE represent the generation of  $APE_Z$  to  $APE_E$ , respectively.  $DKIN_Z$ ,  $DKIN_E$ ,  $DAPE_Z$ , and  $DAPE_E$  depict the dissipation of  $KIN_Z$ ,  $KIN_E$ ,  $APE_Z$ , and  $APE_E$ , respectively.

The three-dimensional spatial mean energy budget can be expressed by

$$d \langle APE_Z \rangle / dt = -\langle CA \rangle + \langle CZ \rangle + \langle GZ \rangle - \langle DAPE_Z \rangle, \tag{21}$$

$$d \langle APE_E \rangle / dt = \langle CA \rangle - \langle CE \rangle + \langle GE \rangle - \langle DAPE_E \rangle, \tag{22}$$

$$d \langle KIN_Z \rangle / dt = \langle CK \rangle - \langle CZ \rangle - \langle DKIN_Z \rangle, \tag{23}$$

$$d \langle KIN_E \rangle / dt = \langle CE \rangle - \langle CK \rangle - \langle DKIN_E \rangle, \tag{24}$$


where  $\langle f \rangle$  indicates the average over the meridional plane.  $DAPE_Z$ ,  $DAPE_E$ ,  $DKIN_Z$ , and  $DKIN_E$  depict the dissipation of  $KIN_Z$ ,  $KIN_E$ ,  $APE_Z$ , and  $APE_E$ , respectively. The left-hand side is approximately zero climatologically, so the dissipation term, which appears as the last term on the right-hand side in Eqs. (21)–(23), was calculated as the residual to close the energy budget. We interpolate the data from  $\sigma$  to pressure levels and calculate vertically mass-weighted averages from 850 to 200 hPa. We obtain a reasonable closure of the energy cycle, where the total of CA and GE is very close to CE, which supplies the eddy kinetic energy ( $KIN_E$ ).

We restrict our discussion to the energy conversions relevant to the storm track activity: conversion of available potential energy from the zonal mean state to eddies (CA), baroclinic eddy energy generation (GE), baroclinic eddy energy conversion (CE), and the eddy kinetic energy ( $KIN_E$ ).

**5 Vertically integrated water cycle**

The climatological mean vertically integrated atmospheric moisture cycle shows the gain and loss of atmospheric water content as a function of latitude according to Eqs. (8) and (9), where the sum of the five different terms is approximately zero at every latitude, confirming climatological balance (Fig. 2). At lower latitudes, moisture is provided by  $E$  from the warm tropical ocean, which is subsequently converged into the inter-tropical convergence zones (ITCZs) in the summer hemisphere and winter hemisphere by  $C_m$ , where  $P_c$  removes the moisture from the atmosphere.

Two latitudinal peaks of  $P_c$  and the latitude of mean upward motion in the tropical latitudes (Figs. 3c–d and S1c–d in the Supplement) indicate the existence of a double ITCZ, which is presumably caused by the prescribed tropical SST



**Figure 2.** Meridional diagram of the simulated zonal mean evaporation (red line), large-scale precipitation (dashed blue line), cumulus precipitation (blue solid line), vertically integrated convergence of northward eddy moisture flux (dashed black line), vertically integrated convergence of northward mean-flow moisture flux (solid black line), evaporation minus precipitation (shading, red colour is for positive values and blue colour is for negative values) for (a) F35, (b) F45, (c) F55, and (d) REF experiments. Unit is  $[\text{kg m}^{-2} \text{d}^{-1}]$ .

(Watt-Meyer and Frierson, 2019). The stronger and more equatorward peak in  $P_c$  in the winter hemisphere compared to the summer hemisphere reflects the prescribed season. The similarity of the moisture cycle in the lower latitudes across all experiments (Fig. 2a–d) suggests that the position and existence of the mid-latitude SST front have little influence on the Hadley circulation (Figs. 3c, d, S1c, d).

In the mid-latitudes, however, the atmospheric moisture budget is sensitive to the latitudinal position as well as the existence of an SST front. Both  $C_e$  and  $P_l$  are maximised on the poleward flank of the SST front (dashed black and blue lines in Fig. 2a–c), which is consistent with the positioning of the storm track (Ogawa et al., 2012, 2016).  $E$  features a stark contrast across the SST front (red lines in Fig. 2a–c), with a notable increase from the poleward toward the equatorward side of the SST front. This increase is mostly due to the surface-saturation-specific humidity being a function of SST through the Clausius–Clapeyron relation and not the mean surface wind speed (not shown).

Similarly,  $P_c$  (solid blue lines in Fig. 2a–c) also features higher values on the equatorward side of the SST front.  $C_m$  is generally less than  $C_e$  near the SST front (solid and dashed black lines in 2a–d), highlighting the dominance of eddy contributions around the SST front. While the amount of air–sea moisture exchange is slightly less in the summer hemisphere, the SST front still strongly influences the atmospheric water budget in both hemispheres.


In REF (Fig. 2d), the absence of the SST front results in smoothed profiles in the extra-tropics without any distinc-

tive peaks or steep transitions as found in the experiments with an SST front (Fig. 2a–c). Given the absence of a peak in  $C_e$ , the moisture convergence by mid-latitude eddies is significantly reduced, which is consistent with a suppression of storm track activity in the absence of an SST front (Nakamura et al., 2008; Sampe et al., 2010, 2013). Comparing REF to the front experiments, it is also evident that the evaporation is strongly influenced by absolute SST, which is due to the Clausius–Clapeyron dependence of the saturation mixing ratio at SST.

## 6 Meridional overturning of atmospheric moisture

The meridional overturning of atmospheric moisture depicted by  $F$  reveals a clear separation of the tropical and extra-tropical cells for F45 (Fig. 3a, c). The moisture converges into the ITCZs in the tropical latitudes, where it is transported upward by the mean circulation (Fig. 3c). The subsiding branch in the subtropical latitudes splits in an equatorward and a poleward component, with significant divergence in the lower troposphere (Fig. 3c). The direction of  $F$  in the mid-latitudes features a clear poleward ascent (Fig. 3a), which is associated with the isentropic up-glide along the isentropes in the warm conveyor belts in extra-tropical cyclones (McTaggart-Cowan et al., 2017).

Both F45 and REF feature a baroclinic zone in the mid-latitudes, evident by the sloping isentropes across the entire troposphere (Fig. 3a, b). Comparing the lower-tropospheric



**Figure 3.** (a) Simulated zonally averaged mean eddy moisture flux (vector,  $[\text{kg m}^{-2} \text{d}^{-1}]$ ), its convergence (shading,  $[\text{kg m}^{-3} \text{d}^{-1}]$ ), and potential temperature (contour, [K]) for F45. (c) Zonally averaged mean-flow moisture flux (vector), its convergence (shading), and mass stream function (contour,  $[10^9 \text{ kg s}^{-1}]$ ) for F45. (b, d) Same as (a, c), but for REF.


isentropes, it is evident that the isentropes in F45 are more compact in the meridional direction compared to REF, implying a stronger low-level temperature gradient and baroclinicity in the presence of an SST front. The low-level temperature contrast from the subtropical to subpolar regions in the atmosphere is similar to the change in the contrast in the SST. Therefore, the SST distribution largely dominates the low-level atmospheric temperature field and baroclinicity.

In the tropical latitudes, the meridional overturning of atmospheric moisture is dominated by  $F_m$  (Fig. 3c), which is consistent with  $C_m$  (solid black line, Fig. 2b) being the main contributor to the divergence in the subtropics and convergence in the ITCZs (red line, Fig. 2b). Thus, the tropical circulation largely resembles the Hadley cell with the inner-tropical  $F$  being almost entirely explained by  $F_m$ . In the vicinity of the extra-tropical SST front, the amplification of  $F_m$  in the poleward direction near the surface is associated with the thermally indirect Ferrel cell (Fig. 3c).

The extra-tropics are dominated by  $F_e$ , which is ascending poleward along isentropic surfaces (Fig. 3a), consistent with the moisture fluxes along warm conveyor belts associated with mid-latitude cyclones (Madonna et al., 2014; Papritz and Spengler, 2015; McTaggart-Cowan et al., 2017). The poleward ascending flux implies a divergence in the lower

troposphere equatorward of the SST front and a convergence in the mid- to upper troposphere in the mid-latitudes, which is more or less centred on the SST front. Consistently,  $C_e$  peaks on the poleward side of the SST front, yielding a peak in  $P_1$  associated with the warm conveyor belts (Fig. 2b).

Varying the position of the SST front yields qualitatively similar results (Fig. S1). In the tropical latitudes,  $F_m$  and its associated divergence/convergence are almost identical across these experiments, whereas  $F_e$  and its associated divergence shift with the latitude of the SST front for F35, F45, and F55 in the lower troposphere (Figs. 3, S1). For REF, while  $F_m$  is again overall unchanged, the absence of the SST front results in a wider latitudinal spread of the lower-tropospheric divergence of  $F_e$ , stretching from the subtropics to the higher latitudes (Fig. 3b). The mid- and upper-tropospheric convergence of  $F_e$  is generally less affected by the position and existence of the SST front and mainly widens with a higher-latitude SST front or if no SST front is prescribed. The latter is also evident in a more widely spread  $P_1$  for F45 and F55 as well as for REF (Fig. 2).




**Figure 4.** (a) Mean moistening tendency due to micro-physics parameterisation (shading,  $[\text{kg m}^{-3} \text{d}^{-1}]$ ), potential temperature (black contour, [K]), and zonal wind (brown contours for a contour interval of  $15 [\text{m s}^{-1}]$  from  $15 [\text{m s}^{-1}]$ ) for F45. (c) Mean moistening tendency due to cumulus parameterisation (shading) and specific humidity (contour,  $[\text{g kg}^{-1}]$ ) for F45. (e) Mean moistening tendency due to diffusive process (shading) and specific humidity (contour) for F45. (b, d, f) Same as (a, c, e), but for REF.

### 7 Moisture flux convergence and parameterised processes

To achieve climatological balance in Eq. (6), the convergence of  $F$  (Fig. 3a, c) is counteracted by  $\dot{q}$ , which consists of the microphysics parameterisation ( $\dot{q}_m$ ), the convection parameterisation ( $\dot{q}_c$ ), and the boundary layer parameterisation ( $\dot{q}_b$ ) (Fig. 4a, c, e), where  $\dot{q}_m$  represents the removal of moisture by large-scale condensation,  $\dot{q}_c$  represents the removal of moisture by convective condensation, and  $\dot{q}_b$  represents the sum of the vertical diffusion of the moisture evaporated

from the ocean surface and the vertical redistribution of the moisture through the atmospheric dry convection. The distributions of  $\dot{q}_m$  and  $\dot{q}_c$  are strikingly different, where the vertical integrals of  $\dot{q}_m$  and  $\dot{q}_c$  (Fig. 4e, g) correspond to  $P_l$  and  $P_c$ , respectively (Fig. 2b).

$\dot{q}_m$  is mainly located in the mid-latitude mid-troposphere, which solely contributes to the removal of moisture from the atmosphere (Fig. 4a) and is balanced by the convergence of  $F_c$  (Fig. 3a). The moisture removal occurring along the isentropic slope, which is in thermal wind balance with the jets (Fig. 4a), corresponds to the precipitation formed along the



**Figure 5.** A schematic diagram illustrating the Lorenz energy cycle for F45 as calculated as the zonally and vertically weighted average from 850 to 200 hPa. Arrows indicate the direction of the positive energy conversion or generation. APE (KIN) stands for the available potential energy (kinetic energy). Subscripts Z and E are for the zonal mean state and eddies, respectively. Values with parentheses are calculated as the residual to close the budget. The unit for APE and KIN is  $[10^5 \text{ J m}^{-2}]$ , and it is  $[10^5 \text{ J m}^{-2} \text{ d}^{-1}]$  for the rest of the calculation. See text for more details.

up-gliding of moist air parcels in cyclones (Papritz and Spengler, 2015).  $\dot{q}_c$ , on the other hand, is balanced by the convergence of  $\mathbf{F}_m$  (Fig. 3c) and features losses of atmospheric moisture in the planetary boundary layer at all latitudes as well as within the vertical towers of the ITCZ (Fig. 4c).  $\dot{q}_c$  in the subtropical lower troposphere adds moisture to the atmosphere (Fig. 4c) and is largely compensated by the divergence of  $\mathbf{F}_m$  (Fig. 3c).

The secondary peak of  $P_c$  in the mid-latitudes (Fig. 2b) is associated with an increasing vertical extent of negative  $\dot{q}_c$  from the tropical latitudes towards the SST front, where  $\dot{q}_c$  drops dramatically (Fig. 4c). The gains of  $\dot{q}_c$  in the subtropical mid-troposphere are associated with a vertical rearrangement of moisture by the convection scheme (Fig. 4c) and are compensated by the divergence of  $\mathbf{F}_c$  and  $\mathbf{F}_m$  (Fig. 3a and c).

The contribution of  $\dot{q}_b$  represents the turbulent uptake of moisture from the ocean and its redistribution in the atmospheric boundary layer (Fig. 4i). The vertical integral of  $\dot{q}_b$  corresponds to  $E$  (red line in Fig. 2b) and accordingly features a stark contrast across the SST front. The moisture supplied by  $\dot{q}_b$  is largely compensated by  $\dot{q}_c$  (Fig. 4c, e). The areas of high climatological mean moisture amount in the subtropical planetary boundary layer appear to be largely determined by regions where  $\dot{q}_b$  is large (Fig. 4e). For REF there is no significant drop in  $\dot{q}_b$  and  $\dot{q}_c$  at the SST front; instead they gradually decrease towards higher latitudes (Fig. 4b, d).

Similar to  $\mathbf{F}$ ,  $\dot{q}$  in the lower latitudes is not sensitive to the position or existence of the SST front (Fig. S2). How-

ever, the stark contrast in vertical extent of  $\dot{q}_b$  and  $\dot{q}_c$  remains anchored to the SST front when the latter is shifted latitudinally (Fig. S4a–b). Similarly,  $\dot{q}_m$  is shifted with the SST front (Fig. S4c–f), which is consistent with the sensitivity of the vertically integrated moisture budget (Fig. 2).


## 8 Diabatic enhancement of storm track activity

The importance of the SST front for both the latitude of the storm track (Ogawa et al., 2012) and the eddy moisture flux convergence shown in this study implies a diabatic enhancement of storm track activity through energising individual storms via surface latent heat exchange (Hualand and Spengler, 2020; Bui and Spengler, 2021). We diagnose the Lorenz energy cycle (Lorenz, 1955; Oort and Peixoto, 1974), where the contribution of the latent heat can be diagnosed as the diabatic eddy energy generation. Considering pressure levels from 850 to 200 hPa, we obtain a reasonable closure of the energy cycle (Fig. 5), where the total of CA and GE is very close to CE, the conversion to eddy kinetic energy ( $\text{KIN}_E$ ).


While the direct impact of diabatic heating by GE ( $0.44 \times 10^5 \text{ J m}^{-2} \text{ d}^{-1}$ ) is only  $\approx 15\%$  of CA, the diabatic heating also alters the temperature anomaly and thus contributes indirectly to CA. Thus, the ratio between the hemispherically averaged GE and CA cannot directly suggest a relative impact of moisture for the storm track activity. Nevertheless, given the systematically organised eddy moisture flux convergence (Fig. 3) as well as the corresponding diabatic heating (Figs. 4), it is worthwhile to discuss the meridional distributions of the vertically averaged energy budget to obtain an indication of the local impact of moisture in the energy budget.

Consistent with Ogawa et al. (2012), the peak latitude of CA is sensitive to the position of the SST front (Fig. 6a). While CE, GE, and  $\text{KIN}_E$  increase for an equatorward shift in the SST front (Fig. 6b–d), the maximum amplitude of CA remains largely unchanged for the latitudinal shift in the SST front (Fig. 6a). The sensitivity of GE is consistent with higher SST yielding enhanced diabatic heating around the SST front in the mid-troposphere (red lines in Fig. 2). The reduction in GE (Fig. 6c) along the equatorward flank of the SST front is related to surface heat fluxes dampening near-surface atmospheric temperature anomalies. The enhancement of GE on the poleward flank of the SST front, on the other hand, is due to mid-tropospheric diabatic heating associated with the eddy moisture flux convergence (Fig. 3). In fact, the double peak structure in GE (Fig. 6c) in the winter hemisphere is hinted at by  $\text{KIN}_E$  (Fig. 6d), suggesting a considerable impact of diabatic processes on  $\text{KIN}_E$ . CA supplying  $\text{APE}_E$  parallel to CE (Fig. 5) also shows a similar double peak (Fig. 6a), suggesting a diabatic contribution to the temperature anomaly in CA (Eq. 16). CA, CE, GE, and  $\text{KIN}_E$  are all amplified in the presence of the SST front compared to REF (Fig. 6a–d), confirming the role of the SST front in intensifying the transient eddy activity (Sampe et al., 2010).





**Figure 6.** (a) CA, (b) CE, (c) GE, and (d)  $KIN_E$  averaged zonally and vertically from 850 to 200 hPa for F35 (red), F45 (black), F55 (blue), and REF (grey).




**Figure 7.** As Fig.2, but for (a) F45+5, (b) REF+5, (c) F45–5, and (d) REF–5.

The lower-latitude peaks in CA, GE, CE, and  $KIN_E$  are always located at a similar latitude near  $33^\circ$  in the winter hemisphere for F45, F55, and REF, where GE in REF is as strong as F45 and F55 (Fig. 6). This is consistent with the eddy moisture flux convergence peaking at a similar latitude (Fig. 2b–d). These similarities of the latitudinal peaks in both storm track activity and eddy moisture flux convergence are consistent with the existence of a secondary maximum in the

SST gradient around  $33^\circ$  for F45, F55, and REF (Fig. 1b). For F35, this weak subtropical SST gradient is joined with the main SST front at  $35^\circ$ , and the storm track activity and water cycle are maximised at the poleward flank of the main SST front.

To disentangle the impact between SST and SST gradient, we repeat F45 and REF with a uniform increase or decrease in SST by 5 K, referred to as F45+5, F45–5, REF+5, and



**Figure 8.** As Fig. 6, but for F45+5 (orange), F45 (black), and F45–5 (green).

REF–5, respectively. The vertically integrated water cycle amplifies (weakens) for higher (lower) SST, without a systematical shift in latitudinal position (Fig. 7). The peak of the moisture flux convergence on the poleward flank of the SST front in F45 is amplified (reduced) by  $\approx 50\%$  for F45+5 (F45–5). This is consistent with a change in the mean moisture flux convergence in conjunction with the strength of the Hadley circulation, balanced by convective condensation, as well as changes in eddy moisture flux convergence together with the steepness of isentropic surfaces, balanced by microphysics condensation (Figs. S3–6).

The uniform changes in SST also modify the storm track activity, yielding enhanced (reduced) storm track activity for higher (lower) SST ( $KIN_E$ ; Fig. 8d). Small changes in CA (Fig. 8a) for the uniform SST change can be understood by the unchanged SST gradient for these experiments. In contrast, GE and CE show a clear increase (decrease) for higher (lower) SST ( $KIN_E$ ; Fig. 8b, c), consistent with changes in  $KIN_E$  (Fig. 8d). The sensitivity of storm track activity to uniform changes in SST (Fig. 8) can hence also explain the sensitivity of the storm track activity to the position of the SST front (Fig. 6), as the mean SST is higher (lower) for F35 (F55).

## 9 Conclusions

Using a set of idealised aqua-planet experiments, we investigated the sensitivity of the atmospheric water cycle and storm track activity to shifting or removing the mid-latitude SST front. When shifting the latitude of the SST front, the tropical circulation and atmospheric water cycle are almost unaf-

ected, with barely any changes in the time-mean circulation and Hadley cell. In the mid-latitudes, however, evaporation and convective and large-scale precipitation, as well as the eddy moisture flux and its convergence, are shifting latitudinally in concert with the SST front.

The atmospheric water budget in the tropical latitudes is dominated by evaporation from the warm ocean surface, providing moisture to the boundary layer which is then transported upward by the convection parameterisation in the subtropics. In the inner tropics, the time-mean circulation associated with the Hadley cell provides upward moisture transport and convergence, where the moisture is released as convective precipitation along the ITCZ. In the mid-latitudes, on the other hand, upward surface latent heat fluxes and convective precipitation are maximised along the equatorward flank of the SST front, whereas large-scale precipitation and atmospheric eddy moisture flux convergence are maximised on the poleward flank of the SST front. The absence of these features in a simulation without an SST front confirms the importance of the SST front for the atmospheric water cycle in the mid-latitudes.

As storm tracks are self-maintained through the diabatic generation of eddy available potential energy (GE) associated with the eddy moisture flux convergence ( $C_e$ ) and microphysics condensation ( $\dot{q}_m$ ), the position of the storm track is diabatically anchored at the SST front. This is consistent with eddy kinetic energy ( $KIN_E$ ) and microphysics condensation ( $\dot{q}_m$ ) being weaker and less confined for the experiment without an SST front, indicating that the SST front latitudinally constrains and thereby enhances the storm track activity.

Given that surface latent heat fluxes are largely determined by the surface saturation mixing ratio at SST through the Clausius–Clapeyron relation, the distributions of evaporation and convective precipitation are strongly tied to the SST. Hence, moving the SST front to higher (lower) latitudes at relatively lower (higher) SST reduces (increases) the strength of the water cycle and thereby the intensity of the storm track. This sensitivity is confirmed through experiments with globally increased or reduced SST, with the water cycle and storm track intensity changing by 50 % for SST changes of  $\pm 5$  K.

Our results demonstrate the importance of both SST and SST fronts to understand the mid-latitude atmospheric water cycle and storm track activity. While the position of the SST front determines the position of the eddy moisture convergence and thus the diabatic heating that energises the storm track, the underlying SST determines the general strength of the water cycle and thus the intensity of the storm track. Our study thus highlights the importance of correctly representing SST and the position of SST fronts to ensure an adequate representation of both the atmospheric water cycle and the storm track.

*Code and data availability.* The data used in this study are publicly available from the NIRD research data archive: <https://doi.org/10.11582/2020.00065> (Ogawa and Spengler, 2020). Figures shown in this study are plotted using the NCAR Command Language (NCL, <https://doi.org/10.5065/D6WD3XH5>, The NCAR Command Language, 2017). Codes can be obtained from the corresponding author.

*Supplement.* The supplement related to this article is available online at: <https://doi.org/10.5194/wcd-5-1031-2024-supplement>.

*Author contributions.* FO and TS are the co-initiators of this study; FO conducted the analysis and was the first author of the main text; TS guided the assembly of the study and helped revise the paper.

*Competing interests.* The contact author has declared that neither of the authors has any competing interests.

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

*Acknowledgements.* Fumiaki Ogawa was supported by MEXT; Grant-in-Aid programs 6102 (Hotspot 2), 24A203 (Hotspot 3), and JPMXD1420318865 (ArCS II); by the Ministry of Environment of Japan; the Environment Research and Technology Development

Fund (grant no. JPMEERF20242001); and by the Research Council of Norway (grant no. 275268) (COLUMBIA). Thomas Spengler was supported by the project BALMCAST (Research Council of Norway project number 324081). This study was supported by UNPACC (Research Council of Norway project number 262220). We also thank Ophélie Nourrisson and Florian Marris for some initial work on this topic. Computing resources were provided by the Japan Agency for Marine Science and Technology (JAMSTEC).

*Financial support.* This research has been supported by the Ministry of Education, Culture, Sports, Science and Technology (grant nos. 6102, 24A203, JPMXD1420318865); the Ministry of the Environment of Japan (grant no. JPMEERF20242001); and the Research Council of Norway (grant nos. 324081, 262220, and 275268).

*Review statement.* This paper was edited by David Battisti and reviewed by one anonymous referee.

## References

- Bryan, F. and Oort, A.: Season Variation of the Global Water Balance Based on Aerological Data, *J. Geophys. Res.*, 89, 11717–11730, 1984.
- Bui, H. and Spengler, T.: On the Influence of Sea Surface Temperature distributions on the Development of Extratropical Cyclones, *J. Atmos. Sci.*, 78, 1173–1188, <https://doi.org/10.1175/JAS-D-20-0137.1>, 2021.
- Deremble, B., Lapeyre, G., and Ghil, M.: Atmospheric Dynamics Triggered by an Oceanic SST Front in a Moist Quasigeostrophic Model, *J. Atmos. Sci.*, 69, 1617–1632, 2012.
- Dey, K. and Döös, K.: The coupled ocean–atmosphere hydrologic cycle, *Tellus A*, 71, 1650413, <https://doi.org/10.1080/16000870.2019.1650413>, 2019.
- Emanuel, K. A.: A Scheme for Representing Cumulus Convection in Large-Scale Models, *J. Atmos. Sci.*, 48, 2313–2329, [https://doi.org/10.1175/1520-0469\(1991\)048<2313:ASFRCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1991)048<2313:ASFRCC>2.0.CO;2), 1991.
- Enomoto, T., Kuwano-Yoshida, A., Komori, N., and Ohfuchi, W.: Description of AFES 2: Improvements for high-resolution and coupled simulations., *High Resolution Numerical Modelling of the Atmosphere and Ocean*, edited by: Kevin, H. and Wataru, O., Springer, 77–97, [https://doi.org/10.1007/978-0-387-49791-4\\_5](https://doi.org/10.1007/978-0-387-49791-4_5), 2008.
- Hartmann, D.: *Global Physical Climatology*, Elsevier Science, 2nd Edn., <https://doi.org/10.1016/C2009-0-00030-0>, 2016.
- Haualand, K. F. and Spengler, T.: Direct and Indirect Effects of Surface Fluxes on Moist Baroclinic Development in an Idealized Framework, *J. Atmos. Sci.*, 77, 3211–3225, <https://doi.org/10.1175/JAS-D-19-0328.1>, 2020.
- Höjgård-Olsen, E., Brogniez, H., and Chepfer, H.: Observed Evolution of the Tropical Atmospheric Water Cycle with Sea Surface Temperature, *J. Climate*, 33, 3449–3470, 2020.
- Hoskins, B. J. and Valdes, P. J.: On the Existence of Storm Tracks, *J. Atmos. Sci.*, 47, 1854–1864, 1990.
- Hotta, D. and Nakamura, H.: On the significance of the sensible heat supply from the ocean in the maintenance of the mean

- baroclinicity along storm tracks, *J. Climate*, 24, 3377–3401, <https://doi.org/10.1175/2010JCLI3910.1>, 2011.
- Kuwano-Yoshida, A., Enomoto, T., and Ohfuchi, W.: An improved PDF cloud scheme for climate simulations., *Q. J. Roy. Meteor. Soc.*, 136, 1583–1597, 2010.
- Le Trent, H. and Li, Z.-X.: Sensitivity of an atmospheric general circulation model to prescribed SST changes: feedback effects associated with the simulation of cloud optical properties, *Clim. Dynam.*, 5, 175–187, <https://doi.org/10.1007/BF00251808>, 1991.
- Lorenz, E. N.: Available Potential Energy and the Maintenance of the General Circulation, *Tellus*, 7, 157–167, <https://doi.org/10.3402/tellusa.v7i2.8796>, 1955.
- Madonna, E., Wernli, H., Joos, H., and Martius, O.: Warm Conveyor Belts in the ERA-Interim Dataset (1979–2010). Part I: Climatology and Potential Vorticity Evolution., *J. Climate*, 27, 3–26, 2014.
- McTaggart-Cowan, R., Gyakum, J. R., and Moore, R. W.: The Baroclinic Moisture Flux, *Mon. Weather Rev.*, 145, 25–47, <https://doi.org/10.1175/MWR-D-16-0153.1>, 2017.
- Nakamura, H., Sampe, T., Tanimoto, Y., and Shimpo, A.: Observed Associations Among Storm Tracks, Jet Streams and Midlatitude Oceanic Fronts, *AGU Monograph*, 147, 329–345, 2004.
- Nakamura, H., Sampe, T., Goto, A., Ohfuchi, W., and Xie, S.-P.: On the importance of midlatitude oceanic frontal zones for the mean state and dominant variability in the tropospheric circulation, *Geophys. Res. Lett.*, 35, L15709, <https://doi.org/10.1029/2008GL034010>, 2008.
- Neiman, P. J., Ralph, F. M., Wick, G. A., Lundquist, J., and Dettinger, M. D.: Meteorological characteristics and overland precipitation impacts of atmospheric rivers affecting the west coast of North America based on eight years of SSM/I satellite observations, *J. Hydrometeorol.*, 9, 22–47, <https://doi.org/10.1175/2007JHM855.1>, 2008.
- Newman, M., Kiladis, G. N., Weickmann, K. M., Ralph, F. M., and Sardeshmukh, P. D.: Relative Contributions of Synoptic and Low-Frequency Eddies to Time-Mean Atmospheric Moisture Transport, Including the Role of Atmospheric Rivers, *J. Climate*, 25, 7341–7361, 2012.
- Ogawa, F. and Spengler, T.: Prevailing Surface Wind Direction during Air–Sea Heat Exchange, *J. Climate*, 32, 5601–5617, 2019.
- Ogawa, F. and Spengler, T.: Output of sensitivity experiments by aqua-planet AGCM on the latitude and existence of SST front, NIRD Research data archive [data set], <https://doi.org/10.11582/2020.00065>, 2020.
- Ogawa, F., Nakamura, H., Nishii, K., Miyasaka, T., and Kuwano-Yoshida, A.: Dependence of the climatological axial latitudes of the tropospheric westerlies and storm tracks on the latitude of an extratropical oceanic front, *Geophys. Res. Lett.*, 35, L15709, <https://doi.org/10.1029/2011GL049922>, 2012.
- Ogawa, F., Omrani, N.-E., Nishii, K., Nakamura, H., and Keenlyside, N.: Ozone-induced climate change propped up by the Southern Hemisphere oceanic front, *Geophys. Res. Lett.*, 42, 10056–10063, 2015.
- Ogawa, F., Nakamura, H., Nishii, K., Miyasaka, T., and Kuwano-Yoshida, A.: Importance of midlatitude oceanic frontal zones for the annular mode variability: Interbasin differences in the southern annular mode signature, *J. Climate*, 29, 6179–6199, <https://doi.org/10.1175/JCLI-D-15-0885.1>, 2016.
- Ohfuchi, W., Nakamura, H., Yoshioka, M., Enomoto, T., Takaya, K., Peng, X., Yamane, S., Nishimura, T., Kurihara, Y., and Ni-nomiya, K.: 10-km mesh meso-scale resolving global simulation of the atmosphere on the Earth Simulator – Preliminary outcome of AFES (AGCM for the Earth Simulator), *J. Earth Simulator*, 1, 8–34, 2004.
- Oki, T., Entekhabi, D., and Harrold, T. I.: The Global Water Cycle, vol. 150 of *Geophysical Monograph Series*, AGU, <https://doi.org/10.1029/150GM18>, 2004.
- Oort, A. H. and Peixoto, J. P.: The Annual Cycle of the Energetics of the Atmosphere on a Planetary Scale, *J. Geophys. Res.*, 79, 2705–2719, 1974.
- Papritz, L. and Spengler, T.: Analysis of the slope of isentropic surfaces and its tendencies over the North Atlantic, *Q. J. Roy. Meteor. Soc.*, 141, 3226–3238, 2015.
- Peixoto, J. P. and Oort, A. H.: The Annual Distribution of Atmospheric Energy on a Planetary Scale, *J. Geophys. Res.*, 79, 2149–2159, 1974.
- Peixoto, J. P. and Oort, A. H.: *Physics of Climate*, American Institute of Physics, 1st Edn., ISBN 978-0-88318-712-8, 1992.
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily high-resolution-blended analyses for sea surface temperature, *J. Climate*, 20, 5473–5496, 2007.
- Sampe, T., Nakamura, H., Goto, A., and Ohfuchi, W.: Significance of a midlatitude SST frontal zone in the formation of a storm track and an eddy-driven westerly jet, *J. Climate*, 23, 1793–1814, 2010.
- Sampe, T., Nakamura, H., and Goto, A.: Potential Influence of a Midlatitude Oceanic Frontal Zone on the Annular Variability in the Extratropical Atmosphere as Revealed by Aqua-Planet Experiments, *J. Meteorol. Soc. Jpn.*, 91, 243–267, 2013.
- Schneider, T., O’Gorman, P. A., and Levine, Z. J.: Water vapor and the dynamics of climate changes, *Rev. Geophys.*, 48, RG3001, <https://doi.org/10.1029/2009RG000302>, 2010.
- Seager, R., Naik, N., and Vecchi, G. A.: Thermodynamic and Dynamic Mechanisms for Large-Scale Changes in the Hydrological Cycle in Response to Global Warming, *J. Climate*, 23, 4651–4668, 2010.
- The NCAR Command Language: (Version 6.4.0), Boulder, Colorado: UCAR/NCAR/CISL/VETS [software], <https://doi.org/10.5065/D6WD3XH5>, 2017.
- Tilina, N., Gavrikov, A., and Gulev, S. K.: Association of the North Atlantic surface turbulent heat fluxes with midlatitude cyclones, *Mon. Weather Rev.*, 146, 3691–3715, 2018.
- Trenberth, K.: Changes in precipitation with climate change, *Clim. Res.*, 47, 123–138, 2011.
- Trenberth, K., Smith, L., Qian, T., Dai, A., and Fasullo, J.: Estimates of the Global Water Budget and Its Annual Cycle Using Observational and Model Data, *J. Hydrometeorol.*, 8, 758–769, 2007.
- Watt-Meyer, O. and Frierson, D. M.: ITCZ Width Controls on Hadley Cell Extent and Eddy-Driven Jet Position and Their Response to Warming, *J. Climate*, 32, 1151–1166, 2019.
- Zhu, Y. and Newell, R. E.: A proposed algorithm for moisture fluxes from atmospheric rivers, *Mon. Weather Rev.*, 126, 725–735, 1998.
- Zolina, O. and Gulev, S. K.: Synoptic Variability of Ocean–Atmosphere Turbulent Fluxes Associated with Atmospheric Cyclones, *J. Climate*, 16, 2717–2734, 2003.