



# From sea to sky: understanding the sea surface temperature impact on an atmospheric blocking event using sensitivity experiments with the ICOSahedral Nonhydrostatic (ICON) model

Svenja Christ<sup>1</sup>, Marta Wenta<sup>1</sup>, Christian M. Grams<sup>1,a</sup>, and Annika Oertel<sup>1</sup>

<sup>1</sup>Institute of Meteorology and Climate Research Troposphere Research (IMKTRO),  
Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

<sup>a</sup>now at: Federal Office of Meteorology and Climatology, MeteoSwiss, Zurich, Switzerland

**Correspondence:** Svenja Christ (svenja.christ@kit.edu)

Received: 30 July 2024 – Discussion started: 19 August 2024

Revised: 29 October 2024 – Accepted: 30 October 2024 – Published: 8 January 2025

**Abstract.** Blocked weather regimes are an important phenomenon in the Euro-Atlantic region and are frequently linked to extreme weather events. Despite their importance for surface weather, the correct prediction of blocking events remains challenging. Previous studies indicated a link between the misrepresentation of blocking events in numerical weather prediction models and sea surface temperature (SST) biases, particularly in the Gulf Stream region. However, the pathway that links SST in the Gulf Stream region and the downstream upper-level flow is not yet fully understood. To deepen our physical understanding of the link between the Gulf Stream SST and downstream atmospheric blocking, we perform sensitivity experiments with varying SST conditions for an atmospheric blocking event in February 2019. This blocking event, which was associated with a winter heat wave with unprecedented temperatures in western Europe, was both preceded and accompanied by several rapidly intensifying extratropical cyclones originating in the Gulf Stream region and crossing the North Atlantic. Those cyclones and their associated rapidly ascending air streams, so-called warm conveyor belts (WCBs), played a crucial role in the development of the upper-level ridge and the blocking event. The ascent of these WCBs, which connect the lower and upper troposphere, was enhanced by moisture uptake during cold air outbreaks (CAOs) in the Gulf Stream region. In this study, we employ sensitivity experiments with the ICOSahedral Nonhydrostatic Weather and Climate Model (ICON) to assess the impact of intense air–sea interactions during CAOs on WCBs and the downstream ridge. In to-

tal five different experiments are used, including idealized and weakened SST gradients and one with increased absolute SST in the Gulf Stream region. Using Eulerian and Lagrangian perspectives, we demonstrate that the SST gradient in the Gulf Stream region affects moisture availability and air temperature in the WCB inflow region and, consequently, WCB ascent. In our case study, stronger SST gradients lead to increased specific humidity and warmer temperatures in the lower troposphere, resulting in more pronounced WCB ascent, while weaker SST gradients are associated with reduced WCB activity. The differences in WCB ascent and outflow properties induced by weakened SST gradients, such as reduced cross-isentropic ascent and outflow heights, subsequently influence the upper-level flow and weaken the downstream ridge. Moreover, experiments with weaker SST gradients show a decrease in cyclone intensity, and vice versa, stronger cyclones are found in experiments with warmer SSTs. To summarize, our results suggest that different SST and SST gradient representations affect the large-scale atmospheric flow via the WCB airstream. Specifically, moisture availability regulated by SST and SST gradients in the WCB inflow region influences subsequent WCB ascent and outflow characteristics, which, in turn, influence the upper-level ridge downstream. The SST in the Gulf Stream region affects WCB characteristics consistently from the inflow, over the ascent to the outflow phase.

## 1 Introduction

Blocking regimes form a persistent, quasi-stationary atmospheric state that interrupts the eastward propagation of mid-latitude weather systems (Michelangeli et al., 1995; Teubler and Riemer, 2016). They are often linked to extreme weather events (e.g., Yiou and Nogaj, 2004; Booth et al., 2017; Schaller et al., 2018; Spensberger et al., 2020; Kautz et al., 2022), as the associated persistent anticyclonic circulation can dominate the weather at a particular location for several days to weeks (Wazneh et al., 2021). To date, the correct prediction of blocking events and associated surface weather remains challenging (Grams et al., 2018; Büeler et al., 2021; Oertel et al., 2023b). Previous research linked the Gulf Stream sea surface temperature (SST) with downstream blocking events in the North Atlantic and western Europe (e.g., O'Reilly et al., 2016; Yamamoto et al., 2021) and suggested that SST biases in the North Atlantic region might contribute to challenges in predicting blocking downstream (Czaja et al., 2019; Roberts et al., 2021; Athanasiadis et al., 2022). However, the physical pathway between SST in the Gulf Stream region and atmospheric blocking events is still not fully understood. In their seminal work Pfahl et al. (2015) showed that latent heat release in ascending air streams is a first-order process in the development of blocking anticyclones. Most of this ascent is confined to the warm conveyor belt (WCB) airstream associated with extratropical cyclones (e.g., Madonna et al., 2014), which ascends into the upper troposphere where it can amplify upper-level ridges, subsequently resulting in blocking (e.g., Grams and Archambault, 2016). WCB airstreams occur frequently in the mid-latitudes and accompany approximately 60% of extratropical cyclones (Carlson, 1980; Eckhardt et al., 2004). The western North Atlantic, particularly the Gulf Stream region, is one of the hotspots for WCB development (Eckhardt et al., 2004; Madonna et al., 2014). WCBs originate in the lower troposphere in the warm sector of extratropical cyclones and typically ascend poleward near the surface cold front (Wernli, 1997). SST variability can affect air–sea interactions and thus modulate low-level moisture in the WCB inflow region, which subsequently affects the associated WCB ascent and outflow characteristics in the upper troposphere (e.g., Schäfler and Harnisch, 2015; Yamamoto et al., 2021; Wenta et al., 2024). In this study, we thus investigate the hypothesis that the WCB airstream links SST variations in the Gulf Stream region to downstream blocking events. In the following, we provide a brief review of previous studies investigating the influence of SST on cyclone dynamics, cold air outbreaks (CAOs), and the impact of WCBs on blocking events. Furthermore, we summarize the impact of moisture availability on WCBs and diabatic heating.

Extratropical cyclones in the North Atlantic and Pacific oceans tend to organize along strong SST gradients (Nakamura et al., 2004). In the North Atlantic, the largest frequency of mid-latitude atmospheric fronts occurs along the

Gulf Stream SST gradient (Berry et al., 2011; Reeder et al., 2021). As a consequence, the influence of SST changes on cyclones is most pronounced in these regions (e.g., Small et al., 2014; Tsopouridis et al., 2021). For example, cyclone intensification is sensitive to latent heat fluxes in their vicinity, which is directly influenced by SST (Vries et al., 2019). Modifying the SST gradient in global atmospheric models has been shown to impact not only the location but also the characteristics of the storm track (Small et al., 2014; Schemm, 2023). The maintenance of the storm track itself is linked to increased low-level baroclinicity (Hotta and Nakamura, 2011; Papritz and Spengler, 2015), which can be sustained in regions like the Gulf Stream due to differential surface heat fluxes (Papritz et al., 2015). On subseasonal-to-seasonal timescales, forecasts could benefit from a higher ocean resolution with improved representation of small ocean eddies (Roberts et al., 2022), while the small-scale SST perturbations may have a relatively small influence on individual synoptic events (Roberts et al., 2021). Nevertheless, Tsopouridis et al. (2021) reported reduced cyclone activity in the North Atlantic as a result of a smoothed SST gradient. In contrast, Bui and Spengler (2021) observed no notable differences in the deepening rates of individual cyclones between smoothed and observed SST conditions. Instead, Bui and Spengler (2021) concluded that the distribution of absolute SSTs, rather than the SST gradient, influences latent heat release and subsequent diabatic cyclone intensification. In line with this, numerical experiments suggest that cyclones respond to a decreasing SST gradient differently, depending on whether the cyclones pass over the warm or cold side of the SST gradient (Booth et al., 2012). Overall, this not only emphasizes the importance of the Gulf Stream for cyclone dynamics in the North Atlantic but also suggests that the influence of absolute SST and SST gradients on the synoptic flow evolution is not yet fully understood.

The passage of cyclones across the Gulf Stream is often linked to the development of CAOs and intense air–sea interactions, driven by the large air–sea temperature differences (e.g., Papritz and Spengler, 2015). The resulting surface heat and moisture fluxes play an important role in regulating the heat and moisture supply to the rapidly ascending air streams (WCBs; Booth et al., 2017). Warmer SSTs enforce more latent heat fluxes and potentially enhance latent heat release and cyclone intensification (Booth et al., 2012; Bui and Spengler, 2021), whereby more intense cyclones tend to have stronger WCBs (Binder et al., 2016). Due to its deep ascent, the WCB airstream connects the boundary layer and the upper troposphere (Wernli, 1997). The ascent of the WCB is characterized by substantial latent heat release (Browning, 1990), and cloud bands forming during WCB ascent can reach lengths of up to 3000 km (Browning et al., 1973). Latent heating enables the cross-isentropic flow of WCB air parcels (Wernli and Davies, 1997; Joos and Wernli, 2012; Madonna et al., 2014). Moreover, the latent heating pattern along the ascent is linked to the characteristic poten-

tial vorticity (PV) changes along WCB trajectories, as PV is produced below the level of maximum latent heat release and reduced above (Wernli and Davies, 1997; Madonna et al., 2014). As a consequence, the WCB performs a net transport of low PV air into the upper troposphere (Wernli and Davies, 1997), where it can contribute to the formation of a negative PV anomaly (Wernli and Davies, 1997; Grams et al., 2011; Joos and Forbes, 2016). Additionally, the divergent outflow of the WCB contributes to the anticyclonic circulation in the upper troposphere (Browning and Roberts, 1994; Steinfeld and Pfahl, 2019). For these reasons, the divergent WCB outflow can play an important role in the maintenance and intensification of the downstream blocking event (Grams et al., 2011; Michel and Rivière, 2011; Pfahl et al., 2015; Grams and Archambault, 2016; Teubler and Riemer, 2016; Steinfeld and Pfahl, 2019; Steinfeld et al., 2020). As WCB ascent substantially influences ridge amplification and blocking intensity, the detailed representation of WCB ascent is important to correctly represent flow properties and minimize forecast error (e.g., Schäfler and Harnisch, 2015; Grams et al., 2018; Berman and Torn, 2019; Steinfeld et al., 2020; Berman and Torn, 2022; Pickl et al., 2023). A key factor for the strengthening of WCB ascent is moisture availability since this can influence subsequent latent heat release and cross-isentropic ascent (Schäfler and Harnisch, 2015; Dacre et al., 2019; Quinting and Grams, 2021; Berman and Torn, 2022; Quinting et al., 2022; Oertel et al., 2023a). Thus, stronger latent heating in response to climate change is expected to influence the size and intensity of blocking anticyclones (Steinfeld et al., 2022). Further, the releases of latent heat can introduce forecast uncertainties (Berman and Torn, 2019, 2022). Therefore, the intensity of the WCB and its impact on PV values in the upper troposphere are closely linked to moisture availability in the lower troposphere (Schäfler and Harnisch, 2015; Schemm et al., 2013).

A primary moisture source of extratropical cyclones is local and located over the western North Atlantic, specifically in the Gulf Stream region (Pfahl et al., 2014; Papritz et al., 2021). In this region, initial cold and dry air from the American continent is heated and moistened by the warm waters south of the SST front during CAOs (Papritz et al., 2021). The advection of cold air across the Gulf Stream leads to intense surface heat fluxes. The moistening of the marine boundary layer is often caused by the passage of a predecessor cyclone, pointing at a possible cyclone-to-cyclone interaction (Sodemann et al., 2008). Specifically, Dacre et al. (2019) and Papritz et al. (2021) demonstrated that the cyclone's moisture typically originates in the cold sector in the pre-cyclone environment of the preceding cyclone. In particular, CAOs caused by the preceding cyclone contribute to this moisture uptake (Wenta et al., 2024).

The described synoptic features are relevant for the synoptic evolution investigated here. In February 2019, an atmospheric blocking event led to the development of a significant winter heat wave in western Europe (Kendon et al., 2020;

Young and Galvin, 2020; Leach et al., 2021). This event was both preceded and accompanied by a series of rapidly intensifying cyclones, leading to the occurrence of intense CAOs in the western and central North Atlantic (Wenta et al., 2024). Wenta et al. (2024) indicated that these cyclones and their associated WCBs may have contributed to the development and maintenance of the block, as they played a key role in forming the upper-level PV anomaly associated with the European blocking event. Furthermore, Wenta et al. (2024) showed that the moisture sources for those cyclones were associated with air–sea interactions during CAOs initiated by the preceding cyclones in the same region, in agreement with the conceptual model proposed by Papritz et al. (2021). These results suggest that cyclones and their associated WCB airstreams provide a mechanistic link of surface processes in the Gulf Stream region with the development of an atmospheric block downstream in February 2019 (see also Kwon et al., 2010; Czaja et al., 2019; Athanasiadis et al., 2022; Wenta et al., 2024).

In this study, we want to explore how sensitive this mechanistic link is to variations of absolute SST and SST gradients in the Gulf Stream region, specifically how absolute SST and SST gradients modulate air–sea interaction, subsequent cyclone development, WCB ascent, and upper-level flow amplification. Therefore, we conduct numerical single-member deterministic sensitivity experiments using the ICON (ICOsahedral Nonhydrostatic) model and address the following main research questions:

1. How do SST gradient and absolute SST perturbations in the Gulf Stream region influence air–sea interactions?
2. How do changes in low-level moisture availability from SST perturbations influence the ascending WCB airstream linking the lower and upper troposphere?
3. Do air–sea interactions over the Gulf Stream region influence upper-level ridge amplification and the formation of the European blocking event through their impact on WCB ascent associated with extratropical cyclones?

In the following section, we provide a detailed description of the methodology (Sect. 2). This is followed by a more detailed introduction to the February 2019 case study (Sect. 3) and the presentation of the results (Sect. 4). We finally conclude with a discussion and summary of the results (Sect. 5).

## 2 Methods

To examine the impact of SST in the Gulf Stream region on WCB ascent and the subsequent development of a downstream ridge, we conduct a detailed case study of a European blocking event (see Sect. 3 for details). The blocking event in February 2019 was preceded and accompanied by a series of rapidly intensifying cyclones, each associated with a

WCB. To determine how variations in SSTs and SST gradients in the Gulf Stream region influence WCB characteristics, numerical simulations with the ICON model (Zängl et al., 2015) were performed, featuring five different prescribed SST configurations in the eastern North Atlantic (see Sect. 2.2 for details).

## 2.1 ICON model setup

The single-member ICON simulations (version 2.6.2.2) are run freely and globally for a lead time of 9 d. The simulations are run with a horizontal resolution of about 13 km (R3B07 grid), along with 90 vertical model levels and a time step of 120 s. This setup corresponds to the operational resolution of the global ICON model used by the German Weather Service. The ICON simulations are initialized from ECMWF's IFS (Integrated Forecasting System) analysis at 00:00 UTC on 18 February 2019. Cloud microphysical processes in the model are represented using a single-moment scheme (Seifert, 2008; Doms et al., 2018), which includes four prognostic hydrometeor categories: cloud water, rain, ice, and snow. The model employs a Tiedtke–Bechtold bulk mass flux scheme for parameterizing convection (Tiedtke, 1989; Bechtold et al., 2008), and the radiation calculations are performed on a reduced radiation grid (R3B06 grid) utilizing the ecRAD scheme (Hogan and Bozzo, 2018). Additionally, the default ICON schemes are applied for subgrid-scale orographic drag (Lott and Miller, 1997), non-orographic gravity wave drag (Orr et al., 2010), and turbulence (Raschendorfer, 2018). Surface fluxes are parameterized using the drag law formulation (Raschendorfer, 2018), which takes into account the horizontal velocity at the lowest model level, the bulk aerodynamic transfer coefficient for turbulent heat exchange at the surface, and the surface temperature.

## 2.2 Design of sensitivity experiments

In total, five separate sensitivity experiments with modified SSTs are conducted, to address the research questions. The reference SST is derived from the ECMWF IFS analysis data, which uses the SST from the Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system (Good et al., 2020). The OSTIA product uses satellite and in situ data from infrared and microwave radiometers (Good et al., 2020). This SST analysis includes large SST gradients associated with the Gulf Stream (Fig. 1a). Throughout each of the 9 d long simulations, the prescribed SSTs remain constant. The five experiments are characterized by the following modifications to the reference SST pattern:

### 1. Control experiment (CNTRL)

The simulation is initialized with SSTs from the ECMWF IFS analysis (Fig. 1a), which has been remapped to the ICON grid. Due to the relatively high resolution of around 9 km, smaller-scale meanders and locally large SST gradients are present.

### 2. Idealized SST gradient experiment (IDEA)

For IDEA, the SST front's small-scale eddies are removed (Fig. 1b) to create an idealized and smooth SST gradient. To achieve this, we apply a Gaussian filter to the two-dimensional SST field in the Gulf Stream region (30 to 55° N and 80 to 25° W). This filter uses a Gaussian kernel with a standard deviation of 3, applied uniformly in both directions. To prevent artificially strong temperature gradients at the border regions of the modified SST field, we included additional smoothing at the border region. As IDEA exhibits only minor differences compared to CNTRL, providing nearly identical results (see Sect. 4), this experiment serves as a reference throughout the study.

### 3. Weak SST gradient experiment (WEAK)

For WEAK (Fig. 1c), a stronger smoothing than for IDEA was applied with a standard deviation of 12 for the Gaussian kernel applied in the Gulf Stream region. This results in a substantially weakened SST gradient at the Gulf Stream SST front.


### 4. Extra weak SST gradient experiment (extWEAK)

In the extWEAK simulation, an even stronger smoothing with a standard deviation of 36 for the Gaussian kernel essentially removed the distinct SST front (Fig. 1e). Moreover, SST was altered in a broader area in the North Atlantic to prevent artificially high SST gradients at border regions. In this setup, the SST transitions gradually and cools progressively from the Equator towards the higher latitudes.

### 5. Warmed SST experiment (P1.5K)

Besides the experiments involving modified SST gradients, the P1.5K experiment features an increase in SST by up to 1.5 K (Fig. 1d). The warming was applied to the IDEA reference experiment. Specifically, SST is increased by 1.5 K in an ellipse centered around the Gulf Stream SST gradient at 41.5° N and 62.5° W. The applied warming gradually decreases towards domain boundaries to prevent the generation of artificial gradients elsewhere in the North Atlantic. In effect, the applied warming is limited to 35 to 55° N and 40 to 80° W.

The SST modifications slightly change the area-mean SST in the region of interest located from 30 to 55° N and from 80 to 25° W (Table 1). This region is of particular interest as most of the cyclones associated with the block pass through, and often intensify, in this region. As expected, CNTRL and IDEA show almost no discernible difference in area-averaged SST, with the only alteration between those experiments being the removal of the small-scale meanders in the Gulf Stream in IDEA. Per definition, P1.5K has the warmest area-averaged SST of 287.27 K (Table 1) and higher



**Figure 1.** Sea surface temperature (SST, shading, in K) and SST gradient (contours, at 0.01, 0.05, and 0.1 K km<sup>-1</sup> colored in grey and black) for the 9 d simulation period for the experiments (a) CNTRL, (b) IDEA, (c) WEAK, (d) extWEAK, and (e) P1.5K. The Gulf Stream region referred to in the text is outlined by the blue box (30 to 55° N and 80 to 25° W).

SSTs in the entire Gulf Stream region with maximum differences reaching up to 1.5 K compared to IDEA. In the WEAK and extWEAK simulations, the SSTs are warmer north of the SST front and colder to the south in comparison to IDEA. As the area south of the SST front in the region of interest is larger, the average SST in the Gulf Stream region is effectively lower in both the WEAK and extWEAK simulations. The coldest area-averaged SST of 285.12 K is found in extWEAK (Table 1).

### 2.3 Warm conveyor belt trajectories

We employ the Lagrangian perspective to identify the WCB airstream as a coherent ensemble of strongly ascending trajectories (Wernli, 1997; Madonna et al., 2014). Specifically, 48 h forward trajectories are computed every hour between 18 and 25 February 2019 with LAGRANTO (Sprenger and Wernli, 2015). Trajectories are started from a 50 km equidistant horizontal grid spanning the North Atlantic region from 80 to 25° W and from 30 to 55° N. They are initiated from 10 chosen vertical levels within the lowermost 2 km, specifically at altitudes of 50, 200, 400, 600, 800, 1000, 1250, 1500, 1750, and 2000 m. The following variables are traced along the trajectories: pressure height, temperature, specific humidity, and potential vorticity.

From all 48 h trajectories, WCB trajectories are selected as those that ascend at least 500 hPa. To avoid double-counting of trajectories, a filter following the approach of Madonna et al. (2014) is applied. In total, approximately 10 % of all 40 710 trajectories are identified as WCB trajectories. We categorize WCB trajectories into three distinct stages based on their pressure ( $p$ ; e.g., Schäfler et al., 2014; Quinting et al., 2022): lower-tropospheric inflow region ( $p \geq 800$  hPa), ascent region ( $800 \text{ hPa} > p > 400$  hPa), and the outflow region in the upper troposphere ( $p \leq 400$  hPa). Using this stratification, we regrid the positions of WCB air parcels onto the Eulerian grid, resulting in consistent hourly masks for WCB inflow, ascent, and outflow, respectively. Furthermore, we apply the nearest neighbor method, as described by Škerlak et al. (2014), to also regrid potential temperature, specific humidity, and potential vorticity traced along the WCB trajectories for inflow, ascent, and outflow, respectively.

### 3 Case study introduction

In the following, we introduce the 9 d case study characterized by the passage of multiple cyclones across the Gulf Stream region and downstream ridge amplification and discuss its representation in the CNTRL simulation. In Febru-

**Table 1.** Averaged SSTs (mean and standard deviation) in the Gulf Stream region (30 to 55° N and 80 to 25° W; see blue box in Fig. 1) for the five different experiments.

	SST	Characteristics
CNTRL	286.38 ± 7.42 K	SST is taken from the IFS analysis, which uses the OSTIA SST analysis (Good et al., 2020).
IDEA	286.34 ± 7.04 K	Small-scale meanders are removed; local SST differences prevail.
WEAK	285.98 ± 5.64 K	SST gradient is smoothed; cooler SSTs south of the SST front and warmer SSTs north of the SST front occur.
extWEAK	285.12 ± 3.57 K	Strong smoothing removes the SST front; substantially cooler SSTs south of the SST front and warmer SSTs north of the SST front prevail.
P1.5K	287.27 ± 6.99 K	SST is increased by up to 1.5 K around the Gulf Stream front.

ary 2019, Great Britain and western Europe encountered unprecedented warmth, with temperatures reaching 10 to 15 K above the climatological average (Kendon et al., 2020). The exceptional temperature anomalies over Europe in February 2019 were caused by a combination of various processes. Crucially, the development of an upper-level ridge extending from northwestern Africa to southern Scandinavia, along with anticyclonic circulation, enabled the southward movement of warm, maritime air masses into western Europe (Young and Galvin, 2020). Spanning 20–27 February, the event is categorized as a European blocking event according to the seven-regime classification (Grams et al., 2017). This is characterized by a positive 500 hPa geopotential height anomaly over the eastern North Atlantic and a concurrent negative anomaly centered over Greenland. The atmospheric blocking was preceded and accompanied by the development of several rapidly intensifying cyclones. Wenta et al. (2024) suggested that those cyclones originating from the North Atlantic provided conditions for both the moistening of air masses over the Gulf Stream and their ascent into the upper troposphere, potentially contributing to the maintenance and amplification of the block.

Our analysis focuses on the synoptic conditions during the evolution of the two most intensive cyclones as well as the subsequent amplification of the upper-level ridge and the blocking onset, which are covered by the 9 d simulation period.

In the following, the synoptic evolution is presented using the results from the ICON CNTRL simulation. Between 18:00 UTC on 18 February 2019 and 00:00 UTC on 20 February 2019, the first cyclone (CY1) moved along the SST gradient in the Gulf Stream region, beginning its intensification at 00:00 UTC on 19 February (Fig. 2a, c). The cyclone then progressed northward by 00:00 UTC on 20 February. During CY1's passage across the North Atlantic, cold continental air was advected over the warm waters south of the Gulf Stream, triggering a CAO (Fig. 2b). This CAO over the warm Gulf Stream waters resulted in significant heat and moisture exchange between the ocean and the atmosphere

(see also Wenta et al., 2024). The WCB trajectories show that the ascent associated with this blocking event, especially at 12:00 UTC on 19 February, occurred in the central North Atlantic, with an inflow from the western part of the North Atlantic. The outflow of WCB trajectories associated with CY1 (Fig. 2c) was located over the eastern North Atlantic, in the developing ridge, as indicated by the PV distribution at 315 K (Fig. 2c).

On 21 February, a second cyclone (CY2) intensified rapidly in the Gulf Stream region (Fig. 2d–f). Over a period of 2 d, as it crossed the North Atlantic, it significantly strengthened and triggered another CAO event (Fig. 2e). CY2 moved into the area where CY1 had previously undergone rapid intensification, impacting the atmospheric boundary layer through moistening and heating. This sequential pattern of CY2 following CY1's track suggests a cyclone–cyclone preconditioning mechanism, during which moisture from the region behind cyclone CY1 is fed into the warm sector of the subsequent cyclone and its associated WCB (Fig. 2f; Papritz et al., 2021; Demirdjian et al., 2023). Indeed, Wenta et al. (2024) demonstrated that a significant portion of the moisture in the ascending WCB airstreams of the February 2019 cyclones originates locally and is linked to the preceding cyclone activity. Similar to CY1, the outflow of CY2 is primarily situated within the established ridge over western Europe, with the upstream WCB ascent predominantly taking place in the central North Atlantic (Fig. 2f). On 24 February, the CY2 WCB trajectories further reinforce the stationary ridge and support its eastward expansion (Fig. 2i).

The synoptic evolution in the ICON simulations, specifically CNTRL and IDEA (see Fig. A1 and Christ, 2023, for details), compares well with the ERA5 reanalysis (Hersbach et al., 2020). Overall, the CNTRL and IDEA simulations are very similar, and only small differences in sea level pressure (SLP) and upper-level PV emerge and subsequently grow with increasing lead time. All key synoptic features that lead to the onset of blocking are consistently depicted in both experiments, and no notable initial divergence is observed in any of the experiments with modified SST conditions. In-

stead, differences between the experiments gradually emerge and intensify as the lead time increases. In the next section, we describe the results of our comprehensive analysis of sensitivity experiments, each incorporating different SST configurations in the Gulf Stream region (Sect. 2.2), and focus on the impacts on air–sea interactions (Sect. 4.1), the moistening and heating of the overlying atmosphere (Sect. 4.2), cyclone evolution (Sect. 4.3), WCB ascent (Sect. 4.4), and finally the evolution of the upper-level ridge (Sect. 4.5).

#### 4 The role of SST perturbations in the Gulf Stream region

Our analysis employs both Eulerian and Lagrangian perspectives. The Eulerian perspective focuses on the evolution of atmospheric conditions within the Gulf Stream region during the development of cyclones CY1 and CY2, whereas the Lagrangian perspective focuses on the evolution and properties of WCB trajectories associated with CY1 and CY2 that connect the lower and upper troposphere. As noted above, the differences between IDEA and CNTRL are small, and we find a very similar synoptic evolution in IDEA and CNTRL (see also Fig. A1), which includes cyclone tracks, WCB ascent, and the upper-level flow evolution. Yet within the 9 d of lead time, small differences in SLP and upper-level PV emerge (Fig. A1e, f), indicating a small but noticeable influence of high-resolution SST on the synoptic evolution. IDEA is taken as a reference for the comparison of the sensitivity experiments, and SST impacts are quantified as the difference between IDEA and any other experiment. One reason for taking IDEA as a reference is that SST modifications in the other experiments are based on IDEA, whereby the CNTRL experiment specifically represents a higher-resolution SST. Yet using CNTRL instead of IDEA as a reference would yield qualitatively similar results due to the similarity of both simulations.

##### 4.1 Impact on air–sea interactions


The impact of SST perturbations on air–sea interactions is investigated through spatially averaged surface heat fluxes in the Gulf Stream region, spanning 30–55° N and 80–25° W. An analysis of the temporal changes in latent and sensible heat fluxes reveals a similar evolution of the surface fluxes across the different experiments (Fig. 3). First of all, the removal of the small-scale variability associated with individual Gulf Stream meanders in IDEA results in only small differences in spatially averaged surface latent and sensible heat fluxes compared to CNTRL (Table 2 and Fig. 3). More pronounced differences arise for the other experiments. The P1.5K experiment consistently shows the highest upward surface fluxes from the ocean to the atmosphere; in contrast the extWEAK experiment displays the lowest fluxes. This difference is particularly pronounced for latent heat flux

(Fig. 3a and Table 2) and also applies to a lesser extent to the sensible heat flux (Fig. 3b). Specifically, a latent heat flux increase of up to  $10 \text{ W m}^{-2}$  for P1.5K and a decrease of  $25 \text{ W m}^{-2}$  for extWEAK compared to IDEA are observed in the Gulf Stream region. Sensible heat flux differences of up to  $5 \text{ W m}^{-2}$  in P1.5K and up to  $20 \text{ W m}^{-2}$  in extWEAK compared to IDEA (Fig. 3b) are present. These differences are particularly pronounced during the passage of cyclones CY1 and CY2, which trigger CAO events in the western North Atlantic (Fig. 2b, e). During these events, the exchange of heat and moisture between the ocean and atmosphere intensifies, resulting from the large contrast between the colder and dry air following the cyclones and the warmer SST. From a temporally averaged perspective (not shown), the P1.5K experiment, with its warmed SST, shows the largest increase in mean upward surface heat fluxes around the Gulf Stream SST front. Conversely, in experiments with weakened SST gradients (WEAK and extWEAK), we observe a reduced magnitude (i.e., less negative) of surface heat fluxes, particularly south of the SST front. These findings align with prior research indicating reduced upward surface fluxes on the warmer, southern side of the SST front when the SST gradient is weakened (Vries et al., 2019; Tsopouridis et al., 2021).

##### 4.2 Impact on air temperature and specific humidity

The changes in sensible and latent heat fluxes impact the air temperature and specific humidity within the atmospheric boundary layer over the Gulf Stream region. Table 2 outlines the average differences in air temperature and specific humidity at 925 hPa in the Gulf Stream region between IDEA and each experiment. The removal of the small-scale variability of the Gulf Stream meanders in IDEA results in only small differences in surface heat fluxes (Table 2 and Fig. 3). Consequently, the differences in area-averaged temperature and specific humidity between CNTRL and IDEA are relatively small throughout the simulation period. Compared to the CNTRL experiment an average air temperature increase of 0.32 and 0.06 K are observed in the P1.5K and the CNTRL experiment, respectively. Likewise, an average specific humidity increase of 0.09 and 0.03  $\text{g kg}^{-1}$  are present. This is consistent with an increase in surface fluxes.

In light of our study's focus on WCBs, we further evaluate the variations in air temperature and specific humidity specifically within the WCB inflow region. This region is primarily located south of the SST front, where SSTs are relatively warm. This region is important for moisture supply for subsequent latent heat release during WCB ascent (Wenta et al., 2024). Here, the Eulerian WCB inflow region is determined based on the occurrence frequency of the positions of WCB air parcels in the IDEA experiment over the entire 9 d simulation period, specifically defined as locations where the trajectories' pressure exceeds 800 hPa. A time-invariant inflow region is justified because the WCB inflow from a La-



**Figure 2.** SST, CAOs, and PV at 315 K during the transition of cyclones CY1 and CY2 across the North Atlantic in February 2019. (a, d, g) Sea level pressure (grey contours, every 5 hPa), 2 PVU potential vorticity (PV) contour at 315 K (black bold), and sea surface temperature (SST, shading, in K) for (a) 12:00 UTC on 19 February 2019, (d) 12:00 UTC on 22 February 2019, and (g) 12:00 UTC on 24 February 2019. (b, e, h) CAO index, defined as  $\theta_{SST} - \theta_{850}$  (blue shading in K), sea level pressure (grey contours, every 5 hPa), and 2 PVU potential vorticity (PV) contour at 315 K (black bold) at the same times shown in panels (a), (d), and (g). (c, f, i) PV at 315 K (shading in PVU), WCB outflow masks (pink shading) at the same times shown in panels (a), (d), and (g), and WCB trajectories initialized 24 h prior to the time shown (colored by pressure; in hPa). All panels show the CNTRL simulation (see Sect. 2.1).


grangian perspective occurs repeatedly in this region. Subsequently, the inflow region is confined to areas where the WCB inflow is consistently present for at least 30 % of the 9 d simulation (Fig. A3). Notably, the location of this Eulerian WCB inflow region remains consistent across different experiments, owing to the similar synoptic evolution across all experiments. In the WCB inflow region, the evolution of the spatially averaged vertical profiles for air temperature and specific humidity shows gradually evolving differences between experiments (Figs. 4 and 5). In the WEAK and more notably in the extWEAK experiments, air temperature is generally slightly lower than in IDEA (Fig. 4b, c) with a simultaneous reduction in moisture content (Fig. 5b, c). Initially, temperature differences are confined to the lower tro-

posphere, but throughout the simulation differences up to the tropopause level arise. The difference is particularly striking in the extWEAK experiment, where the weakening of the SST gradient reduces the specific humidity from the surface up to 500 hPa several days into the simulation (Fig. 5c) with a concomitant decrease in temperature throughout the entire troposphere (Fig. 4c). In contrast, the P1.5K experiment shows an increase in both air temperature (Fig. 4d) and moisture content up to 700 hPa (Fig. 5d), in line with the previously discussed increase in surface sensible and latent heat fluxes (Sect. 4.1). The temperature differences between all experiments are particularly pronounced after the passages of cyclones CY1 and CY2 (Fig. 4) and extend from the boundary layer into the upper troposphere. Overall, the



**Table 2.** Differences between experiments for surface latent and sensible heat fluxes, as well as air temperature, specific humidity, and baroclinicity at 925 hPa. Differences are calculated relative to the idealized SST gradient experiment (IDEA) and are temporally averaged over the whole simulation period (19 to 27 February 2019) and spatially averaged across the Gulf Stream region (30 to 55° N and 80 to 25° W).

	Latent heat flux	Sensible heat flux	Air temperature	Specific humidity	Baroclinicity
IDEA–CNTRL	$1.83 \pm 15.21 \text{ W m}^{-2}$ –2.36 %	$0.39 \pm 9.90 \text{ W m}^{-2}$ –0.80 %	$-0.06 \pm 0.5 \text{ K}$ –0.02 %	$-0.02 \pm 0.29 \text{ g kg}^{-1}$ –0.67 %	$-0.0004 \text{ K km}^{-1}$ –0.9 %
IDEA–WEAK	$-6.64 \pm 21.01 \text{ W m}^{-2}$ 8.56 %	$-2.41 \pm 14.75 \text{ W m}^{-2}$ 4.95 %	$0.01 \pm 0.88 \text{ K}$ 0.0 %	$0.06 \pm 0.44 \text{ g kg}^{-1}$ 2.18 %	$0.001 \text{ K km}^{-1}$ 2.31 %
IDEA–extWEAK	$-15.95 \pm 33.5 \text{ W m}^{-2}$ 20.6 %	$-6.26 \pm 23.04 \text{ W m}^{-2}$ 12.83 %	$0.04 \pm 1.62 \text{ K}$ 0.02 %	$0.16 \pm 0.82 \text{ g kg}^{-1}$ 5.24 %	$0.003 \text{ K km}^{-1}$ 7.07 %
IDEA–P1.5K	$7.42 \pm 10.54 \text{ W m}^{-2}$ –9.58 %	$2.9 \pm 6.77 \text{ W m}^{-2}$ –5.95 %	$-0.32 \pm 0.58 \text{ K}$ –0.12 %	$-0.09 \pm 0.36 \text{ g kg}^{-1}$ –3.16 %	$-0.0006 \text{ K km}^{-1}$ –1.43 %




**Figure 3.** (a) Latent heat flux (LHF, in  $\text{W m}^{-2}$ , upward defined negatively) spatially averaged over the Gulf Stream region (30 to 55° N and 80 to 25° W) for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red). Panel (b) is the same as (a) but for sensible heat flux (SHF). Grey shading outlines the time period of the passage of cyclone 1 (CY1) and cyclone 2 (CY2) in the Gulf Stream region.

spatiotemporal evolution of temperature differences in the experiments illustrates the influence of the prescribed SST perturbations on the atmosphere, whereby differences evolve from the lower troposphere and subsequently progress upward after the passages of CY1 and CY2.


The evolution of temperature and humidity profiles described above is related to altered vertical motion coupled with changes in diabatic heating from cloud formation processes during WCB ascent. The availability of moisture in the WCB inflow region determines the degree of latent heat released during cloud formation and, thus, the cross-isentropic WCB ascent strength (e.g., Oertel et al., 2023a). First of all, we use accumulated precipitation in the North Atlantic (30 to 60° N and 80 to 0° W) as a proxy for the mass conversion of water vapor to the liquid and/or solid state. Throughout the 9 d period the accumulated precipitation dif-

fers between the experiments (Fig. 6). The highest precipitation sums, approximately 6 % more precipitation than in IDEA, occur in the P1.5K experiment, which suggests an enhancement in diabatic heating of the mid-troposphere during WCB ascent. Conversely, experiments with weaker SST gradients, specifically WEAK and extWEAK, accumulate 5 % and 16 % less precipitation, respectively, than IDEA, which, in turn, is associated with reduced diabatic heating.


The variability in precipitation sums suggests that differences in lower tropospheric moisture content across experiments noticeably influence diabatic heating rates from parameterized cloud microphysical processes and convection. The analysis of diabatic heating tendencies from the individual model parameterizations (Oertel et al., 2023a) confirms that cloud microphysics and the convection scheme substantially influence total heating and also differ between the experiments (Figs. A4, A5, and A6 and detailed description in Christ, 2023). Consistent with the differences in surface precipitation, the P1.5K experiment shows enhanced cloud microphysical diabatic heating rates above 3–4 km altitude (Fig. A4c), followed by the CNTRL (Fig. A4a), whereas the heating rates in the WEAK and extWEAK experiments are reduced (Fig. A4b,d). Similarly, the convection parameterization (Fig. A6), which is most active in the lowest 2 km, is more active in the P1.5K experiment, and reduced diabatic heating arises in the WEAK and extWEAK experiments. This indicates that temperature differences propagate from the sea surface into the boundary layer, where the WCB airstream picks up additional moisture and subsequently warms the mid-troposphere during ascent through latent heating, efficiently redistributing temperature and humidity. To conclude, diabatic heating during WCB ascent and associated surface precipitation sums are affected by the availability of moisture in the WCB inflow region, which in turn is influenced by SST perturbations in the Gulf Stream region. In Sect. 4.4, we elaborate in more detail on the impact on WCB ascent.



**Figure 4.** (a)–(d) Evolution of air temperature differences ( $\Delta T$ , shading, in K) spatially averaged in the Eulerian WCB inflow region defined as the area where the WCB inflow frequency in IDEA (WCB trajectories’ pressure larger than 800 hPa) exceeds 30 % during the 9 d simulation period. Differences are shown for (a) IDEA–CNTRL, (b) IDEA–WEAK, (c) IDEA–extWEAK, and (d) IDEA–P1.5K. Grey shading outlines the time period related to the passage of cyclone CY1 and cyclone 2 CY2 in the Gulf Stream region.




**Figure 5.** As Fig 4 but for specific humidity differences ( $\Delta q_v$ , shading, in  $\text{g kg}^{-1}$ ).



**Figure 6.** Evolution of spatiotemporally accumulated surface precipitation sums (in kg) in the North Atlantic ( $30$  to  $60^\circ$  N and  $80$  to  $0^\circ$  W) during the 9 d simulation period for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red). Grey shading outlines the time of the passage of cyclone 1 (CY1) and cyclone 2 (CY2) in the Gulf Stream region.

### 4.3 Impacts on cyclones

Building upon the discussions above, cyclones CY1 and CY2 and their associated WCBs play a critical role in amplifying and transmitting the influence of SST into higher atmospheric layers. Generally, the genesis of cyclones is linked to baroclinicity (Charney, 1947), which provides information on the potential energy available for cyclones. We calculate baroclinicity as the horizontal gradient of equivalent potential temperature at the 925 hPa level. This brings into focus the potential impact of the strength of the SST gradient on cyclone development through changes in low-level baroclinicity in the Gulf Stream region. Spatial averages of low-level baroclinicity in the Gulf Stream region (Fig. 7a, Tab. 2) demonstrate a reduction of baroclinicity in the WEAK and notably the extWEAK experiments in comparison to IDEA with on average 2.3 % and 7 % reduction, respectively. Despite an identical SST gradient, the P1.5K experiment shows a slight increase in baroclinicity compared to IDEA, which



**Figure 7.** (a) Evolution of baroclinicity at 925 hPa ( $\nabla\Theta_e$ , in  $\text{K km}^{-1}$ ) for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red) spatially averaged over the Gulf Stream region (30 to 55° N and 80 to 25° W). (b, c) Evolution of minimum sea level pressure (SLP) in the cyclone center (in hPa) for (b) the first cyclone (CY1) and (c) the second cyclone (CY2) for the experiments shown in (a). Grey shading outlines the time of the passage of cyclone 1 (CY1) and cyclone 2 (CY2) in the Gulf Stream region. Blue shading in (a) indicates the times of CY1 and CY2 shown in (b) and (c), respectively.

can be attributed to differential heating and/or advection. In contrast, for spatial averages, only minor differences between CNTRL and IDEA experiments are present, although for temporal averages locally larger differences arise, reflecting the patterns of surface heat flux differences. This suggests that very small-scale SST meanders have a small effect on baroclinicity in our experiments.


Yet experiments with modified baroclinicity and moisture availability resulting from changes in air–sea interactions show different cyclone development. The analysis of cyclone tracks for CY1 and CY2 illustrates that while the cyclones’ positions (Fig. A2) and their peak intensity time (at 21:00 UTC on 20 February 2019 for CY1 and at 00:00 UTC on 24 February 2019 for CY2) remain consistent across different experiments, their intensity notably varies. The intensity is measured by the minimum SLP at the cyclone center (Fig. 7b,c). CY1 deepens more strongly in CNTRL and P1.5K, and minimum SLP drops to 943 hPa in both experiments, compared to the IDEA with a minimum SLP of 945 hPa (Fig. 7b). A slight reduction in deepening rates is observed for the WEAK and extWEAK experiments, which reach an SLP minimum of 945 and 950 hPa, respectively. Cyclone intensification differences between experiments are substantially larger for CY2 (Fig. 7c) than for CY1, for which differences on the order of 15 hPa occur. CY2 in WEAK has a higher SLP (minimum of 971 hPa) than IDEA, and the minimum SLP in extWEAK is markedly higher (minimum of 980 hPa). Conversely, the P1.5K cyclone demon-

strates the most rapid and strongest deepening to a minimum of 964 hPa. Meanwhile, the evolution of minimum SLP in CNTRL and IDEA is relatively similar with a minimum of 968 hPa; i.e., SLP minimum differences for CY2 reach about 2 hPa. The stronger effect of SST perturbations on CY2 suggests that the impact of modified air–sea interactions on the synoptic evolution increases with lead time, likely also influenced by strong surface fluxes associated with the CAO following CY1. Overall, the reduction of cyclone intensity in experiments with weaker SST gradients (WEAK and extWEAK) aligns with the decreased low-level baroclinicity and reduced diabatic heating (Sect. 4.2).

#### 4.4 Impact on WCB ascent

The deep ascent of WCB airstreams through the entire atmospheric column can connect low-level processes and upper-level flow patterns. To gain insight into the relationship between SST in the Gulf Stream region and WCB ascent, we employ a Lagrangian approach to objectively characterize WCB properties. The WCBs associated with cyclones CY1 and CY2 ascend poleward from the warm sector of the cyclones ahead of the cold front into the upper-level ridge (see, e.g., Fig. 2i). The typical ascent characteristics of identified WCBs are depicted for the trajectories from IDEA (Fig. 8). The average ascent of WCB air parcels starts in the inflow layer, below 800 hPa, and ends in the outflow layer, above 400 hPa (Fig. 8a). Due to cloud formation processes during ascent, specific humidity continuously decreases from  $\sim 7$  to  $\sim 0 \text{ g kg}^{-1}$  (Fig. 8b). The increase in liquid and ice water contents (Fig. 8c) during the ascent signifies the onset of cloud formation processes. Latent heat release, primarily from condensation and depositional growth of ice hydrometeors (e.g., Oertel et al., 2023a), enables the cross-isentropic ascent and increases the WCB air parcels’ potential temperature from, on average, 289 to 305 K (Fig. 8d). The increase in potential temperature resulting from latent heating is linked to a modification of PV. PV is generated below the level of maximum latent heat release and reduced above (Fig. 8e), leading to low PV values in the upper tropospheric WCB outflow region (Fig. 11c) and aligning with the conceptual model of PV changes along WCB ascent (Hoskins et al., 1985; Wernli, 1997; Madonna et al., 2014).

In the subsequent sections, we will explore the differences in WCB trajectory characteristics between the five SST experiments (Table 3). Figure 9 displays the number of WCB trajectories starting their ascent every hour throughout the simulations. Of the total 40 710 trajectories that were started every hour, approximately 2000 to 6000 are identified as WCB trajectories depending on the experiment. The two local maxima in WCB trajectory number on 19 and 22 February are associated with the passages of cyclones CY1 (18 to 20 February 2019) and CY2 (22 to 23 February) across the Gulf Stream region (blue shading in Fig. 9). The overall evolution of WCB trajectory starts is similar across all



**Figure 8.** Mean evolution of (a) pressure ( $p$ , in hPa), (b) specific humidity ( $q_v$ , in  $\text{g kg}^{-1}$ ), (c) liquid and ice hydrometeor contents ( $q_{LI}$ , blue and light-blue lines, in  $\text{g kg}^{-1}$ ), (d) potential temperature ( $\theta$ , in K), and (e) potential vorticity (PV, in PVU) along WCB trajectories. The PV evolution in (e) is centered relative to maximum diabatic heating (DH). The shading shows the mean  $\pm$  standard deviation. All figures are shown for the IDEA WCB trajectories.


experiments. However, the difference in absolute WCB trajectory numbers, and thus mass transport across the troposphere, is considerably large, and relative differences of up to 20%–50% emerge between experiments (Fig. 9b, Table 3). An up to 20% increase in WCB trajectory numbers in P1.5K compared to IDEA indicates a stronger WCB ascent associated with enhanced mass flux. Contrarily, up to 50% fewer WCB trajectories are present in extWEAK (Fig. 9b), indicating substantially weakened WCB ascent. Similarly, the WCB trajectory numbers of WEAK are reduced by up to 20% compared to IDEA. Overall, the difference in the number of WCB trajectories between the IDEA and CNTRL experiments is relatively small but not negligible. On average 4.03% more WCB trajectories ascend in the CNTRL, and differences of up to 20% occur on the third day lead time, generally showing a tendency for higher numbers in the CNTRL experiment. Therefore, small-scale ocean eddies present in the CNTRL influence WCB ascent and, thus, potentially impact the upper-level circulation as suggested in Liu et al. (2021). On average the largest number of WCB trajectories is present in P1.5K (Table 3) with 9.4% more WCB trajectories than in IDEA. A reduction of SST gradient reduces the numbers of WCB trajectories by on average  $-9.43\%$  in WEAK and  $-30.07\%$  in extWEAK. The

consistent differences between experiments at all lead times demonstrate that the SST in the Gulf Stream region impacts the WCBs systematically, at least in this case. Thereby, this influence aligns with the (thermo-)dynamic understanding of WCB ascent.

Ascent characteristics of WCB trajectories are often described by the distribution of their ascent times (e.g., Oertel et al., 2021). The distribution of ascent timescales reveals that the typical duration, in this case, lies within 9–20 h (Fig. 9c). The comparison across all experiments shows a shift in ascent timescale distribution (Fig. 9c, Table 3). Notably, in the extWEAK experiment, the distribution of ascent timescales shifts towards longer durations, indicating slower ascent (Fig. 9c). Specifically, the frequency of ascent timescales below 18 h is substantially reduced (Fig. 9c). In the WEAK experiment, rapid ascent with timescales below 10 h occurs less frequently, whereas in the P1.5K experiment, and to a smaller extent in the CNTRL experiment, rapid ascent is more frequent. To summarize, simulations with a reduced SST gradient are characterized by weaker WCB ascent and a lower frequency of fast ascents. Conversely, higher SST in P1.5K results in an increased number of WCB trajectories, especially in a larger number of rapidly ascending WCB trajectories.

**Table 3.** Differences in WCB characteristics between the five experiments. Shown are temporally averaged absolute and relative warm conveyor belt (WCB) trajectory numbers as well as mean 500 hPa ascent times ( $\tau_{500}$ ), absolute pressure change during WCB trajectories' ascent ( $\Delta p$ ), absolute specific humidity change ( $\Delta q_v$ ), absolute change of potential temperature ( $\Delta\Theta$ ), trajectories' mean minimum pressure ( $\min p$ ), and mean maximum potential temperature ( $\max \Theta$ ). Differences are calculated relative to the idealized SST gradient experiment (IDEA) and averaged over all WCB trajectories for the entire period (19 to 25 February 2019).

	WCB number	$\tau_{500}$	abs. $\Delta p$	$\Delta q_v$	$\Delta\Theta$	$\min p$	$\max \Theta$
IDEA–CNTRL	–124	–0.19 h	0.51 hPa	–0.06 g kg <sup>–1</sup>	–0.19 K	–1.68 hPa	–0.19 K
	–4.03 %	–0.91 %	0.09 %	–0.87 %	–0.99 %	–0.45 %	–0.06 %
IDEA–WEAK	291	–0.63 h	2.48 hPa	0.30 g kg <sup>–1</sup>	0.80 K	–2.90 hPa	1.33 K
	9.43 %	–3.02 %	0.43 %	4.11 %	4.21 %	–0.78 %	0.44 %
IDEA–extWEAK	927	–2.54 h	9.98 hPa	0.74 g kg <sup>–1</sup>	2.20 K	–11.65 hPa	3.57 K
	30.07 %	–12.16 %	1.73 %	10.10 %	11.62 %	–3.13 %	1.17 %
IDEA–P1.5K	–290	0.22 h	–2.35 hPa	–0.17 g kg <sup>–1</sup>	–0.52 K	1.95 hPa	–0.79 K
	–9.40 %	1.05 %	–0.41 %	–2.32 %	–2.72 %	0.52 %	–0.26 %



**Figure 9.** (a) Number of WCB trajectories starting between 18 and 25 February 2019 for IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red). The time axis represents the time of the start of the 2 d WCB trajectory ascent. Panel (b) is the same as (a) but for the WCB trajectory number fraction relative to IDEA (in %). Grey shading in (a) and (b) outlines the time according to the passage of cyclone 1 (CY1) and cyclone 2 (CY2) in the Gulf Stream region. (c) Histograms of the fastest 500 hPa ascent times for all WCB trajectories for the five experiments (bin width 1 h); colors are as in panels (a) and (b).

In addition to differences in the number of WCB trajectories and ascent timescale distributions, other WCB properties are influenced by the SST perturbations (Table 3). These differences are evident not only from the Lagrangian perspective but also from the Eulerian viewpoint, which is discussed in Sect. 4.2. As shown above, WEAK and extWEAK are characterized by reduced moisture and heat supply to the WCB inflow region due to altered air–sea interactions and vice versa for P1.5K and CNTRL. This results in differences in specific humidity as well as potential temperature and pressure changes during the WCB ascent among the experiments (Fig. 10, Table 3). Due to larger specific humidity in the WCB inflow, WCB ascent in P1.5K is associated with the largest average loss of specific humidity of  $9 \text{ g kg}^{-1}$  (Fig. 10b), which translates to the largest diabatic heating of approximately 22 K (Fig. 10c). The slightly lower specific humidity loss in WEAK and extWEAK of on average 8.75 and  $8.4 \text{ g kg}^{-1}$  (Fig. 10b) consequently leads to a reduced diabatic heating of 21 and 19 K, respectively (Fig. 10c).

Consistent with larger moisture loss and diabatic heating, P1.5K shows the deepest ascent quantified by an average absolute pressure change of 595 hPa (Fig. 10a). The WCB trajectory ascent in the WEAK and extWEAK is weaker than in IDEA with pressure differences of on average 585 and 575 hPa, while the differences between IDEA and CNTRL are rather small.

Overall, the average differences in diabatic heating (Figs. 6 and 10c), as well as ascent strength, among the experiments (Fig. 10) are consistent throughout the analysis period. Differences in specific humidity and heat supply to the WCB inflow region in response to changed air–sea interactions lead to different WCB characteristics and diabatic heating during ascent. Intuitively, reduced specific humidity loss during WCB ascent in the weaker SST gradient experiments results in lower diabatic heating and weaker ascent. Vice versa, increased surface fluxes from higher SST in the P1.5K experiment, and to a smaller extent in the CNTRL, lead to faster and more pronounced WCB ascent.


The altered diabatic processes during the WCB ascent also influence the WCB characteristics in the upper troposphere (Table 3). On average, WCB air parcels in the P1.5K experiment reach 320 hPa, which is very close to the average pressure in CNTRL and IDEA of 318 hPa for both experiments. In contrast, in WEAK and extWEAK, average WCB outflow height is lower with mean minimum pressure values of  $\sim 330$  and 345 hPa, respectively (Fig. 11a). Furthermore, differences in average potential temperature in the WCB outflow, i.e., maximum isentropic level, are present and vary between 2 and 10 K (Fig. 11b, Table 3). This aligns with differences in diabatic heating: throughout the simulation reduced cross-isentropic ascent in WEAK and extWEAK compared to IDEA and CNTRL results in lower maximum potential temperature in the WCB outflow, while WCB trajectories in P1.5K reach slightly higher isentropic levels. Although differences in WCB outflow potential temperature are

present, PV values averaged in the WCB outflow (defined as all WCB trajectory positions with pressure values lower than 400 hPa) do not vary substantially between the experiments and amount to on average 0.2 PVU (Fig. 11c).

#### 4.5 Impact on the large-scale flow

In all experiments, the WCB outflow predominantly aligns with the intensifying upper-level ridge, contributing to the formation of the quasi-stationary anticyclone associated with the European blocking. Figure 12 illustrates the impacts of modified WCB outflow properties on the upper-tropospheric ridge characteristics by regridding WCB trajectory outflow positions to the Eulerian grid. The WCB outflow region is located near the western coast of Europe downstream of the Gulf Stream region. Only small spatial differences in WCB outflow location between experiments are present, which is consistent with an overall similar large-scale flow evolution. Yet the northern boundary of the WCB outflow in WEAK and extWEAK is located slightly further south than in IDEA (Fig. 12b, d). In both experiments, the position of the upper-level jet is also displaced further south. We hypothesize that the weaker divergent outflow of WCBs in WEAK and extWEAK influences the amplification of the downstream ridge. A consistent change in WCB outflow location is present in P1.5K. Specifically, the WCB outflow is slightly shifted poleward, which is also associated with a small poleward displacement of the jet stream position region (Fig. 12c). In addition to small spatial shifts, the average potential temperature in the WCB outflow region differs (Fig. 12; see also Fig. 11b and Table 3). The large part of WCB outflow potential temperature in WEAK and extWEAK is lower than in IDEA (Fig. 12b,d), and contrarily, potential temperature in P1.5K is larger than in IDEA (Fig. 12c), which is consistent with WCB trajectories' average maximum isentropic level. Thus, in addition to poleward and equatorward displacement, respectively, the vertical position of WCB trajectories is also influenced by the modification of low-level specific humidity and temperature, with WCB air parcels reaching only a lower outflow height in the experiments with weaker SST gradients and a higher outflow height in the experiment with increased SSTs.

In the following, we illustrate how changes in WCB ascent are also reflected in the large-scale atmospheric circulation. As shown in Fig. 12, the divergent upper-tropospheric WCB outflow influences the amplification of the downstream ridge (see also, e.g., Grams et al., 2011; Pfahl et al., 2015; Steinfeld and Pfahl, 2019; Oertel et al., 2023a). Specifically, the influence of modified air–sea interactions on the large-scale flow modulated by WCB ascent is reflected in the temporal evolution of geopotential height at 500 hPa (Z500) in the western North Atlantic where the downstream ridge develops ( $40$  to  $70^\circ \text{ N}$  and  $30^\circ \text{ W}$  to  $20^\circ \text{ E}$ ). After 3 d of lead time, the influence of WCB trajectories on the upper-level ridge becomes apparent (Fig. 13). The onset of the blocking




**Figure 10.** (a) Evolution of mean absolute pressure change during WCB trajectories' ascent ( $\Delta p$ , in hPa; the date indicates the start time of trajectories) for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red) including mean  $\pm$  standard deviation (shading). Panel (b) is as (a) but for specific humidity change ( $\Delta q_v$ , in  $\text{g kg}^{-1}$ ). Panel (c) is as (a) but for absolute change of potential temperature ( $\Delta \Theta$ , in K). Grey shading outlines the time period of the passage of cyclone 1 (CY1) and cyclone 2 (CY2) in the Gulf Stream region.

associated with an increase in Z500 aligns with the emergence of differences in the Z500 evolution among experiments. The highest Z500 values, i.e., the strongest ridge, are present in P1.5K and CNTRL, while the lowest are found in WEAK and extWEAK. When the upper-level ridge is fully developed and Z500 values peak, the maximum difference between the experiments (extWEAK, P1.5K and CNTRL) amounts to almost 4 gpdm (geopotential decimeter) (Fig. 13). Clear differences in the temporally averaged Z500 fields emerge (Fig. 14). The ridge in experiments with a weakened SST gradient (WEAK and extWEAK) is weaker than in IDEA (Fig. 14b, c), while P1.5K and CNTRL are characterized by overall higher Z500 values (Fig. 14d). In the context of Z500 climatological anomalies, typically ranging from 15–30 gpdm in the Euro-Atlantic sector relative to a seasonal climatology (Grams et al., 2017), the observed average differences of 4 gpdm between the P1.5K, CNTRL, and extWEAK experiments are noteworthy. These discernible

differences underscore the sensitivity of downstream ridge development to modifications in SST in the Gulf Stream region. Our results show that changes in SST mainly impact the strength of WCB ascent while not drastically altering the overall synoptic pattern. As such, it is unlikely to see major changes in the upper-level flow. Yet these findings suggest that blocking intensity in the Euro-Atlantic sector is sensitive to the Gulf Stream SST pattern and mesoscale ocean eddies and influenced by changes in WCB ascent characteristics resulting from modified air–sea interactions.

Overall our findings suggest that SSTs within the Gulf Stream region indeed can affect the downstream flow evolution by influencing the WCB inflow region (Figs. 4 and 5) through modified surface fluxes (Fig. 3), which subsequently modifies WCB ascent characteristics (Figs. 9 and 10). These perturbations then also propagate into the middle to upper troposphere (Fig. 11), where the WCB interacts with the large-scale flow and influences the strength of the down-



**Figure 11.** As Fig. 10 but for (a) mean minimum pressure ( $p$ ) of WCB trajectories (in hPa), (b) mean maximum potential temperature ( $\Theta$ ) of WCB trajectories (in K), and (c) mean potential vorticity (PV) of WCB trajectory outflow (in PVU).

stream ridge (Fig. 13). We conclude that the WCB airstream acts as a mechanistic link between the lower and upper troposphere and can efficiently transfer near-surface perturbations upward, linking the air–sea interactions over the Gulf Stream with the large-scale flow.

## 5 Summary and discussion


This study investigates the effects of SST perturbations in the Gulf Stream region on the evolution of the downstream flow during an atmospheric blocking event in February 2019. A series of sensitivity experiments with the ICON model were conducted to evaluate the impact of SST changes in the Gulf Stream region on the formation and maintenance of the downstream block. Thereby, a particular focus is on improving our understanding of the physical processes that link near-surface processes with the upper-level flow. The sensitivity experiments include five free-running simulations: (i) a control simulation (CNTRL) with SST taken from the ECMWF IFS analysis, (ii) an experiment with an idealized SST gradient devoid of small-scale meanders (IDEA), (iii) a simulation with a reduced Gulf Stream SST gradient

(WEAK), (iv) an extreme scenario for which the SST gradient in the Gulf Stream was almost completely removed (extWEAK), and (v) a simulation with increased SST while preserving the SST gradients (P1.5K). In each of the experiments, the synoptic-scale dynamics, including cyclone formation, WCB airstream ascent, and downstream ridge development, are well represented and largely similar. However, smaller, non-negligible differences arise between experiments. These subtle changes affect both boundary layer dynamics and processes in the upper troposphere, highlighting the relevance of the representation of SST for influencing the chain of events that ultimately impact the upper-level flow evolution.


The comparison of the IDEA and CNTRL experiments, which differ only by smoothing local SST gradients, indicates that small-scale SST meanders have a smaller influence on atmospheric conditions during the 9 d simulation period compared to the other experiments. This aligns with Tsopouridis et al. (2021), who reported limited effects on cyclone dynamics after smoothing SST gradients in the Gulf Stream and Kuroshio currents. Similarly, the subtle differences between the CNTRL and IDEA experiments



WCB outflow | 24. Feb at 00 UTC



**Figure 12.** (a)–(d) Potential temperature difference ( $\Delta\Theta$ , shading, in K) in the WCB outflow at 00:00 UTC on 24 February 2019 for (a) IDEA–CNTRL, (b) IDEA–WEAK, (c) IDEA–extWEAK, and (d) IDEA–P1.5K. The WCB outflow region in IDEA is outlined in black (bold contours), and the WCB outflow for the respective experiments is shown in color. Also shown are sea level pressure (contours, every 5 hPa) and 2 PVU potential vorticity contour at 315 K PV for IDEA (black) and the respective experiment (color).




**Figure 13.** Evolution of geopotential height at 500 hPa ( $Z_{500}$ , in gpdm) for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red) spatially averaged over the ridge region downstream of the Gulf Stream region (40 to 70° N and 30° W to 20° E).

within this short period of only several days are consistent with Roberts et al. (2021), who suggested a minimal impact of small-scale SST meanders on atmospheric dynamics for short lead times. However, the differences are not negligible and indicate an impact of small-scale ocean eddies on the WCB trajectory number and ascent behavior, which subsequently influences the upper-level circulation. In the error growth framework (e.g., Baumgart et al., 2019), such small-scale errors can matter for predictability, in particular when they are projected to the tropopause region by the WCB, where errors grow nonlinearly. Due to the determin-

istic nature of our simulations, it remains open to how large and systematic the differences between IDEA and CNTRL are compared to differences in an initial condition ensemble. Moreover, the role of small ocean eddies may be more significant over longer periods. Roberts et al. (2021) indicated that while SST biases or inaccuracies may not substantially influence individual synoptic events, as demonstrated in our study, their impact could be more pronounced on subseasonal scales. Furthermore, Roberts et al. (2022) suggested that an increased ocean resolution in atmosphere–ocean coupled simulations could lead to a more noticeable overall impact, primarily due to a better representation of the SST evolution over extended timescales. The latter is not accounted for in our experiments as SST patterns remain constant throughout the comparatively short simulation time.

In contrast to the difference between CNTRL and IDEA, larger differences compared to IDEA (which would also be true compared to CNTRL) were found for the WEAK, extWEAK, and P1.5K experiments. In particular, differences in the strength of air–sea interactions, the characteristics of WCB ascent, and the structure of the downstream ridge are present. Differences between experiments are most apparent in the boundary layer and particularly pronounced during the passage of cyclones, which also advect cold continental air over the relatively warm ocean surface, leading to CAOs (Fig. 15a). The low-level air which had been heated and moistened as a consequence of the passage of the first cyclone (CY1) is subsequently drawn into the WCB of the



**Figure 14.** Differences in temporally averaged (21 to 27 February 2019) geopotential height ( $\Delta Z500$ , shading, in gpdm) in the ridge region downstream of the Gulf Stream (40 to 70° N and 30° W to 20° E, blue box) for (a) IDEA–CNTRL, (b) IDEA–WEAK, (c) IDEA–extWEAK, and (d) IDEA–P1.5K. The 2 PVU potential vorticity contour at 315 K PV at 12:00 UTC on 25 February 2019 for IDEA (black) and the respective experiment (in color) are shown as well.


second cyclone (CY2). The latter has also been referred to as a “hand-over mechanism” (Papritz et al., 2021). An increase in SST in the P1.5K experiment leads to enhanced upward latent and sensible heat fluxes, which result in larger low-level moisture. In contrast, the WEAK and extWEAK experiments with colder SSTs south of the Gulf Stream SST front due to the reduced SST gradients are associated with reduced surface fluxes and, thus, low-level moisture. This agrees with Small et al. (2014), who noted lower surface heat fluxes, reduced moisture content, and colder air temperature in the lower troposphere as a response to a weakened SST gradient.

Modifications of low-level moisture are relevant for subsequent flow evolution because the availability of moisture in the WCB inflow region influences its ascent. WCB ascent is more pronounced and on average faster in the P1.5K experiment, in which specific humidity prior to WCB ascent is higher (Fig. 15b). During the ascent higher initial moisture availability leads to a more efficient conversion of water vapor into hydrometeors, resulting in enhanced diabatic heating. In contrast, the WEAK and extWEAK experiments show reduced diabatic heating as well as weaker and slower WCB ascent. This emphasizes the importance of specific humidity content in the WCB inflow region for its ascent behavior, which agrees with previous studies (Schemm et al., 2013; Schäfler and Harnisch, 2015; Oertel et al., 2021, 2023a; Joos et al., 2023). Moreover, differences in low-level moisture availability, together with changes in the temperature distribution and low-level baroclinicity, influence cyclone intensity, while cyclone tracks are not affected. Correlations between moisture availability and cyclone intensity have already been reported by Booth et al. (2012). However, it is ac-

knowledged that the relative contributions of moisture availability, static stability (from surface heating), or baroclinicity to cyclone intensification are not disentangled here.

SST modifications not only influence WCB ascent behavior and trajectory number but also influence the WCB outflow properties in the upper troposphere. Potential temperature in the WCB outflow P1.5K experiment is higher than in IDEA, while it is lower in the WEAK and extWEAK experiments, which follows from differences in diabatic heating during ascent. Thus, WCB trajectories with positive humidity perturbations in their inflow region on average reach higher isentropic levels (see also Oertel et al., 2023a). This impact on WCB outflow properties distinctly contrasts with the effect of stochastically perturbed parameterization tendencies (SPPTs; Buizza et al., 1999) on WCB outflow properties as reported by Pickl et al. (2023). They found that usage of SPPTs increased the number of WCB trajectories, whereas the average diabatic heating or outflow properties, such as outflow potential temperature, remained unchanged. It is noteworthy that the relatively subtle differences observed in the upper-level ridge across the experiments can be attributed to some extent to the duration of the simulations. We hypothesize that longer lead times may result in more pronounced discrepancies. Yet the differences in Z500 between experiments are still remarkable when considered relative to the range of climatological Z500 anomalies. However, we also note that for substantially longer lead times, such as subseasonal-to-seasonal timescales, atmosphere–ocean coupled simulations or SST updates are required.

The results presented here support previous research underscoring the link between Gulf Stream SST and downstream blocking events in the North Atlantic and western




**Figure 15.** Schematic of process chain linking air–sea interactions in the Gulf Stream to the amplification of a blocking event over Europe. (a) The first cyclone (CY1) triggers a cold air outbreak with associated high surface fluxes increasing low-level specific humidity (light-blue shading). (b) The second cyclone (CY2) travels into the region preconditioned by CY1. The WCB of CY2 originates in the region preconditioned by high surface fluxes and ascends, leading to cloud formation and latent heating. This associated divergent WCB outflow amplifies and maintains the downstream ridge (grey dashed contour) associated with a surface high ( $H$ ). The sensitivity of the blocking to SST emerges through the WCB linking humidity conditions near the surface with the upper-level flow.

Europe (e.g., O’Reilly et al., 2016; Yamamoto et al., 2021). We conclude that SST perturbations affect WCB characteristics consistently at every stage of WCB development, i.e., in the inflow, ascent, and outflow stages. The WCB is thus an important link that connects the SST changes in the Gulf Stream region with the dynamics in the middle and upper troposphere, thereby influencing the large-scale circulation and the strength of atmospheric blocking downstream, which is in line with Wenta et al. (2024). Figure 15 illustrates the chain of synoptic events and various interacting processes.


While this study provides valuable insights into the underlying physical mechanisms, it is based on a specific case study and a single model setup. Therefore, its findings cannot be universally applied to other cases and synoptic situations. This is especially relevant as the passage of several intense cyclones across the Gulf Stream region in this case increased the atmosphere’s sensitivity to SST changes. A further limitation of our analysis is the use of an atmosphere-only simulation, where SSTs are static and not dynamically coupled to the ocean. While this setup allows us to isolate the impact of SSTs on synoptic evolution, it omits feedback processes between ocean and atmosphere, such as transient cooling of SST after cyclone passage (Dacre et al., 2020). These effects may be less significant in the cyclone’s warm sector. The simplified SST conditions used in this study were designed to specifically assess the atmospheric sensitivity to underlying SSTs and the propagation of related perturbations through the atmosphere to the upper-level jet level. We believe that the key conclusions and the identified physical mechanisms remain valid regardless of the use of an interactive ocean model or static SST. Yet for future analysis, it would be interesting to include a coupled ocean with a sub-

sequently artificially smoothed SST gradient. Besides, with this setup, we were unable to determine whether the absolute SST values or the local SST gradients play a more important role in cyclone intensification and WCB ascent. To unravel these complex processes and feedback mechanisms and to better understand the role of air–sea interactions, especially in coupled systems, future research is required, including, for example, a coupled ocean model or time-varying prescribed SST fields. Furthermore, to quantify the effects of systematically perturbed lower boundary conditions in comparison to the impacts from uncertainties in initial conditions, ensemble simulations with an initial condition perturbation for the experiments would be beneficial. In addition to the systematic impact of SSTs quantified in our deterministic setup, this would provide uncertainty information and emphasize the SST effects compared to unsystematic variations as a consequence of (chaotic) error growth and intrinsic predictability limits. Nevertheless, the systematic and consistent impacts in our deterministic setup across all lead times and simulations, which are also consistent with the (thermo-)dynamic processes, provide confidence in the described underlying physical mechanisms. Hence, despite the mentioned limitations, we believe that the identified consistent impacts and the process-oriented approach adopted in this study significantly contribute to understanding how different SST representations can affect large-scale atmospheric flow evolution.


Appendix A




**Figure A1.** (a, c, e) Sea level pressure (contours, every 10 hPa) for ERA5 (blue), CNTRL (brown), and IDEA (black). (b, d, f) Potential vorticity contour (PV) at 315 K (shading in PVU) for ERA5 and 2 PVU potential vorticity contour at 315 K for CNTRL (brown) and IDEA (black) at (a, b) 00:00 UTC on 18 February, (c, d) 12:00 UTC on 22 February, and (e, f) 12:00 UTC on 24 February.




**Figure A2.** (a) Cyclone tracks of the first cyclone (CY1) from 10:00 UTC on 18 February to 19:00 UTC on 21 February (hourly) for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red). Sea surface temperature (SST, shading in K, for IDEA). (b) Same as (a) but for the second cyclone (CY2) from 20:00 UTC on 21 February to 09:00 UTC on 25 February.




**Figure A3.** Eulerian WCB inflow region, defined as the area where the occurrence of WCB inflow (WCB trajectories’ pressure larger than 800 hPa) exceeds 30 % (contours for CNTRL (brown), IDEA (black), WEAK (green), extWEAK (light green), and P1.5K (red)) during the 9 d simulation period. This area corresponds to the WCB inflow region definition used in Figs. 4 and 5. In addition, the 70 % frequency is also shown.



**Figure A4.** Evolution of vertical profiles of diabatic heating rates (DH, shading, in  $\text{K h}^{-1}$ ) for IDEA spatially averaged over the Gulf Stream region ( $30$  to  $55^\circ\text{N}$  and  $80$  to  $25^\circ\text{W}$ ). DH rates are shown for (a) the first saturation adjustment (Sat1), (b) the second saturation adjustment (Sat2), (c) microphysics (Micro), (d) turbulence (Turb), (e) convection (Conv), (f) longwave radiation (RadL), (g) shortwave radiation (RadS), and (h) total DH rate. See Oertel et al. (2023a) for a detailed description and overview of heating rates. Panels (a), (c), and (e) also show total hydrometer content (pink contours, in  $\text{g kg}^{-1}$ ).



**Figure A5.** Evolution of differences in diabatic heating rate profiles related to cloud processes, i.e., microphysics and the first saturation adjustment (DH, shading, in  $\text{K h}^{-1}$ ) spatially averaged over the Gulf Stream region ( $30$  to  $55^\circ \text{N}$  and  $80$  to  $25^\circ \text{W}$ ) for (a) IDEA–CNTRL, (b) IDEA–WEAK, (c) IDEA–extWEAK, and (d) IDEA–P1.5K. Also shown is the total diabatic heating rate for IDEA (black contours, at  $0.1 \text{ K h}^{-1}$ ).



**Figure A6.** As Fig. A5 but for diabatic heating rates from convection scheme (DH, shading, in  $\text{K h}^{-1}$ ) spatially averaged over the Gulf Stream region ( $30$  to  $55^\circ \text{N}$  and  $80$  to  $25^\circ \text{W}$ ) for (a) IDEA–CNTRL, (b) IDEA–WEAK, (c) IDEA–extWEAK, and (d) IDEA–P1.5K. Also shown is the convective heating rate for IDEA (black contours, at  $0.1 \text{ K h}^{-1}$ ).

*Data availability.* WCB trajectories and SST patterns from the experiments are publicly available in RADAR4KIT (<https://doi.org/10.35097/2ps96kskhd64de0g>) (Christ et al., 2024).

*Author contributions.* SC, AO, and MW performed the analyses, simulations, and trajectory calculations. SC wrote the first draft of the manuscript. All authors contributed to the design of the study and discussions and revisions of the manuscript.

*Competing interests.* At least one of the (co-)authors is a member of the editorial board of *Weather and Climate Dynamics*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

*Acknowledgements.* The research leading to these results has been carried out within the subproject B8 of the Transregional Collaborative Research Center SFB/TRR 165 “Waves to Weather” (<http://www.wavestoweather.de>, last access: 30 July 2024) funded by the German Research Foundation (DFG). Moreover, Marta Wenta is supported by DFG (grant no. GR 5540/2-1). Christian M. Grams was supported by the Helmholtz Association as part of the Young Investigator Group “Sub-seasonal Predictability: Understanding the Role of Diabatic Outflow” (SPREADOUT; grant no. VH-NG-1243). The contribution of Annika Oertel was partly carried out within the Italia–Deutschland science-4-services network in weather and climate (IDEA-S4S; INVACODA, grant no. 4823IDEAP6). This Italian–German research network of universities, research institutes, and the Deutscher Wetterdienst is funded by the BMDV (Federal Ministry of Transport and Digital Infrastructure). The authors acknowledge support from the state of Baden-Württemberg through bwHPC. The ICON simulations were carried out on the supercomputer HoreKa at Karlsruhe Institute of Technology, Karlsruhe, which is funded by the Ministry of Science, Research and the Arts Baden-Württemberg, Germany, and the German Federal Ministry of Education and Research. We are very grateful to Nedjeljka Žagar for discussions and her feedback on this study.

*Financial support.* This research has been supported by the Deutsche Forschungsgemeinschaft (grant nos. SFB/TRR 165 and GR 5540/2-1), the Helmholtz Association (grant no. VH-NG-1243), and the Bundesministerium für Digitales und Verkehr (grant no. 4823IDEAP6).

The article processing charges for this open-access publication were covered by the Karlsruhe Institute of Technology (KIT).

*Review statement.* This paper was edited by Silvio Davolio and reviewed by two anonymous referees.

## References

- Athanasiadis, P. J., Ogawa, F., Omrani, N.-E., Keenlyside, N., Schiemann, R., Baker, A. J., Vidale, P. L., Bellucci, A., Ruggieri, P., Haarsma, R., Roberts, M., Roberts, C., Novak, L., and Gualdi, S.: Mitigating Climate Biases in the Midlatitude North Atlantic by Increasing Model Resolution: SST Gradients and Their Relation to Blocking and the Jet, *J. Climate*, 35, 6985–7006, <https://doi.org/10.1175/JCLI-D-21-0515.1>, 2022.
- Baumgart, M., Ghinassi, P., Wirth, V., Selz, T., Craig, G. C., and Riemer, M.: Quantitative View on the Processes Governing the Upscale Error Growth up to the Planetary Scale Using a Stochastic Convection Scheme, *Mon. Weather Rev.*, 147, 1713–1731, <https://doi.org/10.1175/MWR-D-18-0292.1>, 2019.
- Bechtold, P., Köhler, M., Jung, T., Doblas-Reyes, F., Leutbecher, M., Rodwell, M. J., Vitart, F., and Balsamo, G.: Advances in simulating atmospheric variability with the ECMWF model: From synoptic to decadal time-scales, *Q. J. Roy. Meteorol. Soc.*, 134, 1337–1351, <https://doi.org/10.1002/qj.289>, 2008.
- Berman, J. D. and Torn, R. D.: The Impact of Initial Condition and Warm Conveyor Belt Forecast Uncertainty on Variability in the Downstream Waveguide in an ECWMF Case Study, *Mon. Weather Rev.*, 147, 4071–4089, <https://doi.org/10.1175/MWR-D-18-0333.1>, 2019.
- Berman, J. D. and Torn, R. D.: The Sensitivity of Downstream Ridge Building Forecasts to Upstream Warm Conveyor Belt Forecast Uncertainty Using MPAS, *Mon. Weather Rev.*, 150, 2573–2592, <https://doi.org/10.1175/MWR-D-21-0048.1>, 2022.
- Berry, G., Reeder, M. J., and Jakob, C.: A global climatology of atmospheric fronts, *Geophys. Res. Lett.*, 38, L04809, <https://doi.org/10.1029/2010GL046451>, 2011.
- Binder, H., Boettcher, M., Joos, H., and Wernli, H.: The Role of Warm Conveyor Belts for the Intensification of Extratropical Cyclones in Northern Hemisphere Winter, *J. Atmos. Sci.*, 73, 3997–4020, <https://doi.org/10.1175/JAS-D-15-0302.1>, 2016.
- Booth, J., Thompson, L., Patoux, J., and Kelly, K.: Sensitivity of Midlatitude Storm Intensification to Perturbations in the Sea Surface Temperature near the Gulf Stream, *Mon. Weather Rev.*, 140, 1241–1256, <https://doi.org/10.1175/MWR-D-11-00195.1>, 2012.
- Booth, J., Dunn-Sigouin, E., and Pfahl, S.: The Relationship Between Extratropical Cyclone Steering and Blocking Along the North American East Coast, *Geophys. Res. Lett.*, 44, 11976–11984, <https://doi.org/10.1002/2017GL075941>, 2017.
- Browning, K. A.: Organization of Clouds and Precipitation in Extratropical Cyclones, 129–153, BAMS, Boston, MA, [https://doi.org/10.1007/978-1-944970-33-8\\_8](https://doi.org/10.1007/978-1-944970-33-8_8), 1990.
- Browning, K. A. and Roberts, N. M.: Structure of a frontal cyclone, *Q. J. Roy. Meteorol. Soc.*, 120, 1535–1557, <https://doi.org/10.1002/qj.49712052006>, 1994.
- Browning, K. A., Hardman, M. E., Harrold, T. W., and Pardee, C. W.: The structure of rainbands within a midlatitude depression, *Q. J. Roy. Meteorol. Soc.*, 99, 215–231, <https://doi.org/10.1002/qj.49709942002>, 1973.
- Bui, H. and Spengler, T.: On the Influence of Sea Surface Temperature Distributions on the Development of Extratropical Cyclones,

- J. Atmos. Sci., 78, 1173–1188, <https://doi.org/10.1175/JAS-D-20-0137.1>, 2021.
- Buizza, R., Milleer, M., and Palmer, T. N.: Stochastic representation of model uncertainties in the ECMWF ensemble prediction system, *Q. J. Roy. Meteorol. Soc.*, 125, 2887–2908, <https://doi.org/10.1002/qj.49712556006>, 1999.
- Büeler, D., Ferranti, L., Magnusson, L., Quinting, J. F., and Grams, C. M.: Year-round sub-seasonal forecast skill for Atlantic–European weather regimes, *Q. J. Roy. Meteorol. Soc.*, 147, 4283–4309, <https://doi.org/10.1002/qj.4178>, 2021.
- Carlson, T. N.: Airflow Through Midlatitude Cyclones and the Comma Cloud Pattern, *Mon. Weather Rev.*, 108, 1498–1509, [https://doi.org/10.1175/1520-0493\(1980\)108<1498:ATMCAT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<1498:ATMCAT>2.0.CO;2), 1980.
- Charney, J. G.: The Dynamics of long waves in a baroclinic westerly current, *J. Atmos. Sci.*, 4, 136–162, [https://doi.org/10.1175/1520-0469\(1947\)004<0136:TDOLWI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1947)004<0136:TDOLWI>2.0.CO;2), 1947.
- Christ, S.: The influence of the sea surface temperature gradient in the Gulf Stream region on cyclone development, warm conveyor belt ascent, and downstream ridge formation during the European winter “heat wave” in February 2019, M.Sc. thesis, University Hamburg, Germany, <https://bwsyncandshare.kit.edu/s/q6Nk6aFamdfaaT5> (last access: 3 January 2025), 2023.
- Christ, S., Oertel, A., Wentz, M., and Grams, C. M.: Research data for “Understanding the sea surface temperature impact on an atmospheric blocking event using sensitivity experiments with the ICOSahedral Nonhydrostatic (ICON) model”, Karlsruhe Institute of Technology [data set], <https://doi.org/10.35097/2ps96kskhd64de0g>, 2024.
- Czaja, A., Frankignoul, C., Minobe, S., and Vanni ere, B.: Simulating the Midlatitude Atmospheric Circulation: What Might We Gain From High-Resolution Modeling of Air-Sea Interactions?, *Curr. Clim. Change Rep.*, 5, 390–406, <https://doi.org/10.1007/s40641-019-00148-5>, 2019.
- Dacre, H. F., Mart inez-Alvarado, O., and Mbengue, C. O.: Linking Atmospheric Rivers and Warm Conveyor Belt Airflows, *J. Hydrometeorol.*, 20, 1183–1196, <https://doi.org/10.1175/JHM-D-18-0175.1>, 2019.
- Dacre, H. F., Josey, S. A., and Grant, A. L. M.: Extratropical-cyclone-induced sea surface temperature anomalies in the 2013–2014 winter, *Weather Clim. Dynam.*, 1, 27–44, <https://doi.org/10.5194/wcd-1-27-2020>, 2020.
- Demirdjian, R., Doyle, J. D., Finocchio, P. M., and Reynolds, C. A.: Preconditioning and Intensification of Upstream Extratropical Cyclones through Surface Fluxes, *J. Atmos. Sci.*, 80, 1499–1517, <https://doi.org/10.1175/JAS-D-22-0251.1>, 2023.
- Doms, G., Forstner, J., Heise, E., Reinhardt, T., Ritter, B., and Schrodin, R.: A Description of the Nonhydrostatic Regional COSMO Model, [http://www.cosmo-model.org/content/model/documentation/newsLetters/newsLetter01/newsLetter\\_01.pdf](http://www.cosmo-model.org/content/model/documentation/newsLetters/newsLetter01/newsLetter_01.pdf) (last access: December 2023), 2018.
- Eckhardt, S., Stohl, A., Wernli, H., James, P., Forster, C., and Spichtinger, N.: A 15-Year Climatology of Warm Conveyor Belts, *J. Climate*, 17, 218–237, [https://doi.org/10.1175/1520-0442\(2004\)017<0218:AYCOWC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0218:AYCOWC>2.0.CO;2), 2004.
- Good, S., Fiedler, E., Mao, C., Martin, M. J., Maycock, A., Reid, R., Roberts-Jones, J., Searle, T., Waters, J., While, J., and Worsfold, M.: The Current Configuration of the OSTIA System for Operational Production of Foundation Sea Surface Temperature and Ice Concentration Analyses, *Remote Sensing*, 12, 720, <https://doi.org/10.3390/rs12040720>, 2020.
- Grams, C. M. and Archambault, H. M.: The Key Role of Diabatic Outflow in Amplifying the Midlatitude Flow: A Representative Case Study of Weather Systems Surrounding Western North Pacific Extratropical Transition, *Mon. Weather Rev.*, 144, 3847–3869, <https://doi.org/10.1175/MWR-D-15-0419.1>, 2016.
- Grams, C. M., Wernli, H., B otcher, M.,  ampa, J., Corsmeier, U., Jones, S. C., Keller, J. H., Lenz, C.-J., and Wiegand, L.: The key role of diabatic processes in modifying the upper-tropospheric wave guide: a North Atlantic case-study, *Q. J. Roy. Meteorol. Soc.*, 137, 2174–2193, <https://doi.org/10.1002/qj.891>, 2011.
- Grams, C. M., Beerli, R., Pfenninger, S., Staffell, I., and Wernli, H.: Balancing Europe’s wind-power output through spatial deployment informed by weather regimes, *Nat. Clim. Change.*, 7, 557–562, <https://doi.org/10.1038/nclimate3338>, 2017.
- Grams, C. M., Magnusson, L., and Madonna, E.: An atmospheric dynamics perspective on the amplification and propagation of forecast error in numerical weather prediction models: A case study, *Q. J. Roy. Meteorol. Soc.*, 144, 2577–2591, <https://doi.org/10.1002/qj.3353>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Hor anyi, A., Mu oz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., H olm, E., Janiskov a, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Th epaut, J.-N.: The ERA5 global reanalysis, *Q. J. Roy. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *J. Adv. Model. Earth Syst.*, 10, 1990–2008, <https://doi.org/10.1029/2018MS001364>, 2018.
- Hoskins, B. J., McIntyre, M. E., and Robertson, A. W.: On the use and significance of isentropic potential vorticity maps, *Q. J. Roy. Meteorol. Soc.*, 111, 877–946, <https://doi.org/10.1002/qj.49711147002>, 1985.
- Hotta, D. and Nakamura, H.: On the Significance of the Sensible Heat Supply from the Ocean in the Maintenance of the Mean Baroclinicity along Storm Tracks, *J. Climate*, 24, 3377–3401, <https://doi.org/10.1175/2010JCLI3910.1>, 2011.
- Joos, H. and Forbes, R. M.: Impact of different IFS microphysics on a warm conveyor belt and the downstream flow evolution, *Q. J. Roy. Meteorol. Soc.*, 142, 2727–2739, <https://doi.org/10.1002/qj.2863>, 2016.
- Joos, H. and Wernli, H.: Influence of microphysical processes on the potential vorticity development in a warm conveyor belt: a case-study with the limited-area model COSMO, *Q. J. Roy. Meteorol. Soc.*, 138, 407–418, <https://doi.org/10.1002/qj.934>, 2012.
- Joos, H., Sprenger, M., Binder, H., Beyerle, U., and Wernli, H.: Warm conveyor belts in present-day and future climate simulations – Part I: Climatology and impacts, *Weather Clim. Dynam.*, 4, 133–155, <https://doi.org/10.5194/wcd-4-133-2023>, 2023.
- Kautz, L.-A., Martius, O., Pfahl, S., Pinto, J. G., Ramos, A. M., Sousa, P. M., and Woollings, T.: Atmospheric blocking and weather extremes over the Euro-Atlantic



- sector – a review, *Weather Clim. Dynam.*, 3, 305–336, <https://doi.org/10.5194/wcd-3-305-2022>, 2022.
- Kendon, M., Sexton, D., and McCarthy, M.: A temperature of 20 °C in the UK winter: a sign of the future?, *Weather*, 75, 318–324, <https://doi.org/10.1002/wea.3811>, 2020.
- Kwon, Y.-O., Alexander, M. A., Bond, N. A., Frankignoul, C., Nakamura, H., Qiu, B., and Thompson, L. A.: Role of the Gulf Stream and Kuroshio–Oyashio Systems in Large-Scale Atmosphere–Ocean Interaction: A Review, *J. Climate*, 23, 3249–3281, <https://doi.org/10.1175/2010JCLI3343.1>, 2010.
- Leach, N. J., Weisheimer, A., Allen, M. R., and Palmer, T.: Forecast-based attribution of a winter heatwave within the limit of predictability, *PNAS*, 118, e2112087118, <https://doi.org/10.1073/pnas.2112087118>, 2021.
- Liu, X., Ma, X., Chang, P., Jia, Y., Fu, D., Xu, G., Wu, L., Saravanan, R., and Patricola, C. M.: Ocean fronts and eddies force atmospheric rivers and heavy precipitation in western North America, *Nat. Commun.*, 12, 1268, <https://doi.org/10.1038/s41467-021-21504-w>, 2021.
- Lott, F. and Miller, M. J.: A new subgrid-scale orographic drag parametrization: Its formulation and testing, *Q. J. Roy. Meteorol. Soc.*, 123, 101–127, <https://doi.org/10.1002/qj.49712353704>, 1997.
- Madonna, E., Wernli, H., Joos, H., and Martius, O.: Warm Conveyor Belts in the ERA-Interim Dataset (1979–2010). Part I: Climatology and Potential Vorticity Evolution, *J. Climate*, 27, 3–26, <https://doi.org/10.1175/JCLI-D-12-00720.1>, 2014.
- Michel, C. and Rivière, G.: The Link between Rossby Wave Breakings and Weather Regime Transitions, *J. Atmos. Sci.*, 68, 1730–1748, <https://doi.org/10.1175/2011JAS3635.1>, 2011.
- Michelangeli, P.-A., Vautard, R., and Legras, B.: Weather Regimes: Recurrence and Quasi Stationarity, *J. Atmos. Sci.*, 52, 1237–1256, [https://doi.org/10.1175/1520-0469\(1995\)052<1237:WRRASQ>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1237:WRRASQ>2.0.CO;2), 1995.
- Nakamura, H., Sampe, T., Tanimoto, Y., and Shimpo, A.: Observed Associations Among Storm Tracks, Jet Streams and Midlatitude Oceanic Fronts, 329–345, *AGU*, <https://doi.org/10.1029/147GM18>, 2004.
- Oertel, A., Sprenger, M., Joos, H., Boettcher, M., Konow, H., Hagen, M., and Wernli, H.: Observations and simulation of intense convection embedded in a warm conveyor belt – how ambient vertical wind shear determines the dynamical impact, *Weather Clim. Dynam.*, 2, 89–110, <https://doi.org/10.5194/wcd-2-89-2021>, 2021.
- Oertel, A., Miltenberger, A. K., Grams, C. M., and Hoose, C.: Interaction of microphysics and dynamics in a warm conveyor belt simulated with the ICON model, *EGUsphere*, 2023, 1–48, <https://doi.org/10.5194/egusphere-2023-259>, 2023a.
- Oertel, A., Pickl, M., Quinting, J. F., Hauser, S., Wandel, J., Magnusson, L., Balmaseda, M., Vitart, F., and Grams, C. M.: Everything Hits at Once: How Remote Rainfall Matters for the Prediction of the 2021 North American Heat Wave, *Geophys. Res. Lett.*, 50, e2022GL100958, <https://doi.org/10.1029/2022GL100958>, 2023b.
- Orr, A., Bechtold, P., Scinocca, J., Ern, M., and Janiskova, M.: Improved Middle Atmosphere Climate and Forecasts in the ECMWF Model through a Nonorographic Gravity Wave Drag Parameterization, *J. Climate*, 23, 5905–5926, <https://doi.org/10.1175/2010JCLI3490.1>, 2010.
- O’Reilly, C. H., Minobe, S., and Kuwano-Yoshida, A.: The influence of the Gulf Stream on wintertime European blocking, *Clim. Dynam.*, 47, 1545–1567, <https://doi.org/10.1007/s00382-015-2919-0>, 2016.
- Papritz, L. and Spengler, T.: Analysis of the slope of isentropic surfaces and its tendencies over the North Atlantic, *Q. J. Roy. Meteorol. Soc.*, 141, 3226–3238, <https://doi.org/10.1002/qj.2605>, 2015.
- Papritz, L., Pfahl, S., Sodemann, H., and Wernli, H.: A Climatology of Cold Air Outbreaks and Their Impact on Air–Sea Heat Fluxes in the High-Latitude South Pacific, *J. Climate*, 28, 342–364, <https://doi.org/10.1175/JCLI-D-14-00482.1>, 2015.
- Papritz, L., Aemisegger, F., and Wernli, H.: Sources and Transport Pathways of Precipitating Waters in Cold-Season Deep North Atlantic Cyclones, *J. Atmos. Sci.*, 78, 3349–3368, <https://doi.org/10.1175/JAS-D-21-0105.1>, 2021.
- Pfahl, S., Madonna, E., Boettcher, M., Joos, H., and Wernli, H.: Warm Conveyor Belts in the ERA-Interim Dataset (1979–2010). Part II: Moisture Origin and Relevance for Precipitation, *J. Climate*, 27, 27–40, <https://doi.org/10.1175/JCLI-D-13-00223.1>, 2014.
- Pfahl, S., Schwierz, C., Croci-Maspoli, M., Grams, C. M., and Wernli, H.: Importance of latent heat release in ascending air streams for atmospheric blocking, *Nat. Geosci.*, 8, 610–614, <https://doi.org/10.1038/ngeo2487>, 2015.
- Pickl, M., Quinting, J. F., and Grams, C. M.: Warm conveyor belts as amplifiers of forecast uncertainty, *Q. J. Roy. Meteorol. Soc.*, 149, 3064–3085, <https://doi.org/10.1002/qj.4546>, 2023.
- Quinting, J. F. and Grams, C. M.: Toward a Systematic Evaluation of Warm Conveyor Belts in Numerical Weather Prediction and Climate Models. Part I: Predictor Selection and Logistic Regression Model, *J. Atmos. Sci.*, 78, 1465–1485, <https://doi.org/10.1175/JAS-D-20-0139.1>, 2021.
- Quinting, J. F., Grams, C. M., Oertel, A., and Pickl, M.: EuLerian Identification of ascending AirStreams (ELIAS 2.0) in numerical weather prediction and climate models – Part 2: Model application to different datasets, *Geosci. Model Dev.*, 15, 731–744, <https://doi.org/10.5194/gmd-15-731-2022>, 2022.
- Raschendorfer, M.: The new turbulence parameterization of LM, *Cosmo news letter no. 1*, Consortium for Small-Scale Modelling, [http://www.cosmo-model.org/content/model/documentation/newsLetters/newsLetter01/newsLetter\\_01.pdf](http://www.cosmo-model.org/content/model/documentation/newsLetters/newsLetter01/newsLetter_01.pdf) (last access: December 2023), 2018.
- Reeder, M. J., Spengler, T., and Spensberger, C.: The Effect of Sea Surface Temperature Fronts on Atmospheric Frontogenesis, *J. Atmos. Sci.*, 78, 1753–1771, <https://doi.org/10.1175/JAS-D-20-0118.1>, 2021.
- Roberts, D., Vitart, F., and Balmaseda, M. A.: Hemispheric Impact of North Atlantic SSTs in Subseasonal Forecasts, *Geophys. Res. Lett.*, 48, e2020GL091446, <https://doi.org/10.1029/2020GL091446>, 2021.
- Roberts, D., Balmaseda, M. A., Tietsche, S., and Vitart, F.: Sensitivity of ECMWF coupled forecasts to improved initialization of the ocean mesoscale, *Q. J. Roy. Meteorol. Soc.*, 148, 3694–3714, <https://doi.org/10.1002/qj.4383>, 2022.
- Schaller, N., Sillmann, J., Anstey, J., Fischer, E. M., Grams, C. M., and Russo, S.: Influence of blocking on Northern European and Western Russian heatwaves in large climate model ensembles,

- Environ. Res. Lett., 13, 054015, <https://doi.org/10.1088/1748-9326/aaba55>, 2018.
- Schemm, S.: Toward Eliminating the Decades-Old “Too Zonal and Too Equatorward” Storm-Track Bias in Climate Models, *J. Adv. Model. Earth Syst.*, 15, e2022MS003482, <https://doi.org/10.1029/2022MS003482>, 2023.
- Schemm, S., Wernli, H., and Papritz, L.: Warm Conveyor Belts in Idealized Moist Baroclinic Wave Simulations, *J. Atmos. Sci.*, 70, 627–652, <https://doi.org/10.1175/JAS-D-12-0147.1>, 2013.
- Schäfler, A. and Harnisch, F.: Impact of the inflow moisture on the evolution of a warm conveyor belt, *Q. J. Roy. Meteorol. Soc.*, 141, 299–310, <https://doi.org/10.1002/qj.2360>, 2015.
- Schäfler, A., Boettcher, M., Grams, C. M., Rautenhaus, M., Sodemann, H., and Wernli, H.: Planning aircraft measurements within a warm conveyor belt, *Weather*, 69, 161–166, <https://doi.org/10.1002/wea.2245>, 2014.
- Seifert, A.: On the parameterization of evaporation of raindrops as simulated by a one-dimensional rainshaft model, *J. Atmos. Sci.*, 65, 3608–3619, <https://doi.org/10.1175/2008JAS2586.1>, 2008.
- Small, R. J., Tomas, R. A., and Bryan, F. O.: Storm track response to ocean fronts in a global high-resolution climate model, *Clim. Dynam.*, 43, 805–828, <https://doi.org/10.1007/s00382-013-1980-9>, 2014.
- Sodemann, H., Schwierz, C., and Wernli, H.: Interannual variability of Greenland winter precipitation sources: Lagrangian moisture diagnostic and North Atlantic Oscillation influence, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2007JD008503>, 2008.
- Spensberger, C., Madonna, E., Boettcher, M., Grams, C. M., Papritz, L., Quinting, J. F., Röthlisberger, M., Sprenger, M., and Zschenderlein, P.: Dynamics of concurrent and sequential Central European and Scandinavian heatwaves, *Q. J. Roy. Meteorol. Soc.*, 146, 2998–3013, <https://doi.org/10.1002/qj.3822>, 2020.
- Sprenger, M. and Wernli, H.: The LAGRANTO Lagrangian analysis tool – version 2.0, *Geosci. Model Dev.*, 8, 2569–2586, <https://doi.org/10.5194/gmd-8-2569-2015>, 2015.
- Steinfeld, D. and Pfahl, S.: The role of latent heating in atmospheric blocking dynamics: a global climatology, *Clim. Dynam.*, 53, 6159–6180, <https://doi.org/10.1007/s00382-019-04919-6>, 2019.
- Steinfeld, D., Boettcher, M., Forbes, R., and Pfahl, S.: The sensitivity of atmospheric blocking to upstream latent heating – numerical experiments, *Weather Clim. Dynam.*, 1, 405–426, <https://doi.org/10.5194/wcd-1-405-2020>, 2020.
- Steinfeld, D., Sprenger, M., Beyerle, U., and Pfahl, S.: Response of moist and dry processes in atmospheric blocking to climate change, *Environ. Res. Lett.*, 17, 084020, <https://doi.org/10.1088/1748-9326/ac81af>, 2022.
- Teubler, F. and Riemer, M.: Dynamics of Rossby Wave Packets in a Quantitative Potential Vorticity–Potential Temperature Framework, *J. Atmos. Sci.*, 73, 1063–1081, <https://doi.org/10.1175/JAS-D-15-0162.1>, 2016.
- Tiedtke, M.: A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large-Scale Models, *Mon. Weather Rev.*, 117, 1779–1800, [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2), 1989.
- Tsopouridis, L., Spengler, T., and Spensberger, C.: Smoother versus sharper Gulf Stream and Kuroshio sea surface temperature fronts: effects on cyclones and climatology, *Weather Clim. Dynam.*, 2, 953–970, <https://doi.org/10.5194/wcd-2-953-2021>, 2021.
- Vries, H. d., Scher, S., Haarsma, R., Drijfhout, S., and Delden, A. v.: How Gulf-Stream SST-fronts influence Atlantic winter storms, *Clim. Dynam.*, 52, 5899–5909, <https://doi.org/10.1007/s00382-018-4486-7>, 2019.
- Škerlak, B., Sprenger, M., and Wernli, H.: A global climatology of stratosphere–troposphere exchange using the ERA-Interim data set from 1979 to 2011, *Atmos. Chem. Phys.*, 14, 913–937, <https://doi.org/10.5194/acp-14-913-2014>, 2014.
- Wazneh, H., Gachon, P., Laprise, R., de Vernal, A., and Tremblay, B.: Atmospheric blocking events in the North Atlantic: trends and links to climate anomalies and teleconnections, *Clim. Dynam.*, 56, 2199–2221, <https://doi.org/10.1007/s00382-020-05583-x>, 2021.
- Wenta, M., Grams, C. M., Papritz, L., and Federer, M.: Linking Gulf Stream air–sea interactions to the exceptional blocking episode in February 2019: a Lagrangian perspective, *Weather Clim. Dynam.*, 5, 181–209, <https://doi.org/10.5194/wcd-5-181-2024>, 2024.
- Wernli, H.: A Lagrangian-based analysis of extratropical cyclones. II: A detailed case-study, *Q. J. Roy. Meteorol. Soc.*, 123, 1677–1706, <https://doi.org/10.1002/qj.49712354211>, 1997.
- Wernli, H. and Davies, H. C.: A Lagrangian-based analysis of extratropical cyclones. I: The method and some applications, *Q. J. Roy. Meteorol. Soc.*, 123, 467–489, <https://doi.org/10.1002/qj.49712353811>, 1997.
- Yamamoto, A., Nonaka, M., Martineau, P., Yamazaki, A., Kwon, Y.-O., Nakamura, H., and Taguchi, B.: Oceanic moisture sources contributing to wintertime Euro-Atlantic blocking, *Weather Clim. Dynam.*, 2, 819–840, <https://doi.org/10.5194/wcd-2-819-2021>, 2021.
- Yiou, P. and Nogaj, M.: Extreme climatic events and weather regimes over the North Atlantic: When and where?, *Geophys. Res. Lett.*, 31, L07202, <https://doi.org/10.1029/2003GL019119>, 2004.
- Young, M. and Galvin, J.: The record-breaking warm spell of February 2019 in Britain, the Channel Islands, France and the Netherlands, *Weather*, 75, 36–45, <https://doi.org/10.1002/wea.3664>, 2020.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Q. J. Roy. Meteorol. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.