

Responses to the reviewers

Large impact of tiny model domain shifts for the Pentecost 2014 MCS over Germany

by Christian Barthlott and Andrew I. Barrett

April 2, 2020

Dear Editor,

This letter accompanies our revised manuscript. We are grateful for the helpful comments, and hope our revision addresses them all. Below we detail the changes made in our revision. We include the text of the review in black, our responses are in blue.

Co-Editor comments

The reviewer’s main criticism refers to the motivation of your work and its connection to the relevant literature. I agree with the reviewer that “reporting on the surprisingly large sensitivity” of your domain shifting technique is a rather weak motivation for a scientific study, and I’d ask to formulate a specific research question or hypothesis that you’re addressing in your study. For instance, if this hypothesis should be that small shifts in the domain can have huge impacts on the prediction skill, than it would be natural to discuss previous studies on the impacts of changed boundary conditions (using other techniques) and also on small (stochastic?) perturbations in the initial conditions in the introduction. Possibly this could be fit into the manuscript by shortening the detailed description of your own previous research on the MCS event. Furthermore, also the discussion in the last section of the paper could be extended along this line. For example, I’m wondering how you can conclude that “model domain shifting could be used to quantify how uncertainties in the initial and boundary conditions contribute to the predictability” without comparing the sensitivity with other techniques (the spread generated by your technique may still be smaller and thus underestimate the effect of these uncertainties, which wouldn’t be a proper quantification in this case).

It was not our original intention to introduce a new method to generate ensembles with perturbed initial/boundary conditions in operational convective-scale ensemble forecasting. Our main goals were (i) to produce a reasonable forecast of this major event by having a model domain large enough to cover all stages of the event and (ii) investigate the impact of the domain size and location (i.e. IBC/LBC) on the simulation results. Due to the large influence of these tiny changes on the simulation results, we strongly believe that this method should be tested for more cases (also with different extents of domain shifting) and other models. It may also be that the high sensitivity is a feature of days with low predictability only, which would be a useful information to have. Therefore, a more systematic evaluation is left for future work. We rephrased our introduction to better motivate this study.

We also followed the suggestion of the reviewer and conducted an additional model run on the reference domain with perturbed lateral boundary conditions. The result of this run together with the requested comments on other boundary-perturbation techniques is now included in a separate section 5 at the end of the paper.

Reviewer 1

Major Comments

1. Motivation

The technique used by the authors is new to the literature on convection-allowing ensemble forecasts. However, there is no clear motivation as to why this technique is preferred over other boundary-perturbation techniques (e.g., Torn et al. 2006). Both I and Reviewer #2 saw this as a flaw in the previous manuscript and it has not been addressed within the introduction. I do not expect the authors to complete companion analyses using other techniques to generate ensemble forecasts (although I

would like to see those comparisons), but I do expect the authors to at the very least discuss the advantages and disadvantages of the various techniques (including their own) which motivates this work. There is no reason to reinvent the wheel in ensemble generation unless sufficient reasoning motivates a more simple technique. What benefit does this technique have over others that make it worthwhile to pursue?

The technique of domain shifting is not new, we already mentioned the study of Henneberg et al. (2018) in the introduction. However, in their study, the simulation domain was considerably smaller and the perturbations were introduced by shifting the domain boundaries by ten to 30 grid points north and eastwards. Our study is the first one to address all 8 directions by just one grid point on a large model domain.

It was not our original intention to introduce a new method to generate ensembles with perturbed initial/boundary conditions in operational convective-scale ensemble forecasting. Our main goals were (i) to produce a reasonable forecast of this major event by having a model domain large enough to cover all stages of the event and (ii) investigate the impact of the domain location (i.e. IBC/LBC) on the simulation results. Due to the large influence of these tiny changes on the simulation results, we strongly believe that this method should be tested for more cases (also with different extents of domain shifting) and other models. It may also be that the high sensitivity is a feature of days with low predictability only, which would be a useful information to have. Therefore, a more systematic evaluation is left for future work.

We also followed the suggestion further below and conducted an additional model run on the reference domain with perturbed lateral boundary conditions. The result of this run together with the requested comments on other boundary-perturbation techniques is now included in a separate section 5 at the end of the paper.

Changes to paper:

We rephrased our introduction at several places to better motivate this study and included a new section 5 towards the end of the paper. Parts of the introduction were also moved to this new section, please see the track-changed version for all changes.

One possible new avenue to pursue is comparing the conducted experiments with parallel forecasts that use small, stochastic perturbations on the boundary conditions of the reference forecast. I suspect that the forecast variability seen in the current forecasts is simply the result of numerical noise and chaos seeding (see Ancell et al. 2018), whereby small perturbations in remote locations of the domain (e.g., the boundary) can induce very large changes in forecast precipitation for unrealistic reasons; the perturbations propagate during each model timestep the distance of the finite differencing scheme, which is typically faster than the speed of sound. The authors could actually use this companion analysis to prove or disprove that their methodology is sound and actually producing meaningful spread in their forecasts, i.e. represents true predictability.

We thank the reviewer for this suggestion and performed another model run on the reference domain, but with stochastic perturbations on the boundary conditions. We added random temperature fluctuations with a Gaussian distribution of zero mean and a standard deviation of 0.01 K at all levels for the model boundaries. The results of this run together with comments on other perturbation techniques is now in a new section 5. In this paper, we wanted to demonstrate the large sensitivity of the model results to a relatively simple technique of perturbing the initial and boundary conditions. The applicability of this technique has to be tested for other (and more) cases and could then also be compared to other operational used methods. Thus, we mentioned a number of other methods in the new section, but did not provide a detailed intercomparison of different methods or rating of our method against others. We believe that this would blow up the paper unnecessarily and should be

done in the mentioned follow-up work.

Changes to paper:

New section 5 and modifications in the Summary/Conclusion section

2. Ensemble Sensitivity

My previous suggestion to use ensemble sensitivity was motivated by wanting to see that the reasoning for forecast differences proposed by the authors (e.g., closer proximity to cold SSTs) could be deduced statistically, and not just subjectively by the authors. For instance, does higher precipitation in Germany during the 1700-0000 UTC timeframe (Fig. 4) statistically relate to higher/lower CAPE, stronger convergence, or upper-level dynamics at an earlier time that can explain the differences in the simulations? The analysis presented is a good start, but maybe insufficient to answer the dynamic questions about what is influencing the forecasts. I suggest the authors consider applying ensemble sensitivity to the fields in Figure 8 to support their claims about what ultimately influences MCS initiation and propagation into Germany.

We thank the reviewer for clarifying the motivation behind the suggestion of ensemble sensitivity analysis. Following their suggestion, Figure 9 has been replaced with a new figure looking at some of the variables from Figure 8 and now with a consistent time range to Figure 10. As is now shown, CAPE and precipitation rate do correlate well with later success of the simulation. However, the ensemble sensitivity analysis cannot be used to describe meaningful differences in the simulations before the convection has formed and organised. Only once it is organised is the sensitivity to CAPE and precipitation rate clear. The initial causes of the different tracks of the convective cells remain unclear but are likely chaotic in origin.

Most variables in Figure 8 only show a difference between successful and unsuccessful simulations at and after 12 UTC. CAPE and precipitation rate are examples of these - and the sensitivity is shown clearly in Figure 9. Low-level convergence and the resulting vertical velocity at 500 m both show differences before 12 UTC; however, Figure 9 shows that the location of these differences are not systematic in the set of simulations as the ensemble sensitivity analysis is largely dominated by noise until after 12 UTC.

Ensemble sensitivity analysis fields of CAPE and CIN were very similar and CIN is therefore omitted. Similarly, precipitable water is very similar to precipitation rate in highlighting the location of the existing convective cell. There really are no large-scale differences in e.g. CAPE, which can also be seen in Figure 10. We believe that the current ensemble sensitivity analysis has proved valuable in highlighting systematic differences between the simulations, but that further analysis will not allow the origin of the differences to be uncovered - largely because of the disparate locations of the initial convection.

Changes to paper:

Figure 9 has been replaced and improved by using different variables consistent with Figure 8. The text in section 4.4 has been modified consistently with the new figure.

3. Forecast Differences

It may be helpful for the eventual readership to actually see forecast difference plots that illustrate the reduced CAPE near the french coastline, enhanced convergence, etc. rather than try to interpret the differences based on equivalent environmental plots. For instance, in Figure 10 the differences between the W, REF, and SE precipitation rates are fairly obvious, but differences in the CAPE fields (and presumably a number of other environmental parameters) are much more difficult to make out. Rather than showing identical forecast plots side-by-side as in Figs. 10, 11, and 12 (and elsewhere in the manuscript), please consider including different plots, which would help elucidate the claims made

in the text more clearly.

We do not believe that difference plots would help the reader much. As the reviewer already mentioned, precipitation differences are fairly obvious in Fig. 10. At that time of convective stage, there are no differences in CAPE between individual model runs (only after 1200 UTC differences can be seen as a cause of convection/no convection). The important point is the location of the cell which is more north in areas of lower CAPE for the unsuccessful runs. Figure 12 is a collection of plots from the same model run for different times, so difference plots cannot be applied here.

Changes to paper:

none

Minor/Specific Comments

- Line 33, Line 36: What is the convective event on 8 June? The only mention of deep convection is the MCS on 9 June (Line 27). Please consider highlighting the preceding day or remove mention of an event on 8 June.

We agree with the reviewer that the events on June 8 are not essential for the reader and removed the results but kept the mentioning of the day.

Changes to paper:

A series of different numerical simulations for the convective events of ~~8 and~~ 9 June 2014 *and the previous day* were performed with the COSMO model, the main findings were

- The COSMO model (in quasi operational set-up, without data assimilation) initialized at 00:00 UTC did not reproduce ~~the events on 8 June only, but not~~ the mesoscale convective system (MCS) on 9 June.
 - The enlargement of the model domain towards the West ~~had the largest effect~~ *improved the precipitation forecast over France only* due to better resolving the initiation and development of deep convection over western France and, later, secondary initiation over northern France. *The MCS over Germany, however, was not simulated even with this larger domain.*
 - ~~The use of a double-moment microphysical scheme improved the rainfall amounts on 8 June only.~~
- Lines 49-51: But this increase in lead time was not observed in the simulations by Barthlott et al. (2017), correct? Lines 37-38 only mention that extending the model domain “had the largest effect”. Please clarify what “largest effect” means in relation to the forecast skill and if this skill improvement corroborates the statement in Lines 49-51.

The increase in lead time mentioned in lines 49-51 is just of theoretical nature if we assume that a convection-permitting model runs over a larger domain and successfully simulates the convection to come. In the study of Barthlott et al. (2017), the enlargement of the domain had a positive impact only for the preceding day and not for the day with the MCS discussed here. We modified the text to make that clearer.

Changes to paper:

The enlargement of the model domain towards the West ~~had the largest effect~~ *improved the precipitation forecast over France only* due to better resolving the initiation and development of deep convection over western France and, later, secondary initiation over northern France. *The MCS over Germany on 9 June 2014, however, was not simulated even with this larger domain.*

- Line 55: “there are various to generate”

We believe that our phrasing is correct here: *“There a various ways of generating an ensemble...”*

Changes to paper:

none

- Line 59: “Recent studies by”

done

- Lines 59-61: How do these studies contribute to ensemble generation (Line 56)? These references seem out of place for what this section is discussing.

There are various ways of generating an ensemble and some of them are cited in that parapgraph. The studies on lines 59-61 address other and new approaches in ensemble modelling and are therefore mentioned here. Please note that these lines are now in section 5.

Changes to paper:

none

- Lines 7,94,182: I urge the authors to consider removing “surprising” and “surprisingly” from the manuscript. The results are NOT surprising when you consider the domain shifting technique as a simple stochastic perturbation, which has been shown to have large impacts on convection by others when considering perturbed lateral boundary conditions (e.g., Clark et al. 2010, Romine et al. 2014). This wording is much too subjective and not scientific. Moreover, the general variability of convective systems can be dramatic within convection-allowing models, as demonstrated in other studies (e.g., Melhauser and Zhang 2012), so I would not consider the current results “surprising”.

We do believe that our technique of domain shifting of just 1 grid point provides surprising or at least unexpected results for this particular case. Given the large model domain and the minor changes at the boundaries, we would not have anticipated such a large dependency. However, we follow the reviewer’s suggestion and eliminate that wording.

Nevertheless, we do believe that scientists can be surprised. A keyword search for “surprising” in the AMS journals of the last 10 years leads to more than 7000 hits.

Changes to paper:

Abstract and Introduction: ~~surprisingly~~

Section 4.2 and Summary: ~~surprising~~ → *unexpected*

- Lines 109-110: What initiated the storms during the night and morning hours? Shortwave trough?

Please see our reply to the fourth subsequent remark. “The showers near Nantes...”

- Line 129: Please specify the date with the initialization time

done

- Line 148: I think this is the first time the authors have mentioned “reference run”. I would suggest highlighting the reference/control simulation in the previous section, maybe stating that before any domain shifting occurs, you run a reference/control simulation of the event to compare shifted experiments against. Section 4 then naturally transitions to the discussion of reference forecast skill.

This is a good suggestion. We now introduce the reference run already in section 3.2.

Changes to paper:

Section 3.2:

At first, we conducted a reference run (REF) with a model domain containing ~~The model domain contains~~ 600×650 grid points, which corresponds to an area of about 1668 km×1807 km.

We also included the reference run in Table 1.

Table 1. *Domain shifting* ~~Overview of the numerical simulations.~~

Section 4.1:

Here we compare our *reference* simulation...

- Line 159: “of of” to “of”

done

- Lines 169-170: What initiated the showers near Nantes? What was the initiating mechanism that ultimately led to the extreme event in Germany later in the forecast period?

The showers near Nantes during the night were initiated by a combination of large-scale forcing and low-level wind convergence. Fig. R.1 shows low-level winds and Q-vector divergence between 0100 UTC and 0230 UTC. Although CAPE is absent, there is a region with Q-vector divergence (negative values) indicating forcing for upward vertical motion which coincides with low-level wind convergence. We added additional information in the text.

Later on, more weak showers formed near the coast which traveled towards northern France. One of these showers then developed into the MCS as sufficient CAPE was present in the afternoon and the wind shear favoured organized long-lived convection by outflow triggering. The proximity of the track of the convective system to the north coast of France and the colder temperatures over the sea than the land then determine the large differences in the model results. These processes are analyzed in detail in section 4.6.

Changes to paper:

“The precursors of these showers were initiated over the sea by a combination of large-scale forcing (determined by Q-vector divergence) and low-level wind convergence (not shown).”

- Line 170: How was this track determined? Subjectively? Please specify

Yes, the track was subjectively determined. We included that information in the text.

Changes to paper:

The *subjectively determined* track of the system...

- Line 174: “too far north”

done

Changes to paper:

~~too far in the North~~ *too far north*

- Lines 184-187: These two sentences are at odds with one another. The first sentence states that there are no systematic responses to the forecasts when shifting the domain in a specific direction. The second sentence actually describes a systematic response resulting in poor forecasts when shifting to the east. Please rectify this discrepancy. Please switch sections 4.2 and 4.3. It makes more sense to discuss the reference forecast, the sensitivity to domain choice, and then the results of the experiment simulations. It reads disjointed as is.




Figure R.1: Left: CAPE (colour shading), precipitation (blue colour shading), and 10-m wind field (arrows); Right: Q-vector divergence (colours), 500 hPa geopotential height (contour lines), and precipitation rate (hatched). From top to bottom: 0100, 0130, 0200, 0230 UTC.

We deleted the second sentence as the E run did produce precipitation over Germany, but somewhat shifted northwards. The statement from the first sentence is valid, i.e. that there is no systematic response to the forecasts when shifting the domain in a specific direction.

We believe that it is not necessary to switch sections 4.2 and 4.3. We first want to show the sensitivity of convective precipitation to domain choice (4.2) and then find possible reasons for this behavior (4.3 Differences in IC, 4.4 Convection-related parameters,...). We therefore decided to keep the sections in this order.

Changes to paper:

~~However, as the precipitation in the E run is more to the North and does not extend as much to the East as the other successful runs, all the eastward shift simulations have poorer prediction.~~

- Lines 210-211: Please reference relevant literature on this topic (e.g., Ancell et al. 2018 and others) to support your claim.

We decided to remove that sentence, as it is rather speculative and we show later that small

displacements of the precursor cell determines whether a large MCS develops or not.

Changes to paper:

~~But given the chaotic nature of the atmosphere in convective weather events and the nonlinearity of the system with many feedbacks involved, these small deviations can determine whether a large MCS develops or not.~~

- Line 219: Please illustrate this box in one of your figures. A reader should be able to see/visualize where this box is placed at a particular time for the analysis presented in this section.

As we already depicted in the text, the box follows the storm along the path depicted in Fig. 6. So for different times, the box has different locations. Moreover, the location of the box would only be meaningful for 30-min rain rates and not for accumulated precipitation. It is not feasible to display all boxes for all times in the same figure (Fig. R.2):




Figure R.2: 24-h accumulated precipitation with boxes for all times (left) and every 6 hours (right).

Changes to paper:

We included the location of 4 boxes corresponding to four times (06:00, 12:00, 18:00, 24:00 UTC) in Fig. 6e.

- Line 225: I do not see the “slight reduction in intensity” within the Fig. 8a black line at 1700 UTC.

You are right, we re-phrased that sentence.

Changes to paper:

“The precipitation rate of the REF run is gradually increasing until 14:30 UTC (Fig. 8a) and remains more or less constant until 17:00 UTC. Then precipitation rate is strongly increasing until 18:30 UTC followed by a slight reduction in intensity before the maximum is reached between 21:00 UTC and 22:00 UTC.”

- Line 225: Between 2100 and 2200 UTC
done
- Lines 231-234: There is a clear bifurcation in directional shear between good and bad experiment forecasts by 1300 UTC. Can the authors please discuss the possible implications of these directional shear differences as they relate to MCS development and propagation?

Both deep layer shear and directional shear are similar in all model runs until 12:00 UTC and show suitable conditions for highly-organized convection. It is only after convection has intensified in the successful runs (12:00–13:00 UTC), that there is a bifurcation of directional shear and speed shear between the good and bad experiment forecasts. Therefore we believe that these differences are a result of the (non)-existent MCS itself and not the driver for it. In all model runs, DLS is higher than 20 m/s until 21:00 UTC which shows the potential for organized convection for all model runs. But because of the processes discussed later in the paper (initial convection too close to colder sea), not all runs simulate the MCS. We further comment on that later in a similar question.

Changes to paper:

none

- Lines 245-246: This is a reasonable assumption...but can the authors actually prove this is what is occurring and not some other mesoscale process?

Our assumption that precipitation is consuming CAPE is further outlined below in similar questions (Lines 265-266/Line 269) to that topic.

Changes to paper:

none

- Line 250: What is the implication of reference simulation and “unsuccessful” runs having similar proportions of CAPE-to-CIN? It appears to me that the absence of CIN is the major contributor to initiation for the “successful” (warmer colors) runs.




Figure R.3: Time series of domain-averaged CIN.

CIN or the absence of CIN is not the major contributor for dividing into successful and not successful runs. We computed domain-averaged CIN (Fig. R.3). It can be seen that until 1100–1200 UTC, CIN is on average similar in all model runs. Differences in CIN emerge between 1300–1600 UTC when convective precipitation is intensifying. We added a remark in the paper.

Changes to paper:

“The fraction of grid points fulfilling that criterion is primarily dominated by the existence of CAPE as domain-averaged CIN is very similar in all model runs until 12:00 UTC (not shown).”

- Lines 265-266: Can the authors show this is indeed the cause of lower CAPE and shear in the successful runs? It reads as speculative and should be backed up with some analysis, at the very least shown to the reviewers

Please see our response to the next question, which covers the same topic.

- Line 269: Can the authors please provide supporting evidence that indeed the lower CAPE and shear is a result of storm modification? This appears to be speculation, and should be corroborated with evidence.

As can be seen in the time series of Fig. 8, CAPE is building up from the early morning hours in all model runs. The runs without the MCS simulate an increase in CAPE until 1700–1800 UTC, after that CAPE is reduced due to a reduction of near-surface temperatures. The precipitation in the successful runs, however, consume that CAPE. The model also simulates a time lag of 2-3 h between the CAPE and precipitation with former leading the latter, which is typical for convection over land. It is also evident from Fig. 10 that convective precipitation consumes CAPE, as in the wake of the MCS, CAPE is significantly reduced. This is also evident from Fig. R.4 in which precipitation is displayed as a function of CAPE. The not successful W run simulates an increase in CAPE until 1930 UTC, then CAPE is reduced due to the reduction of near-surface temperatures. The reference run, on the other hand, simulates an increase in CAPE until 1730 UTC only. Then, CAPE is consumed by the precipitation of the convective system. We believe that this is enough evidence for the cause of lower CAPE in the successful runs.




Figure 9 now does show the sensitivity to some of the fields shown in Figure 8. The text in section 4.4 is changed accordingly.

- Line 290: Please show this box in a figure

Changes to paper:

Box added to Figure 6e.

References

Ancell, B. C., et al, 2018: Seeding Chaos: The Dire Consequences of Numerical Noise in NWP Perturbation Experiments. *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-17-0129.1>.

Clark et al, 2010: Growth of Spread in Convection-Allowing and Convection-Parameterizing Ensembles, *Weather and Forecasting*, 25, 594-612.

Melhauser, C. and F. Zhang, 2012: Practical and Intrinsic Predictability of Severe and Convective Weather at the Mesoscales. *Journal of Atmospheric Sciences*, 69, 3350-3371.

Romine, G. et al., 2014: Representing Forecast Error in a Convection-Permitting Ensemble System. *Monthly Weather Review*, 142, 4519-4541.

Torn. R., G. J. Hakim, and C. Snyder, 2006: Boundary conditions for limited-area ensemble Kalman filters. *Monthly Weather Review*, 134, 2490-2502.

Large impact of tiny model domain shifts for the Pentecost 2014 MCS over Germany

Christian Barthlott and Andrew I. Barrett

Institute of Meteorology and Climate Research (IMK-TRO), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

Correspondence: Christian Barthlott (christian.barthlott@kit.edu)

Abstract. The mesoscale convective system (MCS) that affected Germany at Pentecost 2014 (9 June 2014) was one of the most severe for decades. However, the predictability of this system was very low as the operational deterministic and ensemble prediction systems **completely** failed to predict the event with ~~sufficiently long lead times~~ **more than 12 hours lead time**. We present hindcasts of the event using the COnsortium for Small-scale MOdeling (COSMO) model at convection-permitting (2.8 km) resolution on a large (1668×1807 km) ~~grid, which domain~~. **Using this large domain** allowed us to **successfully** simulate the whole life cycle of the system originating from the French Atlantic coast. ~~Results show that this model configuration successfully reproduces the convective events of that day. However, the low~~ **However, even with the large domain the** predictability of the ~~event was evident by the surprisingly large impact of tiny~~ **MCS is low. Tiny** changes to the model domain ~~·~~ **We produced large changes to the MCS, removing it completely from some simulations. To demonstrate this we** systematically shifted the model domain by **just** one grid point in eight different directions, from which three did not simulate any convection over Germany. ~~The analyses show that~~ **Our analysis shows that there were** no important differences in domain-averaged initial conditions nor in the preconvective environment ahead of the convective system ~~exist. That.~~ **The main reason that** one-third of these seemingly identical initial conditions fails to produce any convection over Germany ~~is intriguing. The main reason for the different model results~~ seems to be the proximity of the track of the initial convective system to the coast and colder sea surface. The COSMO model simulates small horizontal displacements of the precursors of the MCS which then determine if the cells dissipate close to the sea or reach a favorable area for convective development over land and further evolve into an MCS. This study demonstrates the potentially huge impact of tiny model domain shifts on forecasting convective processes in this case, which suggests that the sensitivity to similarly small initial condition perturbations **could be a helpful indicator of days with low predictability and** should be evaluated across other cases, model and weather regimes.

20 *Copyright statement.* TEXT

1 Introduction

An accurate forecast of deep moist convection is of great societal and economic relevance due to multiple risks from heavy precipitation, strong winds, lightning, or hail. Convection-permitting models have provided a step-change in rainfall forecasting

and are used operationally in many parts of the world (Clark et al., 2016). Although progress has been made through higher grid spacing of numerical weather prediction models and better parameterizations of physical processes, quantitative forecasting of convective storms remains a challenge. All forecast centers still suffer from so-called forecast busts (Rodwell et al., 2013), in which a ~~strong-large~~ drop in performance occurs and the forecast skill becomes very low.

The 2014 Pentecost storms over Germany were also partly characterized by a low forecast skill. Following a period of hot weather, a series of convective systems occurred over northwestern Germany leading to significant damages with even six fatalities. The major event took place on Pentecost Monday (9 June 2014) where a mesoscale convective system originating over France traveled across Belgium and hit northwestern Germany in the evening (Mathias et al., 2017). At the German Weather Service, both the deterministic run and all 20 members of the ensemble prediction system failed to predict any severe storms over northern Germany. These events and their poor prediction motivated the study of Barthlott et al. (2017), in which several methods of improving Consortium for Small-scale Modeling (COSMO) model simulations were evaluated, including: a larger model domain, higher grid spacing, a more sophisticated microphysics scheme and different initialization times. A series of different numerical simulations for the convective events of ~~8-and~~ 9 June 2014 ~~and the previous day~~ were performed with the COSMO model, the main findings were :-

- The COSMO model (in quasi operational set-up, without data assimilation) initialized at 00:00 UTC ~~reproduced the events on 8 June only, but not~~ did not reproduce the mesoscale convective system (MCS) on 9 June.
- ~~The enlargement of~~ Enlarging the model domain towards the West ~~had the largest effect~~ improved the precipitation forecast only over France, due to better resolving the initiation and development of deep convection over western France and, later, secondary initiation over northern France. ~~The MCS over Germany, however, was not simulated even with this larger domain.~~
- ~~Changes to~~ Improving both vertical and horizontal grid spacing (highest resolution 1 km) had only minor effects on the simulation results. ~~Use of even higher resolutions up to large-eddy scale as in Barthlott and Hoose (2015) may have helped, but was left for future work.~~
- ~~The use of a double-moment microphysical scheme improved the rainfall amounts on 8 June only.~~
- An increased or reduced initial soil moisture had significant effects on the energy balance of the surface (see e.g., Barthlott et al., 2011), but still no MCS-like system was simulated over Germany.
- Although weaker than observed, later initialization times (03:00 UTC, 06:00 UTC) produced deep convection over Germany due to outflow triggering and secondary cell initiation.

Specific reasons for the model failure of the 00:00 UTC run remained unclear, but the analysis of convection-related variables indicated too high values of convective inhibition (CIN) in northern Germany. As was pointed out by Groenemeijer (2014), extending the model domain to the west and south would **potentially** allow storms to be captured earlier by the model. By

55 enlarging the domain 300 km to the west, the direction from which most severe thunderstorms arrive, the lead time ~~can~~ could be increased by 3 h (assuming a system moving with ~~90~~100 km h⁻¹).

~~Many operational forecast centers produce both a high-resolution forecast and an ensemble of lower resolution to provide a measure of uncertainty (Rodwell et al., 2013). In recent years, the benefits of ensemble techniques over deterministic numerical weather prediction are widely recognized (e.g., Hohenegger and Schär, 2007). The recent advance to convective-scale ensembles can help address the uncertainty associated with convective-scale processes (Barrett et al., 2016). There are various ways of generating an ensemble, such as perturbations to~~ The present study was motivated by the wish to test the hypothesis that an even larger model domain would allow the MCS to be simulated. We aimed to (i) produce a reasonable hindcast of this event by making the model domain large enough to cover all stages of the ~~initial conditions and/or boundary conditions (e.g., Montani et al., 2011; Kühnlein et al., 2014), stochastic physical parameterizations (e.g., Buizza et al., 1999; Berner et al., 2017), or ensemble data assimilation such as ensemble Kalman filter (e.g., Dowell et al., 2004; Zhang et al., 2004; Reich et al., 2011). Recent studies of Schneider et al. (2019) and Keil et al. (2019) have also shown that different assumptions for the amount of cloud condensation nuclei could be included in convective-scale ensemble forecasting, but only if the model employs a double-moment microphysics scheme. Because of the fundamental uncertainties of the simulations due to nonlinearities of the model equations, several studies have noted the significant impact of initial and lateral boundary conditions on the simulation of~~ convective precipitation for some situations (e.g., Hohenegger et al., 2006; Trentmann et al., 2009; Richard et al., 2011; Bouttier and Rayn. that ensemble members with the most accurate initial and boundary conditions are most skilful at predicting the location of convective initiation (Barrett et al., 2015).

~~While trying to get a reasonable model representation of the MCS on 9 June 2014, we conducted several numerical experiments with different domain sizes and domain locations and found remarkably large differences in the simulations.~~ event and (ii) investigate what aspects of the model domain (e.g. size, position) were important to successfully simulate the MCS. For regional climate simulations, the sensitivity to the size and position of the domain chosen is well known (e.g. Miguez-Macho et al., 2004). In their study, the center of the grid was successively moved 17° to the west, 10° to the east, 7° to the north, and 10° to the south. These large changes led to a distortion of the large-scale circulation by interaction of the modeled flow with the lateral boundaries of the nested domain which sometimes had a large effect on the precipitation results. Seth and Giorgi (1998) demonstrated that the domain of a regional climate model must be carefully selected for its specific application. In particular, domains much larger than the area of interest appear to be needed for studies of sensitivity to internal forcings, as the interactions between boundary conditions and internal model forcings played an important role. ~~Similar results were obtained by Landman et al. (2005); in their regional climate simulations, the positioning of the eastern boundary of the regional model domain is of major importance in the life cycle of simulated tropical cyclone-like vortices.~~

85 Beside the influence of different domain sizes, the approach of shifting the model domain boundaries (and keeping the number of grid points constant) has been rarely used for short-range convection-resolving numerical weather prediction. The only study, to the authors' knowledge, was conducted by Henneberg et al. (2018) for examining soil moisture influences on convective precipitation over northern Germany. Perturbations were introduced by shifting the domain boundaries by ten to 30 grid points north and eastwards. Their results have shown that by shifting the model domain, an estimate ~~on~~ of the uncertainty

90 of the model results can be calculated and a ~~sufficient~~ **sufficiently** large model spread can be achieved. A ~~somehow~~ **somewhat** similar technique was used by Schlüter and Schädler (2010) to study the impact of small changes in the synoptic situations on extreme precipitation events. They shifted the large-scale atmospheric fields to north, south, east, and west with respect to the orography by about 28 and 56 km and found that the modeled precipitation can be quite sensitive to small changes of the synoptic situation with changes ~~in~~ **on** the order of 20% for the maximum daily precipitation.

95 ~~Limited-area-ensemble predictions are known to be sensitive to the specification of lateral boundary conditions (Bouttier and Raynaud, 2004). The variability of boundary conditions is essential for representing large-scale uncertainties in limited-area predictions beyond a few hours (e.g. Gebhardt et al., 2011; Vié et al., 2011). Whereas in regional climate simulations, where it is desired to eliminate the dependence of the results on the position of the domain~~ **By investigating what aspects of the model domain (size, position) were important to successfully simulate the MCS, we found that small movements of the model domain, e.g. ,** ~~by spectral nudging (Miguez-Macho et al., 2004), the simple approach of domain shifting could be used to account for errors in the initial and boundary conditions and produce an ensemble with sufficient spread. Thus, it is of interest to further evaluate this simple approach of domain shifting to account for uncertainties in~~ **50 km further west or slightly increasing all dimensions of the model domain, could separate successful simulations of the MCS from complete failures. We then systematically reduced the size of our model domain changes, eventually converging on a single grid-point (2.8 km) shift in location while keeping the**
100 ~~domain size the same. Even with these tiny changes, the representation of the~~ **initial and boundary conditions.** ~~MCS ranges from good to failure.~~

~~This paper reports on the surprisingly large sensitivity of moving the model domain by only 1 grid point~~ **It is these final simulations, which result from a single grid-point shift in the model domain, that are discussed in this paper. We evaluate what the differences between the simulations were ,** ~~and what the origin of these differences was. This gives us further insight into the important physical processes for this event, and helps understand why it was so difficult to predict in the operational forecast models. Furthermore, we explore the potential of model domain shifting to help determine the predictability of convective events within an ensemble modeling framework.~~

2 Synoptic situation and observed precipitation

To describe the synoptic situation of the event, we briefly summarize the analysis from Barthlott et al. (2017). For more details,
115 we refer to that paper and to the synoptic analysis performed by Mathias et al. (2017). The synoptic situation on 9 June 2014 was characterized by a trough stretching across the northern Atlantic Ocean southwards almost to the Canary Islands and an extensive ridge covering central northern Africa, the western Mediterranean Sea, and central Europe (Fig. 1). At the surface, there was a low pressure system named “Ela” corresponding to the upper-level trough. The high pressure system over the continent (“Wolfgang”) dominated the region between the Alps, Poland, and the Black Sea. This configuration was already
120 present on the day before and had progressed only slowly eastward. During the period of 8–10 June 2014, the temperature contrast over Western Europe intensified. Cool Atlantic air masses were present at the eastern edge of the low pressure system, while moist and very warm air of subtropical origin was carried north-eastwards by the strong upper-level south-westerly flow.




Figure 1. Global Forecast System analyses at 9 June 00:00 UTC showing 500 hPa geopotential height (gpm, shading), sea-level pressure (hPa, white contours), and 500 hPa wind barbs.




Figure 2. Meteosat visible satellite pictures of northern France and western Germany from 08:00–18:00 UTC (raw data courtesy of EUMETSAT). The convective system investigated here is marked by the blue circles.

Intense thunderstorms developed in northwestern France and the Benelux countries during the night and in the morning hours of 9 June 2014 and also later in the day due to diurnal surface heating. In the evening, an elongated area of convective storms extended from eastern Spain across western and northern France all the way to Benelux (i.e. Belgium, the Netherlands, and Luxembourg) and northwestern Germany. An intense MCS reached its mature phase in the evening over Benelux and western

Germany, which is in the focus of this study. The analysis of satellite pictures in Fig. 2 reveals that the system originated over the Bay of Biscay in the morning of 9 June. The temporal evolution was characterized by several cycles of intensification and decay. For example at 16:00 UTC (Fig. 2e), an intensification at the northeastern edge of the system took place which lead to the large MCS over Germany in the evening with overshooting tops and signs of gravity waves (Fig. 2f).

3 Method

3.1 COSMO model

All simulations were performed with version 5.3 of the numerical weather prediction model COSMO (CONsortium for Small-scale MOdeling, Schättler et al., 2016). The COSMO model is a nonhydrostatic limited-area atmospheric prediction model initially developed by the Deutscher Wetterdienst (DWD, German Weather Service) which is operationally used by several weather services in Europe. It is based on the fully compressible primitive equations integrated with a two-time level Runge-Kutta method (Wicker and Skamarock, 2002). As previous simulations of Barthlott et al. (2017) showed little sensitivity of the results to model grid spacing, we performed all simulations with 2.8 km horizontal grid spacing and 50 terrain-following vertical levels. This corresponds to the operational used setup at the DWD at the time of the event. For consistency with previous simulations of this case, the changes suggested by Barrett et al. (2019) to minimize timestep-dependent results from the microphysics parameterization were not included. The model uses an Arakawa C-grid for horizontal differencing on a rotated latitude/longitude grid. Initial and boundary conditions come from the ECMWF's Integrated Forecasting System (IFS) analyses with a resolution of 0.125 °. All simulations are initialized at 00:00 UTC on 9 June 2014 with an integration time of 36 h. The time step is set to 25 s. Deep convection is resolved explicitly and a modified Tiedtke-scheme (Tiedtke, 1989) is used to parameterize shallow convection (as did the operational deterministic and ensemble prediction system at that time). A 1D turbulence parameterization based on the prognostic equation for the turbulent kinetic energy after Mellor and Yamada (1974) is applied. No latent heat nudging or other data assimilation technique is used. Instead of the operationally used single-moment microphysics scheme, we use the double-moment scheme of Seifert and Beheng (2006) assuming continental concentrations of cloud condensation nuclei ($N_{CN} = 1700 \text{ cm}^{-3}$). In our configuration, the CCN concentration remains constant and is not varied as, for example, in the study of Barthlott and Hoose (2018) investigating aerosol effects on clouds and precipitation in central Europe.

3.2 Model domain choices

~~The model domain contains~~ At first, we conducted a reference run (REF) with a model domain containing 600×650 grid points, which corresponds to an area of about $1668 \text{ km} \times 1807 \text{ km}$. To be able to simulate the entire life cycle of the convective system, the domain covers France, Benelux, Germany, and the Alps with parts of the neighboring countries (Fig. 3). The sensitivity of the model results to domain shifting is assessed by conducting simulations where the model domain is shifted by one grid point in eight different directions (Table 1). All other model settings remained unchanged.




Figure 3. COSMO simulation domain and model orography (in m asl) of the reference run.

Table 1. ~~Domain shifting~~ Overview of the numerical simulations.

name	domain shifting
REF	none
N	1 grid point towards the North
NE	1 grid point towards the North-East
E	1 grid point towards the East
SE	1 grid point towards the South-East
S	1 grid point towards the South
SW	1 grid point towards the South-West
W	1 grid point towards the West
NW	1 grid point towards the North-West

4 Results

4.1 Reference run

160 Here we compare our ~~simulations~~ reference simulation to radar-derived precipitation from the precipitation analysis algorithm RADOLAN (Radar Online Adjustment), which combines weather radar data with hourly surface precipitation observations of




Figure 4. Radar-derived (a) and simulated (b) accumulated precipitation in mm on 9 June 2014 (17:00–24:00 UTC).

about 1300 automated rain gauges to get quality-controlled, high-resolution (1 km) quantitative precipitation estimates. In the reference run, simulated precipitation on the evening of 9 June occurs over Benelux and northern Germany (Fig. 4b). The area covered by precipitation generally agrees well with that from radar observations (Fig. 4a). However, the simulated precipitation is slightly too far north and. In areas near Cologne, Frankfurt, and south of Karlsruhe, the model produces less precipitation than observed and some single convective cells are not simulated. In contrast, precipitation covers more of the English Channel northern Netherlands and Belgium than observed. As far as the total precipitation amounts are concerned, the COSMO model produces similar values to those observed with slightly lower maximum values. However, radar is not an instrument measuring




Figure 5. Radar-derived (blue contours) and simulated (red contours) 30-min precipitation of 1 mm (solid) and 5 mm (dashed) on 9 June 2014.

precipitation in a quantitative sense (see e.g., Rossa et al., 2005) and differences in the amount do not necessarily indicate a
 170 poor performance of the model. Unfortunately, this radar composite also suffers from missing data at some locations (e.g. over
 Belgium south-west from Cologne) and also different calibrations or Z-R-relationships (obvious from the strong precipitation
 gradient about 100 km north of Cologne).

The temporal evolution of ~~of~~ the convective system from both radar-derived and simulated 30-min precipitation rates is
 presented in Fig. 5. Both systems follow a very similar track. We observe the following two main differences: (i) the model
 175 simulates the convective system too far to the North and (ii) the simulated MCS moves faster towards the East. These differences
 are similar to the simulations of Mathias et al. (2017). Moreover, the observed area covered with rain is larger than simulated.
 Given the overall good agreement in precipitation location and timing with reasonable accumulations, we conclude that the
 reference run serves as a good basis for our sensitivity studies.

4.2 Sensitivity to domain choice

180 The 24 h accumulated precipitation for the REF run and all shifted model runs is ~~displayed on the full model domain shown~~
 in Fig. 6. All model realizations show convective systems initiated near the Bay of Biscay in southwestern France which later
 move ~~in a northeasterly direction~~ ~~towards the north east~~. However, these systems are not related to the life cycle of the MCS
 that forms later over Germany and are not important for this study. The system that later became the MCS started as several




Figure 6. 24-h precipitation (00:00–24:00 UTC on 9 June) amount in mm. The red line indicates the approximated storm track of the REF run. The black boxes in (e) indicate areas for averaging convection-related variables along the storm path for 06:00, 12:00, 18:00 and 24:00 UTC (from left to right). The red box in (e) depicts the area for calculating precipitation over Germany for the ensemble sensitivity analysis presented in section 4.5.

smaller convective showers near the city of Nantes in the morning hours (starting around 06:00 UTC). The precursors of these
185 showers were initiated over the sea by a combination of large-scale forcing (determined by Q-vector divergence) and low-level
wind convergence (not shown). The subjectively determined track of the system in the REF run is marked by the red lines in all
model runs. This first convection initiation is displaced to the North compared to the satellite observations (Fig. 2), which was
nearer Bordeaux, and explains the northward displacement of the MCS track over Germany later in the evening. In addition to
the REF run (Fig. 6e), the runs NW, N, SW S, and, to a lesser extent also run E, successfully simulate convective precipitation
190 over northern Germany. In run E (Fig. 6f), the area with precipitation is too far ~~in the North-north~~ and the system decays too
early, west of Hamburg. The other successful model runs differ slightly from REF in the maximum rain amounts and horizontal
extent of precipitation on the ground. Nevertheless, the results of those runs is rather similar with respect to 24-h accumulated
precipitation. From these accumulations alone, the runs N, NW, or SW seem to be better suited as reference simulation due
to the larger precipitation amounts. However, the analysis of the temporal evolution (not shown) reveals that reference run is
195 closest to observations when both rain distribution and temporal development are considered.

The runs without any deep convection over northern Germany are the runs NE, W, and SE. Except for some weak and
isolated showers north of Cologne in the W run, there is no precipitation simulated in the region of interest. Given that the
model domain was shifted by only one grid point, this pronounced difference in the simulation results is ~~surprising~~unexpected.
All of these unsuccessful runs simulate more precipitation over the English Channel and the coastal regions of the Netherlands
200 than the REF run. Additionally, there is no systematic response of the model to domain shifting in any direction, e.g. there is no
systematic decrease of precipitation when shifting the domain from North to South or East to West and the three unsuccessful
simulations are not adjacent to one another. ~~However, as the precipitation in the E run is more to the North and does not extend
as much to the East as the other successful runs, all the eastward shift simulations have poorer prediction.~~

4.3 Differences in initial and boundary conditions

205 As we shifted the model domain only by one grid point towards the eight possible directions (referred to as Queen's case in
spatial statistics), we expect only small differences in the initial and boundary conditions. This is justified by the difference in
horizontal resolution of the initial data and the one used for the COSMO simulations. The spatial resolution of the IFS analyses
used in this study is approximately 13 km. As the COSMO simulations are run with 2.8 km grid spacing, many of the grid
points used in the preprocessor are the same if they are shifted by $\Delta x = 2.8$ km. This circumstance is illustrated in Fig. 7
210 in which the grid boxes of the input data and the COSMO grid of the southwest corner are displayed. Only for parts of the
model boundary does the domain shifting of the high-resolution grid also imply a different grid point used for interpolation
in the preprocessor of our model. Moreover, even when analyzing only the IFS input data, we do not see large point-to-point
gradients in any meteorological fields near the boundary of the nested COSMO simulations (not shown).

However, small differences are present and assessed quantitatively by domain-averaged meteorological variables at initial-
215 ization time (Table 2). Neither the surface fields (2 m temperature and specific humidity), nor the vertically-integrated variables
(convective available potential energy CAPE, convective inhibition CIN, 2.5–5-km averaged relative humidity RH, liquid water
path LWP, ice water path IWP, and deep layer shear DLS) does the model simulate any large differences in our ensemble of




Figure 7. Southwest corner of the simulation domain with illustration of IFS grid (black) and COSMO grid (blue). Numbers of 1 indicate IFS grid points whereas 0 indicates COSMO grid points.

Table 2. Domain averaged 2 m temperature (T in $^{\circ}\text{C}$), 2 m specific humidity (QV in g kg^{-1}), convective available potential energy (CAPE in J kg^{-1}), convective inhibition (CIN in J kg^{-1}), 2.5–5 km averaged relative humidity (RH in %), liquid water path (LWP in g m^{-2}), ice water path (IWP in g m^{-2}), and deep layer shear (DLS in m s^{-2}) at initialization time.

run	T	QV	CAPE	CIN	RH	LWP	IWP	DLS
NW	18.5	10.35	215.0	297.7	44.8	0.32	2.72	14.1
N	18.5	10.35	214.7	297.6	44.7	0.32	2.74	14.1
NE	18.5	10.36	214.8	297.5	44.7	0.32	2.76	14.1
W	18.5	10.36	215.0	298.1	44.7	0.32	2.72	14.1
REF	18.5	10.36	214.8	298.0	44.7	0.32	2.74	14.1
E	18.5	10.36	214.9	297.9	44.7	0.32	2.76	14.1
SW	18.5	10.37	215.1	298.6	44.7	0.32	2.72	14.1
S	18.5	10.37	214.9	298.4	44.7	0.32	2.73	14.0
SE	18.6	10.37	214.9	298.3	44.6	0.32	2.76	14.0

simulations. For example, the 2 m temperature differs by a maximum of 0.1°C between individual model runs. It is also of interest to investigate if the lateral boundaries (updated every 6 h) show any differences when the model domain is shifted. We therefore calculated averaged profiles for each of the four model boundaries for temperature, specific humidity and both horizontal wind components. The analysis of probability distributions (not shown) reveals that the range of simulated values is identical for all variables and only minor differences in the frequency of occurrence exist. Furthermore, averaged values of

those profiles are compared for every lateral boundary condition file (not shown). The maximum difference of the sensitivity runs to the REF runs is 0.02 K for temperature, 0.01 g kg^{-1} for specific humidity, and 0.1 m s^{-1} for the wind components. 225 We therefore conclude that all differences in the initial and boundary conditions of the domain-shifted model runs are small. ~~But given the chaotic nature of the atmosphere in convective weather events and the nonlinearity of the system with many feedbacks involved, these small deviations can determine whether a large MCS develops or not.~~

4.4 Convection-related parameters

The general preconditions for the initiation of deep moist convection are (i) conditional instability, (ii) a sufficient amount 230 of humidity in the lower and middle troposphere to form clouds, and (iii) a trigger process to bring air parcels to their level of free convection (e.g., Doswell III, 1987; Bennett et al., 2006). Trigger processes are e.g., the reaching of the convective temperature, lifting by convergence zones (e.g., Crook and Klemp, 2000), or terrain-induced ascent (Kirshbaum et al., 2018). The organization and further life cycle is then affected by the vertical wind shear, CAPE, and relative humidity. To assess the state of the atmosphere in the vicinity of the MCS affecting northern Germany, we calculated several convection-related 235 variables averaged over a rectangular box surrounding the convective system. The box has a size of $3^\circ \times 2.5^\circ$ and follows the storm along the path depicted in Fig. 6. The box has been positioned in such a way that the convection is not centered in the domain, but rather on the western edge to better capture the (preconvective) environment into which the storm is moving ~~-(see Fig. 6e for the location of this box at four selected times).~~

Figure 8 presents time series of some of these parameters during the life cycle of the convective storm. The brightly colored 240 lines represent the successful simulations, black is the reference simulations and gray and blue colored lines represent the unsuccessful simulations. The precipitation rate of the REF run is gradually increasing until ~~17:00~~14:30 UTC (Fig. 8a) ~~then there is and remains more or less constant until 17:00 UTC. The precipitation rate is strongly increasing until 18:30 UTC followed by~~ a slight reduction in intensity before ~~a second~~ the maximum is reached ~~at between~~ 21:00 UTC ~~and 22:00 UTC~~. After 22:00 UTC, the precipitation rate decreases and the convective system slowly dissipates. The other successful runs (NW, 245 N, SW, S) show larger precipitation rates and an earlier increase already from 12:00 UTC. As already mentioned earlier, these runs agree less well with observations in terms of precipitation location and timing than the REF run. The runs without an MCS over northern Germany (W, NE, SE) simulate similar precipitation amounts to the other runs until 11:00 UTC, but then rain gradually stops. Only run W simulates longer lasting precipitation until 14:00 UTC and a minor peak from a short-lived cell east of Eindhoven at 17:30 UTC.

250 The 0-6 km deep layer shear (Fig. 8b) is similar in all model runs with values of $27\text{--}30 \text{ m s}^{-1}$. Such high values indicate suitable conditions for highly-organized convection to develop in all runs because the precipitation and outflow become separated from the low-level updraft. Before the storms form there is almost no difference between the speed or direction of the wind shear in any of the simulations.

There is plenty of moisture available for convection, and both the mid-level relative humidity (Fig. 8c) and precipitable 255 water (Fig. 8d) show large values that increase as the storm environment moves further East later in the day. The simulations are again all very similar. The maxima in relative humidity are reached at 19:30 UTC which corresponds to the period with




Figure 8. Time series of convection-related parameters averaged over a $3^{\circ} \times 2.5^{\circ}$ box surrounding and following the convective system. Shear parameters are based on surface and 6-km altitude winds.

highest rain intensities. As the differences in relative humidity between the individual model runs are very small (2–4%), we determine that evaporation or entrainment processes are not responsible for the different model results. Moreover, between 14:00–22:00 UTC, the mid-level relative humidity is always higher than 60% which suggests that the role of entrainment of drier environmental air is probably only small. The same applies for the precipitable water for which all model realizations lie close together until 14:00 UTC (Fig. 8d). At later times, the precipitable water is affected by different rain formation and evaporation processes.

Additionally all simulations show substantial conditional instability, especially later in the day. Before 11:00 UTC, all models produce simulations have similar amounts of CAPE (Fig. 8e). Later on, in simulations with larger precipitation totals, the more CAPE has been consumed. This leaves the runs without an MCS over Germany with the highest CAPE values in the early evening (2500–3300 J kg^{-1}). For convection initiation or development, CAPE alone is not a suitable parameter. We therefore calculated the fraction of grid points, for which CAPE is larger than 600 J kg^{-1} and CIN is lower than 5 J kg^{-1}

(Fig. 8f). Here there is a large contrast in the number of grid-points where convection is expected at 13:00 UTC between the successful (around 10%) and unsuccessful (around 5%) runs. However, the reference run and the unsuccessful runs show rather similar curves. The W run reveals a somewhat lower maximum and a quick decrease afterwards. The secondary maximum occurring at 17:00 UTC corresponds to the aforementioned isolated cell initiated near Eindhoven. **The fraction of grid points fulfilling that criterion is primarily dominated by the existence of CAPE as domain-averaged CIN is very similar in all model runs until 12:00 UTC (not shown).**

Low-level wind convergence (Figs. 8g) is one mechanism for producing lift that leads to convection. The time series of convergence values are very similar to the upward vertical motion in the boundary layer (Figs. 8h) which indicates that the lift is primarily produced by convergence, mostly along convective outflow boundaries. Convergence early in the day can not be solely attributed to surface inhomogeneities or terrain features, because small amounts of rain are already simulated in the morning hours leading to wind convergence at outflow boundaries. Between 08:00–11:00 UTC, the convergence of the unsuccessful runs (NE, SE) is slightly weaker despite similar precipitation rates (Fig. 8a). However, after 11:00 UTC there is a clear split between the successful and unsuccessful simulations, with increased convergence and upward wind velocities in the successful simulations. Of the unsuccessful runs, only W exhibits similar convergence strength and lifting in the boundary-layer as the successful model runs and only until 13:00 UTC.

While this analysis does not completely separate the simulations into successful and unsuccessful subsets, there is information that helps explain the chance of producing an MCS. Clearly increased low-level convergence before 12:00 UTC is a good predictor of the later MCS, as is a large increase in the number of grid points with high CAPE but low CIN at 13:00 UTC. The lower CAPE and reduced deep-layer shear in the successful runs after 15:00 UTC are evidence of the storm modifying its own environment rather than a useful predictor of the MCS. ~~To refine the causes of the differences in these simulations we look at the evolution of the convective cells in more detail.~~

4.5 Ensemble sensitivity analysis

The above analysis has shown that whether the simulation is successful or not can be quantified based on small differences in the environment close to the developing convective cell. However, from this analysis we cannot tell what causes these changes in the preconvective environment. Here we use ensemble sensitivity analysis to help identify the origin of these differences. This analysis determines geographical areas, model variables and times that are correlated with a successful simulation. Although the correlations do not provide evidence of a causal relationship they do provide a starting point for understanding the diverging simulations, as shown in the below analysis and also by Barrett et al. (2015), Bednarczyk and Ancell (2015), Hill et al. (2016), Torn et al. (2017).

Following the above papers, we define the ensemble sensitivity S as

$$S = m a \sigma_x = \frac{\text{cov}(y, x)}{\text{var}(x)} a \sigma_x \quad (1)$$

where m is the regression coefficient between the test variable x and the response function y as calculated at each grid point. A scaling factor a is used to de-emphasize noise in the analysis based on the correlation coefficient r ; $a = 0$ where $r^2 < 0.4$

and $a = 1$ otherwise. Finally the sensitivity is scaled by the ensemble standard deviation of the test variable σ_x , calculated individually at each grid point, to normalize the calculated sensitivity ~~which~~; this enables comparison of sensitivities to different model variables.

We attempted to use ensemble sensitivity analysis on numerous model variables at surface, 850, 500 and 250 hPa levels. However, because the model initial states are nearly identical, the ensemble sensitivity analysis is unable to identify any relationships between convection intensity and typical large-scale drivers of convection before the convection develops. Only after the convection develops can signals be seen in the e.g. upper-level pressure, wind and divergence fields as they are directly modified by the convection. Therefore the analysis below focusses on ~~near-surface model fields~~: CAPE, precipitation rate and vertical velocity. These variables are those shown in Fig. 8abd also discussed in more detail in section 4.6 and Fig. 10.

In Fig. 9, we show the ensemble sensitivity of surface precipitation over Germany to temperature (T) and zonal wind (u) at 200 above the ground, at 09:00, 11:00 and 13:00 UTC. The sensitivity is calculated using the mean precipitation over northern Germany between 12:00–23:00 and 13:00 UTC. The precipitation values used are the mean in the (box bounded by 7 W, 14 W, 52 N, 53.5 N; red box in Fig. 6e), which is 4.4–6.7 mm in the successful ensemble members 1.0 mm in the E-shifted member and less than 0.1 mm otherwise. The resulting sensitivity S has units of mm of precipitation per standard deviation change in x (here T or CAPE, precipitation rate or vertical velocity W). Hence the ensemble sensitivity is interpreted as the change to precipitation for a one standard deviation increase in variable x at that grid point.

Working backwards in time from the right hand column to the left, we can see that the ensemble reproduces the MCS better in members where the temperature across northern France (49–50 N, 1–2 E) is lowest and where the wind speed is highest just before MCS development. The ensemble sensitivity analysis (Fig. 9) shows a correlation between increased precipitation over northern Germany with lower CAPE over northern France earlier in the day (11:00–13:00 UTC; left column of Fig. 9). Precipitation located at the northern end of this region of low CAPE is also correlated with successful simulation of the MCS (middle column of Fig. 9). There is also a weak signal of sensitivity to weaker vertical velocities over the region of sensitivity to low CAPE (right column of Fig. 9), with stronger vertical velocities located to the east of the low temperatures. This is a clear sign of CAPE region in location of updrafts in the different simulations. These signals should be interpreted as a relationship between increased precipitation across northern Germany in ensemble members with a strong surface cold pool in the over northern France earlier in the day (at locations marked by the sensitivity. In the middle column, we see that this signal is already evident negative sensitivity to CAPE). The signal is first evident in sensitivity to CAPE at 11:00 UTC near 48.8 N, 0.4 E and in the left column. However, the signal is much weaker at 09:00 UTC (near 47.8 N, 0.75 W) but only in the wind field at this time as the cold pool is yet to develop. A convective system exists near this location 47.8 N, 0.75 W in all ensemble members at 09:00 UTC but the location at 09:00 appears (not shown) but the decisive factor regarding later MCS development seems to be the decisive factor exact location at this time, with a location farther to the east apparently favored.

This favoring MCS development. This does not show up well in the ensemble sensitivity analysis because the disparate locations of the cells mean that values at individual grid-points are not correlated with later success.

335 The disturbance at 09:00 UTC can be tracked farther back to the western french coast, around 150 km south of Nantes (46 N,
1 W) already at 04:00 UTC (not shown). However, neither the ensemble sensitivity analysis nor more detailed investigation
into the convective disturbances at this time showed any systematic structure of the convective cells that were decisive in
the successful simulation of the MCS later in the day. The important aspect appears to be that by 09:00 UTC that a line of
convection begins to form on the outflow of this convection, ~~as seen in the lower-left panel of figure 9,~~ and that changing the
340 position of that cell by only around 10 km determines whether the convective cell evolves into the MCS which later affects
Germany, or not.

The ensemble sensitivity analysis has helped highlight interesting areas in the development timeline of the convective cells.
However, due to the disparate locations of the convective cells, the grid-point correlations required for ensemble sensitivity
analysis do not help explain how these cells differ in their development. In the next section we evaluate in more detail how the
345 developing convective cells interacted with their environment and what caused the differing convective evolutions.

4.6 Simulation result differences

In this section we discuss horizontal cross-sections of convection-related parameters to elucidate the differences (and their
possible causes) of the different model results. For the sake of brevity, we only compare the reference run to two unsuccessful
runs, namely the ones with the model domain shifted towards the W and the SE. Figure 10 presents a time series of those
350 cross-sections for the region of northwestern France and southern England.

The analysis at 10:00 UTC shows a very similar picture for all simulations at all times, with CAPE increasing to the South
and East, wind is westerly over the English Channel, turning to northerly direction over France and there is low CIN over
France from 11:00 UTC at the latest. The simulations all have weak, disorganized convection over northwestern France and
a more isolated cell at the border between France and Belgium. The region with the convective system of interest is marked
355 with a red circle. However, as time develops, only the REF simulation produces a convective system that moves into the high
CAPE region to the East and later becomes an MCS over Germany. By 10:30 UTC, CIN is less than 5 J kg^{-1} in the REF and
SE run, whereas it is still above that threshold until 11:00 UTC in the W run. The highest rain intensities are also simulated in
the REF run. At 11:00 UTC approximately half of the cell of interest in the W run is over the sea, where CAPE is lower and
CIN higher than over land. In contrast, the area of convective rain in the REF run is separated from the rain over the sea and
360 precipitation intensity remains high. The precipitating area in the SE run is also separated from the larger rain area over the
sea, but the precipitation rate is already weaker than in the REF run. At 11:30 UTC, the cell in the W run is weakening and
lies almost at the coastline, whereas the cell in the REF run still remains almost entirely over land while moving towards the
North-East. The corresponding system in the SE run is also weakening while traveling towards the North-East; approximately
half of the cell is now located over the sea. The systems in the W and REF runs both weaken at 12:00 UTC, but the one from
365 the SE run has already decayed. Only the cell in the REF run intensifies again at 12:30 UTC. In the W run, the cell continues
to move along the coastline while weakening, until it is completely dissolved at 14:00 UTC. In the REF run, however, the cell
stays almost completely over land and intensifies further while moving towards the Netherlands (13:00–14:00 UTC).

The sea surface temperatures along the French coast lie around 15°C and are much lower than the land surface temperatures (around 23°C, not shown). This temperature distribution is similar in all model runs for the preconvective environment. The proximity of the cell to the colder sea surface appears to have a decisive influence on the further life cycle of convection. In the REF run and the other successful runs (not shown), the system stays more or less entirely over land between 11:00–12:00 UTC. In these successful simulations, the systems travels further towards Belgium and Germany (rather than over the sea), where it encounters more favorable convective conditions including higher CAPE, which later allows it to evolve into an MCS.

The isolated cell, to the north west of these plots between 10:00–11:00 UTC, does not appear to be important to the decay of the cell of interest. It is located approximately 150 km upstream. The cell is stronger in the W run leading to a slight reduction of CAPE and therefore creating slightly less favorable environmental conditions in the area into which the main cell would later move. However, it appears that the weakening of the main cell occurred independently of the cell upstream and can rather be attributed to the proximity to the colder sea surface.

In addition to this analysis, we further want to point out that the upper-level dynamics are similar in all model runs in the early stage of the convection over France. Hoskins et al. (1978) showed that the traditional form of the quasi-geostrophic omega equation can be rewritten using the Q-vector and that regions of upward (downward) vertical motion are associated with Q-vector convergence (divergence). We calculated the divergence of the Q-vector at 500 hPa and found no noticeable differences between successful and unsuccessful runs, nor does the model simulate any variations are there meaningful differences in geopotential height (not shown). This indicates that the large scale forcing is similar for these model runs and not responsible for the simulation result differences.

4.7 Further MCS evolution

Having established a possible explanation for the decay of the precursors of the MCS in the previous section, we now analyze the further evolution of the system into a MCS using the reference simulation (Fig. 11). To the east of the system, the model simulates an east-west oriented region of high low-level equivalent potential temperature in the north-central part of Germany, which corresponds to CAPE values between 3000–4000 J kg⁻¹. This CAPE-rich air is advected with easterly winds towards the convective system over the Netherlands. Colliding with the cell's outflow, a strong low-level mass and moisture convergence occurs, which fosters the evolution into a MCS. As already discussed in section 4.4, the 0–6 km deep layer shear shows suitable conditions for highly-organized convection (27–30 m s⁻¹). The maximum rain intensities reach locally up to 22 mm (30 min)⁻¹ with a weakly defined bow-like structure of precipitation, typical of storms with an intense rear-inflow jet. In the wake of the MCS, CAPE is almost entirely consumed. From 23:00 UTC onwards, the MCS is decaying while further traveling towards Poland (not shown).

5 Comparison of methods to perturb initial and boundary conditions

Many operational forecast centers produce both a high-resolution forecast and an ensemble of lower resolution to provide a measure of uncertainty (Rodwell et al., 2013). There are various ways of generating an ensemble, such as perturbations to

400 the initial conditions and/or boundary conditions (e.g., Montani et al., 2011; Kühnlein et al., 2014), stochastic physical parameterizations (e.g., Buizza et al., 1999; Berner et al., 2017), or ensemble data assimilation such as ensemble Kalman filter (e.g., Dowell et al., 2004; Zhang et al., 2004; Reich et al., 2011). Recent studies by Schneider et al. (2019) and Keil et al. (2019) have also shown that different assumptions for the amount of cloud condensation nuclei could be included in convective-scale ensemble forecasting, but only if the model employs a double-moment microphysics scheme. Because of the fundamental uncertainties of the simulations due to nonlinearities of the model equations, several studies have noted the significant impact of initial boundary conditions (IBC) and lateral boundary conditions (LBC) on the simulation of convective precipitation for some situations (e.g., Hohenegger et al., 2006; Trentmann et al., 2009; Richard et al., 2011; Bouttier and Raynaud, 2018) and that ensemble members with the most accurate initial and boundary conditions are most skilful at predicting the location of convective initiation (Barrett et al., 2015).

410 One common approach for accounting for uncertainties in the initial and boundary conditions is that perturbations entering the model from the lateral boundaries can be provided by different driving EPS members as it is the case for COSMO-LEPS (Montani et al., 2011) or COSMO-DE-EPS (Gebhardt et al., 2011; Kühnlein et al., 2014). Such perturbations have been shown to play a more and more important role in the behaviour of the limited-area system as the forecast range increases. However, this methodology needs an algorithm to select representative members from the driving ensemble (Marsigli et al., 2001).
415 Bouttier and Raynaud (2018) showed that the algorithms used for the member selection have a significant impact on the resulting ensembles and that clustering-based methods outperformed random subsampling. Torn et al. (2006) proposed two classes of methods to populate a boundary condition ensemble. The ensemble of boundary conditions is either provided by an ensemble Kalman filter (EnKF) on a larger domain corresponding to a random draw from the probability distribution function for the forecast (or analysis) on the limited-area domain boundary or by a fixed-covariance perturbation technique.
420 Romine et al. (2014) stated that there remains value in randomly perturbing a deterministic forecast for ensemble lateral boundary conditions. They examined convection-permitting ensemble forecasts by drawing initial conditions from a downscaled ensemble data assimilation system. As the control ensemble was underdispersive, it was supplemented also by perturbed lateral boundary conditions. Lateral boundary conditions for the cycled analysis were generated from perturbed forecasts with the fixed covariance method (Torn et al., 2006). This technique led to a modest improvement in spread and the least degradation in
425 systematic bias.

To further evaluate the large forecast variability of our simulations, we conducted another experiment on the domain of the successful reference run. In this new model run, however, we added small, stochastic perturbations on the boundary conditions, namely random temperature fluctuations with a Gaussian distribution of zero mean and a standard deviation of 0.01 K at all levels. The run does not reproduce the MCS over Germany (not shown) and the rain distribution is similar to the other
430 unsuccessful (e.g. runs NE or SE in Fig. 6). This finding indicates that the simple method of domain shifting can produce a result comparable to stochastic LBC perturbations and could therefore be used to estimate the uncertainty in convection-resolving simulations. In agreement with Henneberg et al. (2018) who used domain shifting by 10 and 30 grid points on a smaller domain combined with soil moisture uncertainties, domain shifting does provide a sufficient large model spread for the case analyzed in this study.

435 Generally, the approach of domain shifting to generate different IBCs is attractive due to its relative simplicity and practicality of implementation. An advantageous aspect is that IBCs originate from the same driving model, which avoids errors due to inconsistencies that may occur if the initial condition perturbations are applied independently from the boundary condition perturbations (Caron, 2013). Another advantage is the fact that this methodology does not need an algorithm to select representative members if a driving ensemble is used. However, we want to point out that this study aims at revealing the large
440 forecast variability which can be achieved from domain shifting for this particular case. The applicability of this method needs to be compared with other techniques for this and other cases as well which is left for future work.

6 Summary and conclusions

During Pentecost 2014, following a period of hot weather, a mesoscale convective system formed over France and traveled towards Germany in the afternoon of 9 June. A strong southwesterly flow lead to a favorable environment for deep convection
445 due to the advection of warm and moist air. However, the predictability of this event was very low; neither the operational deterministic nor any member of the ensemble prediction system (both convection resolving) captured the event with more than 12 hours lead time (Barthlott et al., 2017).

Hindcasts of this situation were performed with convection-permitting resolution on a large model domain, enabling the simulation of the whole life cycle of the system originating from the western Atlantic coast. The results show that the MCS
450 was reasonably well represented by the COSMO model in this setup. When compared to radar-derived precipitation rates, the MCS was simulated somewhat shifted to the North and the translation speed was slightly higher than observed.

The low predictability of the event was again evident; moving the model domain by just one grid point changed whether the MCS over Germany is successfully simulated or not. The domain was shifted systematically in eight directions (N, NE, E, SE, S, SW, W, NW) by just one grid point and three of these configurations completely failed to simulate deep convection over
455 Germany on that day, while a fourth had some convection but did not capture the organized MCS. This large impact is **even more surprising unexpected** when considering the comparatively large computational domain of 1668 km × 1807 km.

The evaluation of domain-averaged initial conditions, like low-level temperature, moisture, relative humidity, or wind shear showed only negligible differences. The temporal evolution of convection-related parameters in the vicinity of the storm system also revealed similar conditions in its preconvective environment. The ensemble sensitivity analysis was unable to reveal
460 differences in the upper-level flow between ensemble members, although low-level differences associated with a developing cold pool were identified. An explanation of the large differences in the model results lies in the proximity of the track of the convective system to the north coast of France and the colder temperatures over the sea than the land. The convective system in the successful runs stays more or less entirely over land, allowing it to eventually reach a region favorable for convective organization (with high CAPE, large shear and low CIN), whereas the early convection in the unsuccessful runs moved closer
465 to the coast and had considerable portions located over the sea. This small displacement seems to be the main point deciding if the system decays or is able to live on and intensify into an MCS.

Although perhaps an extreme example, this case is in agreement with many previous studies pointing out the effects of small-scale variability in atmospheric parameters (e.g. Crook, 1996; Weckwerth, 2000). These results emphasize the difficulty of forecasting the location and intensity of convective precipitation due to the chaotic nature of the atmosphere in convective
470 weather events and the nonlinearity of the system with many feedbacks. In this case it is required to capture a chain of events that is dependent on precisely predicting the location of initial convection; only if the outflow of the initial convective system occurs in the right location can the damaging MCS be triggered.

The results of this work suggests that model domain shifting could be used to quantify how ~~uncertainties in the initial and boundary conditions contribute~~ internal model variability contributes to the predictability of an event. However, this single
475 case study needs to be expanded to cover more cases, for example in weather regimes with strong synoptic forcing and more stratiform precipitation and in other models such as ICON (ICOsahedral Non-hydrostatic) (Zängl et al., 2015). Moreover, whether changing the extent of domain shifting (e.g. from 1–10 grid points) is important should be evaluated.

Data availability. COSMO model output is available on request from the authors.

Author contributions. CB and AB both designed the numerical experiments, CB carried them out and performed the data analyses. Both
480 authors contributed equally to the writing of the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The research leading to these results has been done within the subprojects B3 and B1 of the Transregional Collaborative Research Center SFB/TRR 165 “Waves to Weather” (www.wavestoweather.de) funded by the German Research Foundation (DFG). This work was performed on the computational resource ForHLR I funded by the Ministry of Science, Research and the Arts Baden-Württemberg
485 and DFG (“Deutsche Forschungsgemeinschaft”). The authors wish to thank the Deutscher Wetterdienst (DWD) for providing the COSMO model code, the initial and boundary data as well as Radar-derived precipitation data. We are also grateful to U. Schättler (DWD) for his help in setting up the preprocessor int2lm of the COSMO model. Satellite pictures were kindly provided by Folke Olesen (IMK-ASF, KIT).

Financial support. This work was partly funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 257899354 – TRR 165.

490 References

- Barrett, A. I., Gray, S. L., Kirshbaum, D. J., Roberts, N. M., Schultz, D. M., and Fairman Jr, J. G.: Synoptic versus orographic control on stationary convective banding, *Q. J. R. Meteorol. Soc.*, 141, 1101–1113, <https://doi.org/10.1002/qj.2409>, 2015.
- Barrett, A. I., Gray, S. L., Kirshbaum, D. J., Roberts, N. M., Schultz, D. M., and Fairman Jr, J. G.: The utility of convection-permitting ensembles for the prediction of stationary convective bands, *Mon. Wea. Rev.*, 144, 1093–1114, <https://doi.org/10.1175/MWR-D-15-0148.1>,
495 2016.
- Barrett, A. I., Wellmann, C., Seifert, A., Hoose, C., Vogel, B., and Kunz, M.: One step at a time: How model timestep significantly affects Convection-Permitting simulations, *Journal of Advances in Modeling Earth Systems*, 11, 641–658, <https://doi.org/10.1029/2018MS001418>, 2019.
- Barthlott, C. and Hoose, C.: Spatial and temporal variability of clouds and precipitation over Germany: multiscale simulations across the
500 “gray zone”, *Atmos. Chem. Phys.*, 15, 12 361–12 384, <https://doi.org/10.5194/acp-15-12361-2015>, 2015.
- Barthlott, C. and Hoose, C.: Aerosol effects on clouds and precipitation over central Europe in different weather regimes, *J. Atmos. Sci.*, 75, 4247–4264, <https://doi.org/10.1175/JAS-D-18-0110.1>, 2018.
- Barthlott, C., Hauck, C., Schädler, G., Kalthoff, N., and Kottmeier, C.: Soil moisture impacts on convective indices and precipitation over complex terrain, *Meteorol. Z.*, 20, 185–197, <https://doi.org/10.1127/0941-2948/2011/0216>, 2011.
- 505 Barthlott, C., Mühr, B., and Hoose, C.: Sensitivity of the 2014 Pentecost storms over Germany to different model grids and microphysics schemes, *Q. J. R. Meteorol. Soc.*, 143, 1485–1503, <https://doi.org/10.1002/qj.3019>, 2017.
- Bednarczyk, C. N. and Ancell, B. C.: Ensemble Sensitivity Analysis Applied to a Southern Plains Convective Event, *Mon. Wea. Rev.*, 143, 230–249, <https://doi.org/10.1175/MWR-D-13-00321.1>, 2015.
- Bennett, L. J., Browning, K. A., Blyth, A. M., Parker, D. J., and Clark, P. A.: A review of the initiation of precipitating convection in the
510 United Kingdom, *Q. J. R. Meteorol. Soc.*, 132, 1001–1020, <https://doi.org/10.1256/qj.05.54>, 2006.
- Berner, J., Achatz, U., Batté, L., Bengtsson, L., de la Cámara, A., Christensen, H. M., Colangeli, M., Coleman, D. R. B., Crommelin, D., Dolaptchiev, S. I., Franzke, C. L. E., Friederichs, P., Imkeller, P., Järvinen, H., Juricke, S., Kitsios, V., Lott, F., Lucarini, V., Mahajan, S., Palmer, T. N., Penland, C., Sakradzija, M., von Storch, J.-S., Weisheimer, A., Weniger, M., Williams, P. D., and Yano, J.-I.: Stochastic Parameterization: Toward a New View of Weather and Climate Models, *Bull. Amer. Meteor. Soc.*, 98, 565–588,
515 <https://doi.org/10.1175/BAMS-D-15-00268.1>, 2017.
- Bouttier, F. and Raynaud, L.: Clustering and selection of boundary conditions for limited area ensemble prediction, *Q. J. R. Meteorol. Soc.*, 144, 2381–2391, <https://doi.org/10.1002/qj.3304>, 2018.
- Buizza, R., Miller, M., and Palmer, T.: Stochastic representation of model uncertainties in the ECMWF ensemble prediction system, *Q. J. R. Meteorol. Soc.*, 125, 2887–2908, 1999.
- 520 Caron, J.-F.: Mismatching Perturbations at the Lateral Boundaries in Limited-Area Ensemble Forecasting: A Case Study, *Mon. Wea. Rev.*, 141, 356–374, <https://doi.org/10.1175/MWR-D-12-00051.1>, 2013.
- Clark, P., Roberts, N., Lean, H., Ballard, S. P., and Charlton-Perez, C.: Convection-permitting models: a step-change in rainfall forecasting, *Meteorol. Appl.*, 23, 165–181, <https://doi.org/10.1002/met.1538>, 2016.
- Crook, N. A.: Sensitivity of moist convection forced by boundary layer processes to low-level thermodynamic fields, *Mon. Wea. Rev.*, 124,
525 1767–1785., 1996.
- Crook, N. A. and Klemp, J. B.: Lifting by convergence lines, *J. Atmos. Sci.*, 57, 873–890., 2000.

- Doswell III, C. A.: The Distinction between Large-Scale and Mesoscale Contribution to Severe Convection: A Case Study Example, *Wea. Forecasting*, 2, 3–16, [https://doi.org/10.1175/1520-0434\(1987\)002<0003:TDBLSA>2.0.CO;2](https://doi.org/10.1175/1520-0434(1987)002<0003:TDBLSA>2.0.CO;2), 1987.
- 530 Dowell, D. C., Zhang, F., Wicker, L. J., Snyder, C., and Crook, N. A.: Wind and temperature retrievals in the 17 May 1981 Arcadia, Oklahoma supercell: Ensemble Kalman filter experiments, *Mon. Wea. Rev.*, 132, 1982–2005, 2004.
- Gebhardt, C., Theis, S., Paulat, M., and Ben Bouallègue, Z.: Uncertainties in COSMO-DE precipitation forecasts introduced by model perturbations and variation of lateral boundaries, *Atmos. Res.*, 100, 168–177, <https://doi.org/10.1016/j.atmosres.2010.12.008>, 2011.
- Groenemeijer, P.: Summary of the evaluation of DWD NWP products and visualizations at the ESSL Testbed 2014, Available online at <https://www.essl.org/cms/publications> [Accessed on 24 July 2019], 28 pp., 2014.
- 535 Henneberg, O., Ament, F., and Grützun, V.: Assessing the uncertainty of soil moisture impacts on convective precipitation using a new ensemble approach, *Atmos. Chem. Phys.*, 18, 6413–6425, <https://doi.org/10.5194/acp-18-6413-2018>, 2018.
- Hill, A. J., Weiss, C. C., and Ancell, B. C.: Ensemble Sensitivity Analysis for Mesoscale Forecasts of Dryline Convection Initiation, *Mon. Wea. Rev.*, 144, 4161–4182, <https://doi.org/10.1175/MWR-D-15-0338.1>, 2016.
- Hohenegger, C. and Schär, C.: Predictability and Error Growth Dynamics in Cloud-Resolving Models, *J. Atmos. Sci.*, 64, 4467–4478, 540 <https://doi.org/10.1175/2007JAS2143.1>, 2007.
- Hohenegger, C., Lüthi, D., and Schär, C.: Predictability Mysteries in Cloud-Resolving Models, *Mon. Wea. Rev.*, 134, 2095–2107, <https://doi.org/10.1175/MWR3176.1>, 2006.
- Hoskins, B. J., Draghici, I., and Davies, H. C.: A new look at the omega-equation, *Q. J. R. Meteorol. Soc.*, 104, 31–38, 1978.
- Keil, C., Baur, F., Bachmann, K., Rasp, S., Schneider, L., and Barthlott, C.: Relative contribution of soil moisture, boundary-layer 545 and microphysical perturbations on convective predictability in different weather regimes, *Q. J. R. Meteorol. Soc.*, 145, 3102–3115, <https://doi.org/10.1002/qj.3607>, 2019.
- Kirshbaum, D. J., Adler, B., Kalthoff, N., Barthlott, C., and Serafin, S.: Moist Orographic Convection: Physical Mechanisms and Links to Surface-Exchange Processes, *Atmosphere*, 9, 80, <https://doi.org/10.3390/atmos9030080>, 2018.
- Kühnlein, C., Keil, C., Craig, G. C., and Gebhardt, C.: The impact of downscaled initial condition perturbations on convective-scale ensemble 550 forecasts of precipitation, *Q. J. R. Meteorol. Soc.*, 140, 1552–1562, <https://doi.org/10.1002/qj.2238>, 2014.
- Landman, W. A., Seth, A., and Camargo, S. J.: The Effect of Regional Climate Model Domain Choice on the Simulation of Tropical Cyclone-Like Vortices in the Southwestern Indian Ocean, *J. Climate*, 18, 1263–1274, <https://doi.org/10.1175/JCLI3324.1>, 2005.
- Marsigli, C., Montani, A., Nerozzi, F., Paccagnella, T., Tibaldi, S., Molteni, F., and Buizza, R.: A strategy for high-resolution ensemble prediction. Part II: Limited-area experiments in four Alpine flood events, *Q. J. R. Meteorol. Soc.*, 127, 2095–2115, 555 <https://doi.org/10.1002/qj.49712757613>, 2001.
- Mathias, L., Ermert, V., Kelemen, F. D., Ludwig, P., and Pinto, J. G.: Synoptic analysis and hindcast of an intense bow echo above Western Europe: The Pentecost storm 2014, *Wea. Forecasting*, 32, 1121–1141, <https://doi.org/10.1175/WAF-D-16-0192.1>, 2017.
- Mellor, G. L. and Yamada, T.: A hierarchy of turbulence closure models for planetary boundary layers, *J. Atmos. Sci.*, 31, 1791–1806, 1974.
- Miguez-Macho, G., Stenchikov, G. L., and Robock, A.: Spectral nudging to eliminate the effects of domain position and geometry in regional 560 climate model simulations, *J. Geophys. Res.*, 109, <https://doi.org/10.1029/2003JD004495>, d13104, 2004.
- Montani, A., Cesari, D., Marsigli, C., and Paccagnella, T.: Seven years of activity in the field of mesoscale ensemble forecasting by the COSMO-LEPS system: Main achievements and open challenges, *Tellus Ser. A*, 63, 605–624, 2011.
- Reich, H., Rhodin, A., and Schraff, C.: LETKF for the nonhydrostatic regional model COSMO-DE, *COSMO Newsletter*, 11, 27–31, 2011.

- Richard, E., Chaboureaud, J. P., Flamant, C., Champollion, C., Hagen, M., Schmidt, K., Kiemle, C., Corsmeier, U., Barthlott, C., and Di
565 Girolamo, P.: Forecasting summer convection over the Black Forest: a case study from the Convective and Orographically-induced Pre-
cipitation Study (COPS) experiment, *Q. J. R. Meteorol. Soc.*, 137 (S1), 101–117, <https://doi.org/10.1002/qj.710>, 2011.
- Rodwell, M. J., Magnusson, L., Bauer, P., Bechtold, P., Bonavita, M., Cardinali, C., Diamantakis, M., Earnshaw, P., Garcia-Mendez, A.,
Isaksen, L., et al.: Characteristics of occasional poor medium-range weather forecasts for Europe, *Bull. Amer. Meteor. Soc.*, 94, 1393–
1405, <https://doi.org/10.1175/BAMS-D-12-00099.1>, 2013.
- 570 Romine, G. S., Schwartz, C. S., Berner, J., Fossell, K. R., Snyder, C., Anderson, J. L., and Weisman, M. L.: Representing Forecast Error in a
Convection-Permitting Ensemble System, *Mon. Wea. Rev.*, 142, 4519–4541, <https://doi.org/10.1175/MWR-D-14-00100.1>, 2014.
- Rossa, A., Bruen, M., Frühwald, D., Macpherson, B., Holleman, I., Michelson, D., and Michaelides, S.: Use of Radar Observations in
Hydrological and NWP Models, Available at <http://www.cost.eu/media/publications> [Accessed on 24 July 2019], 2005.
- Schättler, U., Doms, G., and Schraff, C.: A description of the nonhydrostatic regional COSMO-model, Part VII: User’s Guide, Available
575 online at <http://www.cosmo-model.org> [Accessed on 24 July 2019], 227 pp., 2016.
- Schlüter, I. and Schädler, G.: Sensitivity of Heavy Precipitation Forecasts to Small Modifications of Large-Scale Weather Patterns for the
Elbe River, *J. Hydrometeorol.*, 11, 770–780, <https://doi.org/10.1175/2010JHM1186.1>, 2010.
- Schneider, L., Barthlott, C., Hoose, C., and Barrett, A. I.: Relative impact of aerosol, soil moisture, and orography perturbations on deep
convection, *Atmos. Chem. Phys.*, 19, 12 343–12 359, <https://doi.org/10.5194/acp-19-12343-2019>, 2019.
- 580 Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part I: Model description,
Meteorol. Atmos. Phys., 92, 67–82, 2006.
- Seth, A. and Giorgi, F.: The Effects of Domain Choice on Summer Precipitation Simulation and Sensitivity in a Regional Climate Model, *J.*
Climate, 11, 2698–2712, [https://doi.org/10.1175/1520-0442\(1998\)011<2698:TEODCO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2698:TEODCO>2.0.CO;2), 1998.
- Tiedtke, M.: A comprehensive mass flux scheme for cumulus parameterization in large-scale models, *Mon. Wea. Rev.*, 117, 1779–1800,
585 1989.
- Torn, R. D., Hakim, G. J., and Snyder, C.: Boundary Conditions for Limited-Area Ensemble Kalman Filters, *Mon. Wea. Rev.*, 134, 2490–
2502, <https://doi.org/10.1175/MWR3187.1>, 2006.
- Torn, R. D., Romine, G. S., and Galarneau, T. J.: Sensitivity of Dryline Convection Forecasts to Upstream Forecast Errors for Two Weakly
Forced MPEX Cases, *Mon. Wea. Rev.*, 145, 1831–1852, <https://doi.org/10.1175/MWR-D-16-0457.1>, 2017.
- 590 Trentmann, J., Keil, C., Salzmann, M., Barthlott, C., Bauer, H.-S., Lawrence, M., Leuenberger, D., Wernli, H., Wulfmeyer, V., Corsmeier, U.,
and Kottmeier, C.: Multi-model simulations of a convective situation in low-mountain terrain in central Europe, *Meteorol. Atmos. Phys.*,
103, 95–103, <https://doi.org/10.1007/s00703-008-0323-6>, 2009.
- Vié, B., Nuissier, O., and Ducrocq, V.: Cloud-resolving ensemble simulations of Mediterranean heavy precipitation events: Uncertainty on
initial conditions and lateral boundary conditions, *Mon. Wea. Rev.*, 139, 403–423, <https://doi.org/10.1175/2010MWR3487.1>, 2011.
- 595 Weckwerth, T.: The effect of small-scale moisture variability on thunderstorm initiation, *Mon. Wea. Rev.*, 128, 4017–4030., 2000.
- Wicker, L. J. and Skamarock, W. C.: Time-splitting methods for elastic models using forward time schemes, *Mon. Wea. Rev.*, 130, 2088–
2097, 2002.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M:
Description of the non-hydrostatic dynamical core, *Q. J. R. Meteorol. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.
- 600 Zhang, F., Snyder, C., and Sun, J.: Impacts of Initial Estimate and Observation Availability on Convective-Scale Data Assimilation with an
Ensemble Kalman Filter, *Mon. Wea. Rev.*, 132, 1238–1253, 2004.




Figure 9. Ensemble sensitivity of precipitation-precipitation over Germany to temperature-CAPE (top-row-left column) and zonal-wind component-precipitation rate (bottom-row-middle column) both-and vertical velocity at 200500 m above the surface height (right column). The units are mm per standard deviation change in the ensemble, with positive values indicating that ensemble members produced more precipitation over Germany when the temperature-CAPE, precipitation rate or wind-speed-vertical velocity in the marked locations was larger in the ensemble. A signal of increased precipitation for lower temperatures-and-increased-wind-speed-develop-CAPE values develops throughout the morning, consistent with a large developing cold pool (left column). The precipitation in the successful simulations is at the northern end of this cold pool (middle column); however, the location of the low-level updrafts is too variable in the ensemble to be seen with this analysis (right column).




Figure 10. Convective available potential energy (color shading, in J kg^{-1} , 30-min precipitation rate (blue color shading, in mm (30 min)^{-1} , and 10-m wind field (arrows) between 14:00–18:00 UTC on 9 June. Gray areas indicate low-level wind convergence larger than $0.35 \cdot 10^{-3} \text{ m s}^{-1}$ and hatched areas represent regions where convective inhibition is smaller than 5 J kg^{-1} . Left: domain shifted to W; Middle: reference run; Right: domain shifted to SE. The red circle indicates the region of the convective cell developing into a MCS.




Figure 10. Continued.




Figure 11. As Panels as in Fig. 10, but showing the development of the MCS over Benelux and Germany between 16:00–21:00 UTC in the REF simulation.