

A process-based anatomy of Mediterranean cyclones: From baroclinic lows to tropical-like systems

Emmanouil Flaounas¹, Suzanne L. Gray², Franziska Teubler³

¹Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

5 ²Department of Meteorology, University of Reading, Reading, UK

³Johannes Gutenberg-Universität Mainz, Mainz, Germany

Correspondence to: Emmanouil Flaounas (Emmanouil.flaounas@env.ethz.ch)

Abstract. In this study, we address the question of the atmospheric processes that turn Mediterranean cyclones into severe storms. Our approach applies on-line potential vorticity (PV) budget diagnostics and piecewise PV inversion to WRF model simulations of the mature stage of 100 intense Mediterranean cyclones. We quantify the relative contributions of different processes to cyclone development and therefore deliver, for the first time, a comprehensive insight into the variety of cyclonic systems that develop in the Mediterranean from the perspective of cyclone dynamics.

In particular, we show that all 100 cyclones are systematically influenced by two main PV anomalies: a major anomaly in the upper troposphere, related to the baroclinic forcing of cyclone development, and a minor anomaly in the lower troposphere, related to diabatic processes and momentum forcing of wind. Among the diabatic processes, latent heat is shown to act as the main PV source (reinforcing cyclones), being partly balanced by PV sinks of temperature diffusion and radiative cooling (weakening cyclones). Momentum forcing is shown to have an ambiguous feedback, able to reinforce and weaken cyclones while in certain cases playing an important role in cyclone development. Piecewise PV inversion shows that most cyclones develop due to the combined effect of both baroclinic and diabatic forcing, i.e. due to both PV anomalies. However, the stronger the baroclinic forcing, the less a cyclone is found to develop due to diabatic processes. Several pairs of exemplary cases are used to illustrate the variety of contributions of atmospheric processes to the development of Mediterranean cyclones: (i) cases where both baroclinic and diabatic processes contribute to cyclone development; (ii) cases that mainly developed due to latent-heat release; (iii) cases developing in the wake of the Alps; and (iv) two unusual cases, one where momentum forcing dominates cyclone development and the other presenting a dual surface pressure centre. Finally, we focus on ten medicane cases (i.e. tropical-like cyclones). In contrast to their tropical counterparts—but in accordance with most intense Mediterranean cyclones—most medicanes are shown to develop under the influence of both baroclinic and diabatic processes. In discussion of medicane driving processes, we highlight the need for a physical definition of these systems.

30 1 Introduction

The Mediterranean basin is among the most cyclogenetic regions in the world (Petterssen 1956; Alpert [et al., et al.](#) 1990; Maheras [et al., et al.](#) 2001; Neu [et al., et al.](#) 2013). Systems developing in this region range from weak orographic lows to intense cyclones or even tropical-like cyclones with severe socio-economic impacts (Nissen [et al., et al.](#) 2010; Raveh-Rubin and Wernli, 2015). However, the term *Mediterranean cyclones* is typically used in the scientific literature to define a geographical origin rather than a cyclone category with specific dynamical characteristics. Indeed, the state-of-the-art is lacking a systematic analysis of the dynamics of these systems. In this regard, this study focuses on the most intense Mediterranean cyclones and in particular on the processes that turn these systems into severe storms.

Concerning the most intense Mediterranean cyclones, there is a general consensus that Mediterranean cyclogenesis is provoked by intruding upper-tropospheric systems that trigger baroclinic instability (Fita et al. 2006; Kouroutzoglou et al. 2011; Claud et al. 2010; Flaounas et al. 2015). Such systems typically correspond to potential vorticity (PV) streamers, equivalent to southern deviations of the Polar Jet that are a direct result of wave breaking over the Atlantic Ocean (Raveh-Rubin and Flaounas, 2017). After cyclogenesis takes place, baroclinic forcing may further intensify surface cyclones (e.g. Prezerakos et al. 2006; Lagouvardos et al. 2007) favouring, in parallel, convection in cyclone centres due to forced large-scale ascent. Recent studies used lightning observations to show that about one third of the most intense Mediterranean cyclones are associated with deep convection close to their centre (Galanaki [et al., et al.](#) 2016; Flaounas [et al., et al.](#) 2017). Due to latent heat release, convection serves as a net source of PV in the lower-to-mid troposphere and thus reinforces cyclonic circulations. Consequently, both baroclinic instability and convection may act in synergy to intensify Mediterranean cyclones (Fita [et al., et al.](#) 2006; Chaboureau [et al., et al.](#) 2012; Miglietta [et al., et al.](#) 2017). Nevertheless, the processes that play primary roles in the development and intensity of Mediterranean cyclones is still an open question.

Medicanes (portmanteau of the words *Mediterranean hurricanes*) are rare cyclonic systems (1-3 occurrences per year) and have been proposed in several past studies to share similar dynamics with tropical cyclones (e.g. Emanuel, 2005; Fita et al. 2007). This similarity is mainly attributed to the visual characteristics of medicanes, i.e. a cloudless "eye" in the centre of a spiral cloud coverage (Tous and Romero 2013), but also due to cyclone phase diagrams that showed axisymmetric warm core structures (Cavicchia [et al., et al.](#) 2014). To our knowledge the identification of all known medicane cases has been based on subjective criteria and, in particular, on the observation of cloud coverage satellite images (Tous and Romero 2013; Nastos [et al., et al.](#) 2017). However, the similarity of medicanes to their tropical counterparts is not clear from the perspective of atmospheric processes. Although deep convection and wind-induced surface heat exchanges have been shown to sustain the cyclonic circulation of several medicane cases (e.g. Miglietta et al. 2013; Dafis et al. 2018; Miglietta and Rotunno, 2019), an upper-tropospheric PV streamer or a cut-off is typically located close to medicanes (Tous and Romero, 2013; Nastos [et al., et al.](#) 2017). This suggests a non-negligible contribution of baroclinic forcing to medicane development. Indeed,

several known medicane cases have been shown to develop due to the combined effect of both baroclinic and diabatic processes (Fita ~~et al., et al.~~ 2006; Chaboureau ~~et al., et al.~~ 2012; Carrio ~~et al., et al.~~ 2017; Miglietta and Rotunno, 2019).
65 Furthermore, a recent case study of a medicane showed that the axisymmetric warm core structure may be reached when there is a warm seclusion and a relatively weak cut-off system (Fita and Flaounas, 2018). This raises a question of the driving processes of medicanes and whether other Mediterranean cyclones, not diagnosed as medicanes, are diabatically driven by convection in their centre.

70 Baroclinic instability and convection are two of the most common processes ~~that affect~~~~that universally affect all the~~ dynamics of extratropical cyclones. However, Mediterranean cyclones are also affected by the complex geography of the region. For instance, the relatively small size of the Mediterranean Sea and the arid areas of North Africa have an indirect effect on cyclones by limiting their supply of water vapour (Flaounas ~~et al., et al.~~ 2019). On the other hand, the high mountains that surround the Mediterranean basin have a direct effect on cyclones by interacting with the air flow and
75 triggering PV streamers of topographic origin. Such streamers are usually referred to as PV banners (e.g. Aebischer and Schär, 1998) and correspond, i.e. to filaments of diabatically produced PV at the wake of mountains (Rotunno et al. 1999; Epifanio and Durran, 2002; Schar et al. 2003; Flamant et al. 2004). The potential role of such PV production in Alpine cyclogenesis, one of the most cyclogenetic areas of the Mediterranean basin, has long been suggested to be important (e.g. Bleck and Mattocks, 1984) and have been recently reviewed by Buzzi et al. (2020). Especially when combined with the
80 upper-tropospheric PV streamer, PV banners ~~are suggested to~~may even reinforce cyclones (Tsidulko and Alpert, 2001; McTaggart-Cowan et al. 2010a, 2010b). Therefore, the uniqueness of the region adds a higher degree of complexity to the processes implicated in Mediterranean cyclone development.

Cyclones may be conceptualized as the outcome of PV anomalies that extend to different atmospheric levels. ~~(e.g. Chagnon et al., 2013; Martínez-Alvarado et al., 2016; Büeler and Pfahl, 2017; Attinger et al., et al. 2019).~~ As a diagnostic variable, PV has been especially helpful to provide deep insights into the development mechanisms of different types of cyclones, including tropical cyclones (e.g. Möller and Montgomery, 2000; Chan et al. 2002) and mid-latitude storms (e.g. Boettcher and Wernli, 2011). As a result, the quantification of the relative contribution of different atmospheric processes to the total atmospheric PV is equivalent to delineating the contribution of these processes to cyclone dynamics. For instance, Gray (2006), Chagnon and Gray (2015) and more recently Spreitzer et al. (2019) used PV budget diagnostics to analyse the impact of diabatic processes to the dynamical structure of the upper troposphere. In the same direction, several studies used PV-budget diagnostics to investigate the role of convection and more generally of latent heat in the dynamical structure of cyclones (Martínez-Alvarado et al. 2016; Büeler and Pfahl, 2017; Attinger et al. 2019). By extension, piecewise PV
90 inversion of these individual contributions is expected to reproduce the partition of the low-level wind circulation that is related to each atmospheric process, i.e. ~~to~~ the contribution of these processes to the cyclone itself (defined as a vortex).

Applying piecewise PV inversion to individual PV tracers is a well-established approach to quantify the relative contribution of different processes to the development of cyclones (e.g. Davis and Emanuel 1991; Wu and Emanuel 1995; Stoelinga, 1996; Huo et al. 1999; Bracegirdle and Gray 2009; Schlemmer et al. 2010; Seiler, 2019). For instance, Stoelinga (1996) and Ahmadi-Givi et al. (2004) quantified the contribution of diabatic processes to the intensification of two mid-latitude storms. The use of PV-budget diagnostics or piecewise PV inversion to perform a process-based classification of cyclones is still missing from the state-of-the-art. Such an approach would extend the analysis of Čampa and Wernli (2012) who analysed the vertical PV profile of extratropical cyclones of different geographical origin and would be complementary to earlier efforts by Deveson et al. (2002), followed by Plant et al. (2003) and Gray and Dacre (2006) where extratropical cyclones were classified to three types according to the contribution of upper and lower troposphere to their development. Despite the considerable advancements in analysing and classifying tropical and mid-latitude cyclones from a PV perspective, Mediterranean cyclones have been rarely been the object of comprehensive studies in which their dynamical structure is analysed in a climatological context. In this study, we use regional climate modelling to simulate 100 of the most intense Mediterranean cyclones and we analyse their dynamics using PV-budget diagnostics and piecewise PV inversion. In particular, we address the following specific objectives:

- 1 - Quantify the relative contribution of different processes to the dynamics of Mediterranean cyclones, defined by the spatial variability of PV during their mature stage; and
- 2 - Delineate the role of different processes in cyclone development and thus provide insights into the yet uncharted theoretical continuum between tropical-like cyclones (i.e. “purely” diabatic) and baroclinic lows.

In the next section, we introduce the modelling, diagnostics and PV inversion methods that we used to address these objectives. Section 3 places the simulated cyclones in a climatological context and then sections 43 and 54 present our main results. Finally, section 56 hosts a summary of the analysis and the conclusions of this study.

2 Methods

2.1 Cyclone cases selection and modelling approach

To build a climatology of cyclones, we applied the cyclone tracking method of Flaounas et al. (2014) to the hourly relative vorticity fields at 850 hPa, taken from the ERA5 reanalysis of the European Centre of Medium-Range Forecasts in the 10-year period from 2008 to 2017 (Copernicus Climate Change Service; C3S, 2017). ~~Out of all tracked cyclones, we retained the 100 most intense ones after subjectively excluding cyclonic circulation features that did not present clear enclosing mean sea level pressure contours.~~ Out of all tracked cyclones, we retained the 100 most intense ones. However, the tracking method is sensitive to identifying long lasting relative vorticity local maxima and in several cases these maxima lacked a clear

130 cyclonic structure, i.e. they did not correspond to structured wind vortices. These tracked features were characterised as "false positives" and were subjectively excluded from our dataset of 100 most intense cyclones. Our main criterion to retain a system is the existence of a clear relative vorticity maximum that is outlined by enclosing mean sea level pressure contours. The relative vorticity and mean sea level pressure fields at the time of cyclones' mature stages are shown in the supplementary material. Out of 100 cyclones, ten cases were diagnosed as medicanes in recent studies (Nastos [et al., et al.](#) 2018; Di Muzio [et al., et al.](#) 2019). These cases are shown as cases 7, 14, 24, 28, 46, 56, 65, 69, 75, 76 of the supplementary material.

135

A simulation has been performed for each individual cyclone, using the Weather, Research and Forecasting model (WRF, version 4.0; Skamarock et al. 2008). All simulations have been initialised 36 hours before the cyclone's mature stage, at the time of maximum relative vorticity in ERA5, and last for 72 hours in total. Initial and boundary conditions have been taken from the hourly fields of the ERA5 reanalysis at 0.25 degrees of grid spacing. All the simulations used a common domain (shown in Fig. 1) which is composed of 350x200 grid points with a grid spacing of 15 km. In the vertical, [the model top was set at 50 hPa and](#) we used 34 hybrid eta levels that follow terrain topography in the lower troposphere, gradually transitioning to constant pressure levels in the higher troposphere. The model parametrisation was also common for all simulations and consists of the Single-Moment five-class microphysics scheme (WSM5; Hong et al. 1998, 2004), the Dudhia shortwave radiation scheme (Dudhia 1989), the Rapid Radiative Transfer Model longwave radiation scheme (Mlawer et al. 1997), the Yonsei University planetary boundary layer (PBL) scheme (Hong [et al., et al.](#) 2006; Hong, 2010), and the Kain-Fritsch convective parameterization (Kain 2004). [This combination of physical parametrisation was shown to adequately reproduce Mediterranean cyclones \(e.g. Flaounas et al. 2019; Fita and Flaounas, 2018; Miglietta and Rotunno, 2019\). Especially the use of a five-class microphysics scheme was shown to yield simulations with a small track error in two medicane cases, when compared to the use of other microphysical parametrisations \(Miglietta et al. 2015; Pytharoulis et al. 2018\). It is however noteworthy that the absence of hail or graupel results in unrealistic representation of the microphysical properties of deep convective clouds.](#)

150

2.2 Potential vorticity tracers

A module has been implemented into WRF that calculates PV at every model time-step according to the following equations:

155

$$Q = \frac{-g}{\mu} \xi_a \cdot \nabla \theta \quad (\text{Eq. 1}),$$

$$\frac{dQ}{dt} = \frac{-g}{\mu} \left(\xi_a \cdot \nabla \dot{\theta} + \nabla \times F \cdot \nabla \theta \right) \quad (\text{Eq. 2}),$$

where Q is PV, g is the acceleration due to gravity, f is the Coriolis parameter, μ is the atmospheric column dry air mass, ξ_a is the three dimensional field of absolute vorticity, \mathbf{F} is a two dimensional field that represents a mixing wind acceleration term that includes surface friction, turbulent and diffusive processes, θ is potential temperature and $\dot{\theta}$ is its time derivative. PV is calculated on eta (η) model levels that range from 0 to 1 ($\eta = (P_\eta - P_s)/\mu$, with P_η representing pressure on eta level η and P_s the surface pressure). Equations 1 and 2 follow the formulation of Cao and Xu (2011) that This choice has the advantage of avoiding wind and temperature interpolations—or approximations—in atmospheric levels other than the ones in model outputs (Cao and Xu 2011). After calculating total PV, a PV-budget diagnostic is activated at every model time step. This diagnostic follows previous methodological approaches where the total atmospheric PV is decomposed into several non-conserved partitions and one conserved partition (Lamarque and Hess, 1994; Stoelinga, 1996; Gray, 2006). All these PV partitions are treated by the model as scalars, subject to advection and thus hereafter are referred to as PV tracers.

Each non-conserved PV tracer is attributed to one parametrised process that acts as a source or a sink of atmospheric PV. Overall, six non-conserved PV tracers are used. These derive from latent heat release by microphysics (q_{mp}) and convection (q_{cu}) parametrisations; turbulent fluxes of temperature by the planetary boundary layer parametrisation (q_{bl}); atmospheric warming/cooling by shortwave (q_{sw}) and longwave (q_{lw}) radiation; and a mixed term of momentum acceleration applied to air masses (q_{mo}), which is also provided by the boundary layer parametrisation. As explained below, the conserved PV tracer (q_{co}) represents the contribution of the large-scale flow to PV and it is only subject to advection. In contrast, non-conserved PV tracers are, in addition to advection, also subject to accumulations of net gains and losses of PV. These gains and losses of PV represent the right-hand side terms of Eq. 2 and derive from the net temperature and momentum forcings of the model physical parameterisations to the primitive equations. More precisely and following the annotation in Stoelinga (1996), each PV tracer is calculated at every model grid point as follows:

$$q_x(\tau) = q_x(0) + \int_0^\tau (-\mathbf{v} \cdot \nabla q_x + G_x) dt \quad (\text{Eq. 3}),$$

$$\text{where} \quad G_x = \frac{-g}{\mu} \xi_a \cdot \nabla T_x \quad (\text{Eq. 4}).$$

$$G_{(x=mo)} = \frac{-g}{\mu} (\nabla \times \mathbf{F}) \cdot \nabla \theta$$

\mathbf{v} represents the three-dimensional wind field and q_x represents any PV tracer at a given time step τ , related to process x (co , mp , cu , bl , sw , lw or mo). The term $q_x(0)$ represents the value of PV tracers in the model's initial conditions, while PV sources and sinks are represented by the term G_x . The latter is calculated by Eq. 4, using the full model wind outputs and the

temperature tendencies from model parametrisations (represented by the term T_x). Finally, it is noteworthy that G_{mo} is calculated using F from the boundary layer parametrisation and the full model θ .

190

Given that q_{co} represents the PV partition that is related to the large-scale flow, the total atmospheric PV at the initial time ($q_{co}(0)$ in Eq. 3) and boundary conditions are entirely attributed to q_{co} , while the term G_x is set to zero in Eq. 3. As a result, q_{co} is constantly introduced into the domain by prescribing the full atmospheric PV of the large-scale flow—regardless of whether it is produced by non-conservative processes—and it is interpreted as the PV partition that incorporates the dynamics that are "external" to the simulation domain (i.e. the Mediterranean region). On the other hand, non-conserved PV tracers at initial time ($q_x(0)$ in Eq. 3) and boundary conditions are all set to zero. This approach suggests that diabatic processes start taking place only after the initial conditions of the simulation and only within the simulation domain. Setting the initial conditions 36 hours before the cyclone's mature stage is a rather sufficient period to capture cyclogenesis. Indeed, the characteristic lifetime of Mediterranean cyclones is of the order of two days (Flaounas et al. 2015). Therefore, non-conserved PV tracers accumulate PV sources and sinks due to specific processes (G_x term in Eq. 4) and thus "memorize" the processes that lead cyclones to intensify. At any given model time step, the total PV equals the sum of all tracers:

200

$$Q = q_{co} + q_{mp} + q_{cu} + q_{bl} + q_{sw} + q_{lw} + q_{mo} + q_r \text{ (Eq. 5).}$$

205 As for PV tracers, potential temperature may be also treated as a tracer (θ tracers hereafter), subject to advection and to accumulations of temperature tendencies (T_x) as produced from the physical parametrisations. Consequently, one conserved and several non-conserved θ tracers may be considered, similarly to PV (except for momentum as that does not apply direct changes to temperature):

210

$$\theta = \theta_{co} + \theta_{mp} + \theta_{cu} + \theta_{bl} + \theta_{sw} + \theta_{lw} + \theta_r \text{ (Eq. 6).}$$

~~Finally, it is noteworthy that m~~Models are not numerically bound to conserve PV at every model time step (Saffin et al, 2016). Therefore, in Eqs 5 and 6, a residual occurs represented by the term with the subscript r . This residual can be mainly attributed to the dynamical core of the model and to adjustments made by the model to the prognostic variables in the primitive equations. However, the residual term does not attain large values and its average within a 300 km radius around cyclone centres is of the order of 0.2 PVU. Higher residuals are only found in the first model level with a median of -0.17 PVU, but ranging between -0.5 and -1 PVU in 14 cases out of 100. Finally, it is noteworthy that version 4.0 of the WRF model uses moist potential temperature to allow consistent treatment of moisture in the calculation of pressure in its dynamical core. For the purposes of our analysis we could also use the moist potential temperature as a basis for calculating a generalized PV field for a moist atmosphere. Schubert et al. (2001) showed that such a moist PV field still complies with

220

the invertibility principle and thus it incorporates all information that defines the atmospheric state. However, to be consistent with previous studies, we performed our analysis with all PV fields in WRF being calculated with potential temperature.

2.3 Piecewise PV inversion

225 Applying piecewise PV inversion to individual PV tracers is a well established approach to quantify the relative contribution of different processes to the development of cyclones (e.g. Davis and Emanuel 1991; Wu and Emanuel 1995; Stoeligna, 1996; Huo et al. 1999; Bracegirdle and Gray 2009; Schlemmer et al. 2010; Seiler, 2019).

230 The contribution of different processes to cyclonic wind circulation has been determined by inverting individual PV anomalies under nonlinear balance (Charney 1955; Davis and Emanuel, 1991). Following StoelignaStoeligna (1996), every PV tracer may be considered as a PV anomaly, except for q_{co} , which is approximately equal to the full PV field in the upper troposphere. Therefore, q_{co} anomalies have been defined as differences from three-day averages of q_{co} , i.e. the duration of each simulation. A three-days time period might be short to define PV anomalies in cases of slowly evolving upper tropospheric systems. However, anomalies need to be consistently defined for all 100 cyclones and Mediterranean cyclones
235 are systems of relatively short lifespan. Therefore, we consider that a three-days average is a fair compromise for the application of piecewise PV inversion in this study. To perform piecewise PV inversion we used the "subtraction approach" of Davis (1992) which has already been applied in past studies to separate the advective contributions of upper-tropospheric and low-tropospheric PV and potential temperature anomalies to upper-tropospheric PV anomalies (Piaget et al., et al. 2015; Teubler and Riemer 2016; Schneidereit et al., et al. 2017 and Teubler and Riemer 2020). According to this approach, the
240 partition of the atmospheric state (wind and geopotential) due to individual PV tracers (hereafter referred to as the balanced fields) is obtained by subtracting the balanced state related to the full PV inversion from the state related to the full PV minus the individual PV tracers. For instance, the balanced wind field, \bar{U} , due to q_{mp} is equal to

$$\bar{U}(q_{mp}) = \bar{U}(Q) - \bar{U}(Q - q_{mp}) \text{ (Eq. 7).}$$

245 All model outputs were interpolated to a regular longitude-latitude grid with a spacing of 1 degree and to pressure levels ranging from 950 to 100 hPa with a step of 50 hPa. Finally, following Stoeligna (1996), the piecewise PV inversion of individual PV tracers was performed by prescribing the relevant θ tracers in the boundary conditions at the top and bottom atmospheric pressure levels as Neumann boundary conditions (925 and 125 hPa respectively), except for q_{mo} which is not
250 associated with a relevant θ tracer.

Numerous sensitivity tests have been performed to converge on a universal application of piecewise PV inversion for all 100 simulations. The method results have first been evaluated on the basis of accurate cyclone locations, by comparing cyclone locations in the balanced relative vorticity fields (calculated using the balanced wind outputs from PV inversion) to the ones in the model outputs. Despite the use of interpolated fields, the balanced wind fields showed remarkably accurate cyclone locations with an average bias of less than 1 degree. In addition, the validity of the method has been evaluated through the comparison in the cyclone centres of (1) the balanced vorticity field at 850 hPa after inverting the full PV with (2) the sum of the balanced vorticity fields at the same pressure level after inverting the conserved and all non-conserved PV tracers. After applying PV inversion we have retained 94 cases. Results showed that for these cases, the ratio between the two fields ranged between 0.9 and 1.1. This suggests that the inversion processes of individual PV tracers may accurately explain the contribution of the different processes to cyclonic circulation by at least 90%. In the other cases six cases, the ratio exceeded 1.1, presenting a maximum of 1.4.

3. The 100 cyclone cases in a climatological context

Figure 1 shows the locations and depths of the 100 simulated cyclones at the time of their mature stage (i.e. when they reach maximum relative vorticity at 850 hPa). The cases tend to be concentrated over maritime areas, especially over the western ~~and eastern~~ Mediterranean Sea. Fewer systems are found ~~over the Aegean Sea~~ in the eastern Mediterranean, while a small group of systems reach their maximum intensity over the continental areas of North Africa. Concerning the seasonal distribution of their occurrence, Fig. 2a shows that most cyclones occur in winter with fewer in autumn and spring, while none have been observed in summer with ~~an~~ the exception of three cases in June. Although there is a certain arbitrariness in the selection of the cases, the spatial and seasonal distribution of the 100 cases is consistent with previous climatological studies of intense Mediterranean cyclones (Campins et al. 2011; Flaounas et al. 2018).

Similarly, the mean sea level pressure in the centre of our 100 cyclones is comparable to relevant climatological distributions. Indeed, Fig. 2b shows that the 5th, 50th and 95th simulated cyclones present values of 985, 998 and 1005 hPa, respectively. This is comparable to the results of Flaounas et al. (2018) where intense cyclones were tracked in different regional climate models during a 20-year period. In their results the 5th, 50th and 95th percentiles of cyclones depth roughly ranged between 1000-1005 hPa, ~~985-990~~ 995-1000 hPa and 985-990 hPa, respectively (their figure 8). Therefore, our cyclone cases may be plausibly considered as representative of the climatology of intense Mediterranean cyclones.

Finally, to gain an insight into the dynamics of the 100 cyclones, Fig. 3 shows the composite averages of equivalent potential temperature and wind at 850 hPa for all 100 simulated cyclones, as well as their mean sea level pressure and PV at 300 hPa. Composite fields are spatially centred on the cyclone centres and temporally centred at the time of cyclones' mature stage (i.e. when cyclones have their maximum relative vorticity). Figure 3 clearly shows that cyclones develop in a baroclinic

environment where equivalent potential temperature depicts frontal structures. The composite PV streamer is also clear at
285 300 hPa, presenting a thinning north-south structure mostly overlapping with the composite cyclone's cold sector. The
highest PV values are slightly to the south of the composite cyclone's centre. These results are in fair agreement with
previous studies on the dynamical structure of 200 intense Mediterranean cyclones, tracked in a 20-year climatology
(Flaounas [et al., et al.](#) 2015; their figure 3),

4. A process-based anatomy of Mediterranean cyclone dynamics

290 In the following, we analyse the dynamical structure of the 100 cyclones using vertical and horizontal profiles of the full
atmospheric PV and PV tracers. Vertical profiles have been averaged within a radius of 100 km around cyclone centres. This
radius is small compared to the characteristic size of the cyclones, but sufficient to encompass PV anomalies in their centres.
This analysis will provide insight into the processes that form PV anomalies close to cyclone centres and thus play an
important role in their dynamics.

295 4.1 Dynamical structure of cyclones: conserved and non-conserved PV

~~In the following, we analyse the dynamical structure of the 100 cyclones using vertical and horizontal profiles of the full
atmospheric PV and PV tracers. Vertical profiles have been averaged within a radius of 100 km around cyclone centres. This
radius is small compared to the characteristic size of the cyclones, but sufficient to encompass PV anomalies in their centres.
This analysis will provide insight into the processes that form PV anomalies close to cyclone centres and thus play an
important role in their dynamics.~~

- *Total PV vertical profile*: All cyclones present a similar vertical PV profile (Fig. 4a) with a local maximum of the order of
1.3 PVUs at 850 hPa and a dramatic increase of PV above 400 hPa. Campa and Wernli (2012) showed that such a profile is
typical for extratropical cyclones where the local maximum of PV at lower levels is related to diabatic heating and high PV
305 values in the upper troposphere are related to PV streamers; or, more generally, stratospheric air masses that overlap with
cyclone centres. Indeed, the PV values in Fig. 3 are consistent with the ones in the vertical profiles of Fig. 4.

- *Conserved PV*: The PV streamer in Fig. 3 corresponds to stratospheric air that intrudes into the Mediterranean and thus is
prescribed in the boundary conditions of the domain. As a result, the conserved component of PV in levels above 400 hPa
310 (q_{co} ; Fig. 4b) presents values comparable to the total PV (Fig. 4a). Indeed, the fields of both q_{co} and total atmospheric PV
present similar amplitude and shape when comparing their horizontal cross sections at 300 hPa in Figs 5a and 3,
respectively.

In contrast to the upper troposphere, the vertical profile of q_{co} tends to be rather invariant below 400 hPa, presenting an average value of the order of 0.5 PVUs (Fig. 4b). However, q_{co} at low levels exceeds 1 PVU in several cases, especially in the first model levels. Figure 5b shows that higher values of q_{co} are related to a low-level PV streamer that overlaps with the cold sector of the cyclones and wraps cyclonically around their centre. Being only subject to advection, q_{co} is strongly dependent on the total PV in the initial conditions of the simulations. The origin of this low-level streamer is plausibly related to prominent values of PV close to orographic features. Indeed, the initial conditions of PV at 850 hPa are consistent with Fig. 5b, showing values of the order of 0.5 PVUs close to mountains (not shown).

The role of mountains in acting as PV sources has been demonstrated previously by Schär and Durran (1997). The authors used idealised simulations to show that PV anomalies may be generated in the wake of mountains due to internal dissipation of the air flow. Therefore, mountains plausibly act as a [reservoirsource](#) of relatively high PV values that potentially reinforces cyclones. This hypothesis has been confirmed in a case study by McTaggart-Cowan et al. (2010b). The authors performed several sensitivity tests on the dynamics of a medicane that developed close to the Alps. Results showed that removal of the mountains had a detrimental effect on cyclogenesis, inhibiting its occurrence.

- *Non-conserved PV*: The non-conserved component of PV is represented by the sum of all PV tracers apart from q_{co} (i.e. $q_{mp}+q_{cu}+q_{bl}+q_{sw}+q_{lw}+q_{mo}$). Figure 4c presents a vertical alternation of signs: negative values are observed from the surface (around 975 hPa) to 950 hPa, thereafter positive values exist from 950 to 600 hPa, with a maximum at 850 hPa, and finally, negative values are observed between roughly 600 and 300 hPa. From the point of view of atmospheric dynamics, this alternation of signs suggests that diabatic processes tend to act as a sink of PV close to the surface, as a source in the lower-to-mid troposphere, and again as a sink in the upper troposphere. It is noteworthy that q_{sw} presents especially low values with an average in cyclone centres that is close to zero (not shown). Therefore, analysis solely based on q_{sw} is omitted hereafter.

In the lowest layer (from surface to 950 hPa), the negative values of the non-conserved PV tracers tend to get balanced by the vertically invariant positive values of q_{co} (Fig. 4b), resulting in a PV average close to zero (Fig. 4a). In the middle layer (from 950 to 600 hPa), the diabatic processes are strongly characterized by the local maximum at 850 hPa (Fig. 4a). This local maximum is expected to be further enhanced by the typically positive values of low-level q_{co} (Fig. 5b). However, Fig. 5d shows that the low-level local maximum of non-conserved PV tracers tends to be concentrated close to cyclone centres. This characteristic is in contrast to the broader, low-level conserved PV-streamer in Fig. 5b, suggesting that diabatic processes are expected to be directly related to the low-level cyclonic circulation. Finally, in the upper troposphere (from 600 to 300 hPa), the non-conserved PV tracers act as a rather weak sink of PV. Indeed, Fig. 5c presents significantly weak values when compared to the ones shown in Fig. 5a. Nevertheless, non-conserved PV tracers tend to erode the amplitude of the upper-level PV streamer and thus partly contribute to the thinning of its meridional structure in Fig. 3. It is noteworthy that values in Figs 5a and 5c are rather smoothed since they correspond to composite averages. In single cases, the shape of PV

streamers is expected to present sharp edges due to diabatic processes that may locally act as strong sinks of PV (e.g. Gray, 2006; Chagnon et al. 2013; Spreitzer [et al., et al.](#) 2019).

350 | 4.2 Dynamical structure of cyclones: ~~processes~~ contribution of processes to non-conserved PV

- *Latent heat:* Figure 4d shows that latent heating acts as a major source of PV within the lower and mid troposphere. Indeed, positive values of $q_{mp}+q_{cu}$ are observed from surface to 500 hPa. On the other hand, $q_{mp}+q_{cu}$ presents rather weak negative values from 500 to 300 hPa, acting as a sink of PV. As deduced from Eq. 2, this alternation of sign in Fig. 4d is related to the vertical gradient of latent heating. Indeed, Fig. 6a shows that $\theta_{mp}+\theta_{cu}$ presents its maximum value at around 500 hPa, i.e. the level where its vertical gradient turns from positive to negative.

Therefore, the vertical profile of $\theta_{mp}+\theta_{cu}$ is fairly consistent with that of $q_{mp}+q_{cu}$. In contrast, the vertical profile of instantaneous tendencies of latent heat release ($T_{mp}+T_{cu}$) in Fig. 6b is not consistent, presenting its local maximum at 800 hPa, i.e. in lower levels than 500 hPa as is the case for $\theta_{mp}+\theta_{cu}$ in Fig. 6a. Given that $\theta_{mp}+\theta_{cu}$ accumulate and transport instantaneous tendencies of latent heat release, this inconsistency can be attributed either ~~uniquely one of~~, or most probably to the combination of two possible effects: convection is weakened at the time of the cyclones' mature stage when compared to previous times and high $T_{mp}+T_{cu}$ values are upwardly advected within the convective atmospheric column. To clarify this issue, we used the tracks of all 100 cyclones and produced similar profiles to Fig. 6b at a time 12 hours before the cyclones' mature stage. Results showed that the local maximum of $T_{mp}+T_{cu}$ was indeed located at an atmospheric level close to 500 hPa (not shown), plausibly suggesting that this inconsistency is due to convection being weakened when cyclones reach their mature stage. As a result, the $q_{mp}+q_{cu}$ tracers are capable of including the "dynamical memory" of convection from previous times, i.e. convection may still reinforce cyclones in subsequent times to its maximum activity. This result is consistent with Miglietta [et al., et al.](#) (2013) and Galanaki [et al., et al.](#) (2016) who found that the maximum intensity of Mediterranean cyclones is preceded by deep convection in cyclone centres, but also with Price [et al., et al.](#) (2009) and Whittaker [et al., et al.](#) (2015) who showed that the maximum intensity of tropical cyclones comes in subsequent times, after the peak of convective activity. It is also noteworthy that $\theta_{mp}+\theta_{cu}$ presents negative values from the surface to 850 hPa. Recent results on the water budget of Mediterranean cyclones showed that these negative values are related to rain evaporation (Flaounas [et al., et al.](#) 2019).

375 | Finally Regarding the vertical level of latent heat contribution to cyclone dynamics, Fig. 4d shows that $q_{mp}+q_{cu}$ presents its maximum at 900 hPa, i.e. below the local maximum of non-conserved PV at 850 hPa (Fig. 4c). Figure 7a shows that $q_{mp}+q_{cu}$ acts indeed as a significant source of PV at 900 hPa. However, the positive values of $q_{mp}+q_{cu}$ are spread along a southwest-to-northeast direction, i.e. they follow the frontal structure of the cyclones where convection is favoured. Comparing this spread of positive values of $q_{mp}+q_{cu}$ to the narrowly concentrated positive values for all non-conserved PV tracers in Fig. 5de, it is rather evident that other PV tracers act as sinks of PV, tending to counterbalance the PV sources due to latent heat, especially

along the cyclone fronts. It is noteworthy that although frontal structures are evident in Figs 3 and 7, their exact location, intensity and vertical PV profile are case dependent. As a result, latent heating-produced PV maxima along the cold fronts and their associated PV tendencies right below and above these maxima (Lackmann 2002) will be at some extent inconsistent between the cyclones. This will affect values in the composite averages of Fig. 7a. Such inconsistencies are also expected in composite averages of other PV tracers close to the surface (e.g. Figs 7c and 7e), where PV is typically negative in the cold sector, due to small or negative static stability, and positive values in the warm sector (Vanniere et al. 2016).

Finally, despite the important role of sea surface temperature (SST) in modulating convection (e.g. Miglietta et al. 2011; Pytharoulis, 2018), in this study we found no seasonal dependence or direct relationship between SSTs and $q_{mp}+q_{cv}$ in low atmospheric levels (not shown). This is partly due to the diversity of the 100 cyclones' dynamics and partly due to the fact that all systems occur in different regions and different seasons. For instance, convection might differ between two systems with similar underlying SSTs due to different large scale forced ascent. Similarly, two cyclones that occur in the same season, but in different areas are very likely to present considerable differences in their underlying SSTs and consequently to convection. In fact, the investigation of the role of SST in cyclone dynamics from the perspective of PV tracers is not a straightforward issue due to the trait of $q_{mp}+q_{cv}$ PV tracers to accumulate the diabatically produced PV. As a result, an air mass that has an impact on cyclone dynamics might have experienced the PV gains due to latent heat in remote areas to cyclones centre.

- *Momentum exchanges with environment:* The vertical profile of q_{mo} is highly variable among the 100 cyclone cases (Fig. 4e). At the pressure level of 850 hPa, there are 73 cases where q_{mo} is positive and thus functions as a source of PV (with a maximum of the order of 2 PVU) and 27 cases where q_{mo} is negative and thus functions as a sink (with a minimum of the order of -0.5 PVU). This is consistent with Stoelinga (1996) who analysed the role of q_{mo} in the development of a case study of a cyclone development and showed that indeed this PV tracer may act as both a source and a sink of PV. To understand the ambiguous contribution of q_{mo} to cyclones development, we expand Eq. 4 into the following three terms:

$$G_{mo} = \frac{-g}{\mu} \left[\left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) \cdot \frac{\partial \theta}{\partial \eta} + \left(\frac{\partial F_x}{\partial \eta} \right) \left(\frac{\partial \theta}{\partial y} \right) - \left(\frac{\partial F_y}{\partial \eta} \right) \left(\frac{\partial \theta}{\partial x} \right) \right] \text{ (Eq. 8)}$$

Adamson et al. (2006) used numerical simulations to analyse in depth the role of q_{mo} in the development of an idealized cyclone. In their analysis, the first term is responsible for the "Ekman generation" of PV and is proportional to the contribution of the PBL parametrisation to the cyclone's relative vorticity. As such, this term typically corresponds to a PV sink due to wind deceleration by frictional forces, i.e. frictional forces impose an anticyclonic circulation. On the other hand, the second and third term are responsible for the "baroclinic" generation of PV and are expressed by the dot product

between the horizontal component of the ~~rotational~~curl of \mathbf{F} and the horizontal gradient of θ . In fair accordance with Adamson et al. (2006), Fig. 7b shows that positive values of q_{mo} tend to be concentrated on the northeast side and close to the centre of the cyclones and negative values tend to be concentrated on the southwest side. The baroclinic generation of PV (i.e. the two last terms in Eq. 8) were shown by Adamson et al. (2006) to correspond to a significant PV source, where positive anomalies were suggested to be progressively advected by warm conveyor belts upwards and eventually polewards and towards the west side of cyclones centres. Therefore, the function of q_{mo} as a source or a sink of PV in cyclones' centre depends on the vertical component of the curl of \mathbf{F} , on the magnitude of baroclinicity and its relative location to the horizontal component of the curl of \mathbf{F} , but also to the advection of q_{mo} in cyclone centre. However, the relatively small size of the Mediterranean basin, its sharp land-sea transitions and the short lifetime of cyclones may not always allow a clear formation of frontal structures. It is thus plausible to suggest that the contribution of q_{mo} to cyclone development may vary significantly from case to case with a general tendency to reinforce cyclones, i.e. the average in Fig. 4e presents positive values at 850 hPa. An exemplary case study is used in section 5.4 to provide further details on the baroclinic generation of PV in q_{mo} . Figure 7b provides further insight into the ambiguous sign of q_{mo} . Positive values tend to be concentrated on the northeast side and close to the centre of the cyclones and negative values tend to be concentrated on the southwest side. This finding is consistent with Adamson et al. (2006) where q_{mo} is shown to produce positive PV anomalies in the northeast side of an idealized cyclone, developed in a dry environment. In a case study of an extratropical cyclone, Stoelinga (1996) analysed the role of q_{mo} and showed that indeed this PV tracer may act as both a source and a sink of PV.

~~The first term is related to a clear PV sink due to wind deceleration by frictional forces (imposing an anticyclonic circulation). On the other hand, the second and third term is expressed by the dot product between the horizontal component of the rotational of \mathbf{F} and the horizontal gradient of θ . Therefore, q_{mo} may act as a source of PV that reinforces the low-level cyclone. However, this source depends on the direction of the fronts and wind deceleration. The relatively small size of the Mediterranean basin, its sharp land-sea transitions and the short lifetime of cyclones may not always allow a clear formation of frontal structures. It is thus plausible to suggest that the contribution of q_{mo} to cyclone development may vary significantly from case to case with a general tendency to reinforce cyclones, i.e. the average in Fig. 4e presents positive values at 850 hPa.~~

- *Turbulent fluxes of temperature*: The vertical profile of q_{bl} (Fig. 4f) shows that turbulent fluxes of temperature tend to act as a source of PV within the very first model levels and as a sink right above. Indeed, Fig. 7c shows that q_{bl} at 1000 hPa presents overall positive values, mostly concentrated along the cold front and close to the composite cyclone centre. Clearly, the values of q_{bl} in Fig. 7c are unrealistically high, compared to typical values of total PV in the troposphere. Such high values may be attributed to the sharp vertical gradient of temperature fluxes within the surface layer, i.e. within the two first model levels. Indeed, Fig. 6b shows a positive vertical gradient in the average profile of temperature turbulent fluxes (T_{bl}) in the first model levels. However, it is noteworthy, that the vertical profile of T_{bl} is subject to diurnal variabilities of the PBL height affecting thus its vertical gradient and the magnitude of q_{bl} . ~~It is also noteworthy that~~ In addition, q_{bl} presents a high

variability in amplitude and sign over the cyclone centres in Fig. 4f. It acts as a source of PV in the first model levels for 74 cases and as a sink of PV for the remaining 26 cases. For these 26 cases, the composite average of q_{bl} at 1000 hPa presents a structure consistent with that shown in Fig. 7c (not shown). However, the cyclone centre in these cases is slightly displaced
450 with respect to the area where q_{bl} presents high values. Consequently, the vertical profiles of q_{bl} in Fig. 4f are relatively sensitive to the locations of the cyclone centres. Consistent with the monotonic negative vertical gradient of θ_{bl} in Fig. 6a, q_{bl} acts as a sink of PV in the atmospheric levels above the PBL (roughly set at 950 hPa). Finally, negative values of q_{bl} at 850 hPa (Fig. 7d) tend to overlap with the area where $q_{mp}+q_{cu}$ presents high positive PV values in Fig. 7a. This result suggests that turbulent fluxes of temperature act as a sink of PV and thus tend to counteract the role of latent heat in reinforcing
455 cyclonic circulation.

- *Longwave radiative forcing of temperature*: The average vertical profile of q_{lw} (Fig. 4g) shows overall negative values within the whole atmospheric column, especially within the PBL. These negative values are consistent with negative vertical gradients observed for both θ_{lw} and T_{lw} in Figs 6a and 6b, respectively. As for q_{bl} , the vertical profile of q_{lw} in Fig. 4g presents
460 unrealistically low PV values within the first model levels, when compared to total PV values in the troposphere. This is due to a sharp decrease of T_{lw} within the first model levels. Indeed, positive values of T_{lw} in the first model level are attributed to high concentrations of water vapour that absorb longwave radiation, while negative values in the levels above are mostly related to cooling, due to thermal radiation of air masses. In fact, the average vertical profiles of q_{bl} and q_{lw} in Fig. 4 seem to be symmetric and thus tend to counter each other, especially within the very first model level. Indeed, the correlation
465 coefficient between the values of q_{bl} and q_{lw} at the first model level for all 100 cyclones is -0.96. Moreover, Figs 7c and 7e show similar spatial distributions for the fields of q_{bl} and q_{lw} . It is plausible that turbulent fluxes of water vapour and temperature are subject to the same physical mechanism and thus provoke similar vertical temperature gradients of T_{bl} and T_{lw} close to the surface. As a result, q_{bl} and q_{lw} are of similar amplitudes. On the other hand, and despite being highly correlated, q_{bl} and q_{lw} are not of equal absolute values. Differences between the 100 cyclone centres at the first model level
470 present an average of -1.3 PVUs and a standard deviation of 1.8 PVU. This suggests a high variability between the two fields, which in turn is strongly dependent on several factors such as background water vapour concentration and whether cyclone centres are located above the maritime or continental areas. In the atmospheric levels above the PBL, q_{lw} presents negative values mostly concentrated in cyclone centres (Fig. 7f). The sum of q_{bl} and q_{lw} (Fig. 4h) shows that the two PV tracers act as a sink of PV in the lower troposphere and thus are expected to weaken the cyclone.

475

- *Synopsis*: Upper-tropospheric PV is mostly composed of the conserved PV tracer (q_{co}), while the local maximum of PV anomaly at 850 hPa is mainly due to non-conserved PV tracers (Fig. 4c). Figure 8 summarizes the average PV tracer profiles
in the lower troposphere. Latent heating and momentum acceleration are expected to amplify the cyclone by acting as PV sources ($q_{cu}+q_{mp}$ and q_{mo} , shown by the blue and black lines, respectively), while turbulent fluxes and cooling due to
480 longwave radiation are expected to weaken the cyclone by acting as PV sinks ($q_{bl}+q_{lw}$, shown by a red line). The vertical

profile of latent heat PV tracers ($q_{cu}+q_{mp}$) has its maximum at 925 hPa, where latent heat release ($\theta_{cu}+\theta_{mp}$ in Fig. 6a) sharply increases before it reaches its maximum value at around 500 hPa. On the other hand, the profile of $q_{bl}+q_{lw}$ has its minimum at the level of 950 hPa, i.e. roughly at the top of the PBL. It is of no surprise that the vertical profiles of $q_{cu}+q_{mp}$ and $q_{bl}+q_{lw}$ in Fig. 8 tend to present vertically symmetric structures, longwave radiation and turbulent fluxes are expected to increase proportionally with the intensity of convection. This direct relationship is evident in the scatterplot of Fig. 9, where $q_{cu}+q_{mp}$ and $q_{bl}+q_{lw}$ are compared at the level of 850 hPa for all 100 cyclones and are shown to present a correlation coefficient of -0.87. However, by considering mean sea level pressure in cyclone centres as a measure of intensity (depicted by dot sizes in Fig. 9), it is rather evident that there is no systematic pattern that relates $q_{cu}+q_{mp}$ or $q_{bl}+q_{lw}$ to cyclones intensity. Therefore, the role of other processes in cyclone intensification is expected to be crucial, especially momentum mixing and baroclinic instability (i.e. the role of q_{co} to cyclonic circulation).

5. Relating conserved and non-conserved PV to cyclones intensity

In the previous section we used PV tracers to gain a deeper insight into the role of different atmospheric processes in forming the dynamical structure of cyclones. However, delineating the contribution of baroclinic and diabatic processes to cyclone development is not a trivial task. To this end, we use piecewise PV inversion to define the partition of the atmospheric state related to conserved and non-conserved PV. Figure 10 presents in a scatterplot the values of the balanced relative vorticity in cyclone centres at 850 hPa, calculated from the balanced flow fields after inverting q_{co} anomalies and the sum of all non-conserved PV tracers (i.e. $q_{mp}+q_{cu}+q_{bl}+q_{sw}+q_{lw}+q_{mo}$). Red circles in Fig. 10 depict the medicanes in our dataset. Values correspond to averages within 300 km from the cyclone centres at the time of their mature stage.

Figure 10 shows a negative linear trend (correlation coefficient -0.65) which suggests that the more a cyclonic circulation is due to baroclinic forcing (q_{co}), the less it is due to diabatic processes. In fact, the majority of cases—about 60 cyclones—were found to be related to higher values of relative vorticity due to baroclinic forcing (q_{co}) than to diabatic processes. Most systems have positive values of balanced relative vorticity due to both baroclinic forcing and diabatic processes; however, negative values of vorticity in Fig. 10 are not automatically be interpreted as cyclolytic contributions to cyclone development. For instance, negative values of balanced relative vorticity in Fig. 10 may be caused by a sharp anticyclonic steering of weak winds, embedded within a broader cyclonic wind field. Therefore, piecewise PV inversion is used in this section to better understand which processes play a primary role in cyclone development, rather than to quantify precisely the contribution of different processes to cyclone wind circulation. In the following sections we present and discuss several exemplary cases of Mediterranean cyclones.

510 5.1 Synergistic forcing from baroclinic and diabatic processes

Figure 11 shows the balanced wind and geopotential fields for two cyclone cases during their mature stage. Both cases have positive contributions to development from both baroclinic and diabatic processes according to Fig. 10. The first cyclone is located in the central Mediterranean and corresponds to a rather deep cyclone with a minimum mean sea level pressure value of 988 hPa (within the top 10% of our cases; Fig. 2). The second cyclone (Fig. 11b) is less deep (994 hPa) and is
515 located to the north of Sicily. Comparing the balanced vorticity fields from the full PV inversion (Figs 11a and 11b) and the cyclone centres from the simulation outputs (cyclones #5 and #78 in supplementary material), it is clear that PV inversion yielded an accurate representation of both systems.

Figures 11c and 11d show that both cases are clearly associated with a PV streamer that intrudes into the Mediterranean from the north, wrapping cyclonically around the cyclone centres. Figure 11c shows that the cyclone centre is close to the local
520 minimum of geopotential, as imposed by q_{co} anomalies at 850 hPa, while the centre of the second cyclone in Fig. 11d is embedded within a broader and rather weaker cyclonic circulation. Consistent with the imposed circulation by q_{co} , both cyclone centres are also collocated with the local minimum of the geopotential field imposed by the non-conserved PV tracers at 850 hPa (Figs 11e and 11f). As a result, baroclinic and diabatic forcings synergistically contribute to the
525 development of both cyclones. Such a scenario is expected to be typical for most intense cyclones, the majority of which are concentrated within the range of positive relative vorticity values in Fig 10. Finally, it is noteworthy that Fig. 11e shows a filament of high PV values over the Aegean Sea with a south-north direction, and another one that extends southwards from the cyclone centre. Both filaments seem to outline the frontal structure of the cyclone and are coincident with the high relative vorticity values in both Fig. 11a and the supplementary material (cyclone #5).

530 5.2 Diabatically driven cyclones

Figure 12 presents two cases where diabatic processes contribute strongly to cyclone development. Both cyclones are located within the central Mediterranean and present a clear low-pressure centre. Especially, the cyclone in Fig. 12a is among the deepest ones in our dataset (982 hPa). In contrast, the cyclone in Fig. 12b is one of the weakest ones with a central mean sea level pressure of 1010 hPa. In both cases, q_{co} presents a weak PV anomaly in the upper troposphere with a rather irregular
535 shape (Figs 12c and 12d). In the first case, this anomaly imposes a rather weak balanced wind field at 850 hPa (Fig. 12c), while in the second case the imposed cyclonic circulation is clearly displaced from the cyclone's centre (Fig. 12d).

In contrast to the conserved component of PV, the non-conserved PV in Figs 12e and 12f presents high PV values that are concentrated in the cyclone centres. The inversion of such high values results in balanced geopotential fields with strong
540 gradients, both presenting local minima over the cyclone centres (Figs 12e and 12f). To further clarify the role of convection in these two cases, Fig. 13 shows the fields of $q_{mp}+q_{cu}+q_{bl}+q_{lw}$ at 850 hPa, as well as their imposed balanced geopotential and

wind fields. In both cases, the PV tracers are fairly similar to Figs 12e and 12f, while the imposed balanced fields clearly show a cyclonic circulation around the cyclone centres. Differences between Fig. 13 and Figs 12e and 12f are majorly attributed to q_{mo} (q_{sw} is negligible); however, these differences are not changing the primary role of convection in forming the cyclonic circulation. Therefore, Figs 12 and 13 present two cases where baroclinic instability plays a secondary role to their development and thus they can be considered diabatically driven cyclones.

5.3 Mountain forcing of cyclones

Figure 14 shows two examples that correspond to Alpine cyclogenesis. The relative vorticity field at 850 hPa presents high values that overlap with the western side of the Alpine mountain chain (Fig. 1) in both the actual model output and the balanced fields in Figs 14a and 14b. Cyclogenesis in the cases of Fig. 14—as for the majority of Mediterranean cyclones—is related to an intruding PV streamer. As shown in Figs 14c and 14d, during the mature stage of the cyclones, the streamer is located over the islands of Corsica and Sardinia imposing a cyclonic circulation over the region. In the first case (Fig. 14a), the cyclonic circulation associated with the full PV distribution is centred over the Gulf of Genoa and extends all over the Mediterranean region. In contrast, the streamer in the second case is located on the east side of a low-level ridge (Fig. 14b) and therefore its imposed cyclonic circulation is displaced towards the east, over the Adriatic Sea (Fig. 14d). Despite the differences between the two cases, both PV streamers impose strong northerly winds that pass over the western side of the Alps. These winds are expected to increase the horizontal gradient of wind speed between the western and eastern side of the Alps and thus to reinforce relative vorticity.

In both cases, PV banners clearly stand out as filaments of high PV values that enter into the cyclone centre from the western flank of the Alps (Figs 14e and 14f). Piecewise inversion of non-conserved PV tracers shows that these filaments impose a cyclonic circulation that is centred over the Alps and reinforces the cyclones (Figs 14e and 14f). To better understand the processes leading to these filaments, Fig. 15 further decomposes the non-conserved PV tracers at 850 hPa into $q_{mp}+q_{cu}+q_{lw}+q_{bl}$ and q_{mo} . In both cyclones the high PV values are clearly related to high values of q_{mo} , while $q_{mp}+q_{cu}+q_{lw}+q_{bl}$ seems to play a secondary role among the non-conserved PV tracers. It is, however, important to stress that these cyclones would be unlikely to form without the upper-tropospheric PV streamers. ~~After all, the PV banners are formed due to the interaction of a northerly wind with the Alps. On the other hand, it is possible that PV banners play a crucial role in the development of the cyclone. This result is consistent with the sensitivity tests of McTaggart-Cowan et al., et al. (2010b) who showed that the removal of the Alps inhibited cyclogenesis of a medecane. In fact, baroclinic forcing and topographic PV production contribute both to lee cyclogenesis. However, disentangling their relative contributions is a rather challenging issue since both processes take place in parallel (Buzzi et al. 2020 and references therein). Nevertheless, PV tracers may provide further insights into this complex problem. As discussed in section 4.1, mountains are likely to play the role of a "constant" PV source due to their interaction with the air flow. This constant PV source is imprinted into the simulations' initial conditions and corresponds to the low-level streamer of q_{co} that wraps cyclonically around the cyclone centres in Fig.~~

575 [5b. However, Fig. 15 shows that large part of high PV values close to mountains is related to \$q_{mo}\$ and thus to Ekman and baroclinic PV production \(last two terms in Eq. 8\). In these regards, PV production plays an important role in lee cyclogenesis. This is consistent with the sensitivity tests of McTaggart-Cowan et al. \(2010b\) who showed that the removal of the Alps inhibited cyclogenesis of a medicane.](#)

580 [After the stage of cyclogenesis, Mediterranean cyclones are still influenced by mountains interactions with the air flow due to the complex geography of the region and its sharp land-sea transitions. For instance, an air mass might reinforce a cyclone when it reaches to its centre due to experiencing earlier PV production while interacting with topography. However, the time period during which air masses sustain their high values of \$q_{mo}\$ is rather uncertain, i.e. the "dynamical memory" of \$q_{mo}\$, similar to that discussed for latent heating in section 4.2. Therefore, it is rather difficult to discern the areas where the](#)
585 [influence of mountains is more important. Such analysis would benefit from a Lagrangian approach \(e.g. Attinger et al. 2019\) where the evolution and origin of high values of \$q_{mo}\$ could be backtracked to mountains. However, this is currently out of the scope of this study and will be the subject of a following research study.](#)

5.4 Two peculiar cases of Mediterranean cyclones

In this section, we present two rather peculiar cases to illustrate the variability of the implicated dynamics in cyclone
590 development.

The first case is presented in Fig. 16a where the cyclone centre is nested within an elongated, meridional structure of high values of relative vorticity at 850 hPa. This elongated structure is fairly consistent with the direct model outputs (cyclone #24 in the supplementary material). However, in the model outputs there is also a secondary local maximum of relative vorticity
595 south of the actual centre. The balanced fields in Fig. 16a instead capture this secondary centre as a southward extension of high values of relative vorticity. Such dual-centred cyclones have been found in several locations in the extratropics (Hanley and Caballero, 2012). Figure 16**cb** shows that the primary cyclone centre (depicted by the magenta dot) is collocated with the local minimum of the imposed geopotential field by q_{co} . Therefore, this cyclone is clearly intensified by baroclinic instability. Interestingly, the geopotential field imposed by the non-conserved PV also presents a local minimum, but this
600 minimum is coincident with the secondary centre of the cyclone located to the south of the primary centre. Consequently, a dual centre of a cyclone may be formed due to different processes and a cyclone could be characterized as a diabatically or baroclinically driven system depending on the choice of its representative centre. A case of a Mediterranean cyclone with a dual centre has been recently shown by Carrio [et al., et al.](#) (2020). In their case study analysis, the authors showed that a secondary cyclone centre was nested within a main, primary cyclone. The latter corresponded to baroclinic low pressure
605 cyclone, related to a clear frontal structure, while the secondary cyclone centre was related to a local pressure minimum generated by convection.

The second cyclone case (Fig. 16b) corresponds to one of the few cyclones in our dataset that presents exceptionally high values of q_{mo} at 850 hPa. This rather deep cyclone (990 hPa) is located to the southwest side of Sardinia and is related to a rather "irregular" PV streamer, when compared to the previous exemplary cases (Fig. 16d). The streamer presents an east-west direction along 37°N, enters the region from the Atlantic Ocean and has lower values of PV than the other examples shown. The geopotential field imposed by q_{co} at 850 hPa is characterized by an anticyclonic circulation centred over Italy and a cyclonic circulation centred over North Africa. The cyclone centre in Fig. 16b is located between these two opposite circulations and, therefore, q_{co} does not contribute significantly to the cyclonic circulation, at least during the cyclone mature stage. On the other hand, the sum of all non-conserved PV tracers in Fig. 16f shows rather high values that are concentrated around the cyclone centre. This leads to a strong geopotential gradient in Fig. 16f centred over the cyclone centre. The q_{mo} term represents about 80% of these high positive values (not shown) and therefore momentum forcing has a leading role in the development of the cyclone.

In our knowledge, this is the first study that shows a cyclone system where the mature stage is self-sustained by momentum forcing within the PBL. It is however noteworthy that the convection parametrisation that we use in this study lacks application of momentum adjustments to the wind field. It is thus possible that part of the high q_{mo} values in Fig. 16f is related to the response of PBL parametrisation to convection. To test this hypothesis we performed an additional simulation where the Kain-Fritsch convective parameterization (Kain 2004) was replaced by the one of Tiedke (Zhang and Wang, 2017). The latter provides momentum adjustments, introduced to the model as an additional momentum forcing to the model primitive equations. Therefore, we introduced an additional PV tracer to Eq. 5. This new PV tracer is calculated similarly to q_{mo} , whereas F in Eq. 4 represents the momentum adjustments by the Tiedke convection parametrisation. Results from this new simulation are highly consistent with the original simulation and q_{mo} still presents exceptionally high PV values. In fact, the new PV tracer is of the order of 0.2 PVU in absolute values close to the cyclone centre (not shown).

This case study that we present in Fig. 16f is of particular interest for gaining further insights into the ambiguous role of q_{mo} in acting as both a source and sink of PV in cyclone centres. For this reason, we further decomposed the PV budget by introducing two additional PV tracers. These new PV tracers combine the three terms of Eq. 8: the first PV tracer accumulates the Ekman PV generation due to the first term of Eq. 8 (hereafter q_{mo1}) and the second PV tracer accumulates the baroclinic PV generation due to the second and third term of Eq. 8 (hereafter q_{mo2}). Figure 17a shows the potential temperature at 850 hPa, along with the fields of wind, q_{mo} , and q_{mo1} at the time of the cyclone's mature stage. The cyclone has a clear frontal structure where high values of q_{mo} are concentrated along the warm front. This configuration is fairly similar to the idealized simulation results of Adamson et al. (2006). In contrast to q_{mo} , q_{mo1} is acting cyclolytically, by imposing negative PV values around the cyclone centre. Therefore, it is clear that Ekman PV generation is weakening the cyclonic circulation and thus the high PV values of q_{mo} close to the cyclone centre are by default related to q_{mo2} . To better understand this PV source, Fig. 17b shows the vertical cross section of F_x along the green dashed line of Fig. 17a. In addition, Fig. 17b

645 shows the instantaneous baroclinic PV generation, i.e. the sum of the second and third terms of Eq. 8. PV sources are located within the PBL, in the eastern side of the cyclone (black contours in Fig. 17b) where the vertical gradient of F_x is negative. Although baroclinic PV production by momentum forcing within the PBL is expected to be common in extratropical cyclones, the conditions that lead q_{mo} to dominate dynamics of a cyclone are yet to be determined. This will be the object of a forthcoming study.-

5.5 Medicanes from a PV-based perspective

650 Medicanes have been long been proposed to share similar dynamics with tropical cyclones (e.g Emanuel, 2005; Fita et al. 2007). Nevertheless, a PV streamer or a cut-off system is commonly present close to medicanes and therefore, baroclinic forcing is expected to contribute to the development of these systems (Tous and Romero 2013; Flaounas et al., et al. 2015; Nastos et al., et al. 2018). This raises the question of whether medicanes can be considered as "purely" diabatic systems, driven by latent heat release as in their tropical counterparts. The ten cases included in this study have been diagnosed as medicanes due to their spiral cloud coverage and cloudless "eye" (Nastos et al., et al. 2018; Di Muzio et al., et al. 2019). The application of piecewise PV inversion to the conserved and non-conserved PV components is a sufficient method to identify 655 whether these medicanes indeed share similar dynamics with tropical cyclones, or at least are "mostly" driven by diabatic processes.

Figure 10 places most medicanes (depicted by red circles) within the range of positive contributions from both baroclinic and diabatic forcings. However, two medicanes have close to a zero contribution from q_{co} , including medicane Trixie (Di Muzio et al., et al. 2019) which corresponds to the case presented in Fig. 12b. In addition, a single medicane case is shown by Fig. 660 10 to be an outlier with considerable contributions to its relative vorticity from both baroclinic and diabatic processes: 5.7 and $2.5 \times 10^{-5} \text{ s}^{-1}$, respectively. The other medicanes in Fig. 10 are similar to the mixed cases presented in section 5.4.1 and thus the involvement of baroclinic forcing through the PV streamer cannot be neglected. This finding agrees with previous studies where factor separation techniques showed that the PV streamer plays a primary role in the development of 665 medicanes, especially when interacting with diabatic processes (Fita et al., et al. 2006; Carrio et al., et al. 2017).

In this study, we focus on the mature stage of cyclones, i.e when cyclones attain their maximum intensity. In earlier stages of their lifecycles, the forcing from the PV streamer will only be more important since baroclinic instability is the main cyclogenesis processes. Nevertheless, the contribution of high PV values in the lower troposphere to cyclone development 670 during the cyclogenesis stage cannot be neglected. Indeed, the cyclogenesis of several medicane cases—along with other cases—coincided with, or came after, major convective events (Miglietta et al., et al. 2013; Flaounas et al., et al. 2015; Fita and Flaounas, 2018). As discussed in section 4, the memory effect of PV tracers may be important for the intensification of cyclones even after the peak of convective activity.

675 While a cyclone develops, increasing values of diabatically produced PV in its centre may eventually lead to tropical transition, i.e. when a cyclone initially formed as a baroclinic low is eventually only sustained by latent heat release (Davis and Bosart, 2004). Our results show that this transition cannot be excluded in the Mediterranean; however, this seems to be a rather rare case scenario, especially due to the short lifespan of cyclones and the sharp land-sea transitions. Such cases would be outliers and found towards the top left of the scatterplot shown in Fig. 10. However, Fig. 10 shows that, for the
680 cases considered here, these outliers were not identified as medicanes, probably due to not complying with the phenomenological criterion of a cloudless "eye" in their centre. Therefore, a new definition of medicanes that resides in atmospheric dynamics should be established as systems that are sustained, or strongly driven, by diabatic processes.

6. Summary and conclusions

From the PV-thinking perspective, a cyclone may be conceptualized as the amalgamation of different PV anomalies. In this
685 framework, we implemented into WRF, and applied, an online diagnostic to 100 simulations of the most intense systems that occurred during a 10-year period, from 2008 to 2017. This diagnostic decomposed PV into different components, each related to a parametrized physical process, and enabled piecewise PV inversion to provide further insights into the role of different processes in cyclone development.

690 Our first objective aimed at quantifying the relative contribution of different processes to the PV-structure of Mediterranean cyclones. The vertical PV structure of the 100 cyclones showed two main contributions: one in the upper troposphere, related to the intrusion of the PV streamer into the Mediterranean, and one in the lower troposphere, related to the net production of PV due to diabatic processes. While the upper-tropospheric anomaly is almost solely composed of the conserved PV tracer (q_{co}), the one in the lower troposphere is the outcome of the sum of several contrasting, non-conserved PV tracers. More
695 precisely, latent heat release is shown to correspond to a net source of PV from ground level until the mid troposphere, approximately at 500 hPa. This net PV production was shown to be partly counteracted by sink processes related to longwave radiation and turbulent diffusion of temperature. Consequently, $q_{mp}+q_{cu}$ and $q_{lw}+q_{bi}$ present two vertical profiles almost symmetric about zero. Eventually, their sum corresponds to a positive PV anomaly at around 850 hPa, i.e. the level where cyclones typically have their maximum intensity in terms of relative vorticity. Momentum mixing and frictional
700 forces were shown to have an ambiguous effect on cyclone intensity. Indeed, q_{mo} was shown to present both negative and positive signs in the lower troposphere though its average contribution over all 100 cyclones was shown to be positive. Interestingly, several cases presented exceptionally high q_{mo} values, comparable to the PV produced due to latent heat release. Finally, our results showed that q_{co} in the lower atmospheric levels may attain high values close to the mountains. Such values pre-exist cyclogenesis and are expected to enhance the amplitude of the local maximum of PV in cyclone
705 centres at 850 hPa, but also to balance negative PV values in the first model levels due to $q_{lw}+q_{bi}$.

Our second objective aimed at analysing cyclone development with regards to their baroclinic and diabatic forcing. In fact, Fig. 9 showed that there is hardly a straightforward relationship between the positive low-level PV anomalies due to latent heat release ($q_{mp}+q_{cu}$) and cyclone intensity. Consequently, diabatic processes alone do not function as an index of cyclone intensity. Indeed, piecewise PV inversion showed that most cyclones developed through the contribution of both baroclinic and diabatic forcings. In fact, Fig. 10 showed that the two forcings tend to have a negative, linear relationship: the more a cyclone is diabatically driven, the less baroclinic instability contributes to its mature stage. Piecewise PV inversion is a method that incorporates uncertainties due to technical challenges in its application. Therefore, our results distinguish the processes that play primary roles in cyclone development, rather than quantify their precise contribution. In this context, several exemplary cases of Mediterranean cyclones were analysed using inverted conserved and non-conserved PV tracer fields. In most cases, the contribution of the upper PV streamer was shown to be important for the surface cyclone. In fact, cyclone centres were typically located in the centre, or within, the low-level cyclonic circulation imposed by q_{co} . A few cases were shown to be mainly driven by diabatic processes. In these cases, the PV streamer was still present, but imposed a rather weak low-level cyclonic circulation. Other exemplary cases of Alpine cyclogenesis along with the case presented in Fig. 16b showed that the role of q_{mo} is not negligible and may even play a primary role in sustaining the cyclonic circulation. Finally, we focused on medicanes and, in particular, on the processes that drive their development. Results showed that most of the known medicanes cases developed due to the forcing of both diabatic and baroclinic processes, although some cases like Trixie were shown to have developed under the influence of a strong diabatic forcing.

This study sheds some light on the variable contribution of different atmospheric processes to the development of Mediterranean cyclones. In this regard, PV, and more precisely PV tracers, is a perfectly adequate method to translate processes into cyclone dynamics. However, PV is a complex variable and several methodological challenges are met when it is used for such analyses. Consequently, future work could use improved implementations of piecewise PV inversion that are more consistent with the primitive equations of the model and adequate more suitable for fine scale grids (e.g. Decker, 2010). In addition, other case studies that are mainly driven by q_{mo} should be investigated. It would come as a surprise if such cases are only found in the Mediterranean region. Therefore, an analysis using PV tracers should be also applied to a wide variety of other cyclones within the storm tracks. Such an effort should include the application of piecewise PV inversion to a much wider range of cyclones than the 100 cases used in this study. This will enhance our insights into the process-based continuum of cyclones, as shown in Fig. 10, and would likely reveal a wide number of diabatically driven cyclones. While focusing on the most intense cyclones is a reasonable approach to understand the main intensification processes of Mediterranean cyclones less strong cyclones are also relevant to high impact weather, especially concerning the occurrence of heavy rainfall (Flaounas et al. 2018). In subsequent studies, we thus aim to extend our analysis to more cyclones to understand their driving processes but this time from the perspective of impacts. Finally, it is in the perspectives of this study to provide a new definition of medicanes, based solely on atmospheric dynamics: Mediterranean cyclones that are sustained,

740 or strongly driven, by diabatic processes. This definition would allow to the community to expand its focus to a broader
group of cyclones than the limited number of cases, currently qualified as medicanes.

Code availability: The PV-tracers diagnostic corresponds to a single module, written in Fortran and its implementation
requires additional modifications to several parts of the original WRF code. The whole PV-tracers package is adapted to
745 WRF version 4.0 and might demand minor, additional modifications for its implementation in subsequent versions of the
model. The diagnostic and all relevant source code modifications are freely available upon request.

Acknowledgements: We are grateful to Baruch Ziv and to two anonymour Reviewers for their constructive comments and
suggestions that greatly improved this article. EF is thankful to Oscar Martinez-Alvarado for his help and support ~~in~~while
750 developing the PV budget diagnostic for WRF and to John Methven for ~~the~~ interesting discussions on the interpretation of
PV tracers.

Financial support EF acknowledges funding through the H2020 European Research Council (INTEXseas (grant no.
787652)). FT acknowledges funding through the German Research Foundation Grant RI 1771/4-1 and the Transregional
755 Collaborative Research Center SFB / TRR 165 “Waves to Weather” (www.wavestoweather.de) funded by the German
Research Foundation (DFG).

References

Adamson, D. S., Belcher, S. E., Hoskins, B. J. and Plant, R. S.: Boundary-layer friction in midlatitude cyclones, Q. J. R.
Meteorol. Soc., 132(614), 101–124, doi:10.1256/qj.04.145, 2006.

760

Aebischer, U. and Schär, C.: Low-Level Potential Vorticity and Cyclogenesis to the Lee of the Alps, J. Atmos. Sci., 55,
186–207, doi:10.1175/1520-0469(1998)055<0186:LLPVAC>2.0.CO;2, 1998

Ahmadi-Givi, F., Graig, G. C. and Plant, R. S.: The dynamics of a midlatitude cyclone with very strong latent-heat release,
765 Quarterly Journal of the Royal Meteorological Society, 130(596), 295–323, <https://doi.org/https://doi.org/10.1256/qj.02.226>,
2004.

Alpert, P., Neeman, B. U. and Shay-El, Y.: Climatological analysis of Mediterranean cyclones using ECMWF data, Tellus
A, 42(1), 65–77, doi:10.3402/tellusa.v42i1.11860, 1990.

770

Attinger, R., Spreitzer, E., Boettcher, M., Forbes, R., Wernli, H. and Joos, H.: Quantifying the role of individual diabatic processes for the formation of PV anomalies in a North Pacific cyclone, *Q.J.R. Meteorol. Soc.*, 145(723), 2454–2476, doi:10.1002/qj.3573, 2019.

775 [Bleck, R. and Mattocks, C.: A preliminary analysis of the role of potential vorticity in Alpine lee cyclogenesis, *Beiträge zur Physik der Atmosphäre*, 57\(3\), 357-368, 1984.](#)

[Boettcher, M. and Wernli, H.: Life Cycle Study of a Diabatic Rossby Wave as a Precursor to Rapid Cyclogenesis in the North Atlantic—Dynamics and Forecast Performance, *Monthly Weather Review*, 139\(6\), 1861–1878, <https://doi.org/10.1175/2011MWR3504.1>, 2011.](#)

780

Bracegirdle, T. J. and Gray, S. L.: The dynamics of a polar low assessed using potential vorticity inversion, *Q.J.R. Meteorol. Soc.*, 135(641), 880–893, doi:10.1002/qj.411, 2009.

785 Büeler, D. and Pfahl, S.: Potential Vorticity Diagnostics to Quantify Effects of Latent Heating in Extratropical Cyclones. Part I: Methodology, *J. Atmos. Sci.*, 74(11), 3567–3590, doi:10.1175/JAS-D-17-0041.1, 2017.

[Buzzi, A., Davolio, S. and Fantini, M.: Cyclogenesis in the lee of the Alps: a review of theories, *Bull. of Atmos. Sci.& Technol.*, <https://doi.org/10.1007/s42865-020-00021-6>, 2020.](#)

790

Čampa, J. and Wernli, H.: A PV Perspective on the Vertical Structure of Mature Midlatitude Cyclones in the Northern Hemisphere, *J. Atmos. Sci.*, 69(2), 725–740, doi:10.1175/JAS-D-11-050.1, 2012.

Campins, J., Genovés, A., Picornell, M. A. and Jansà, A.: Climatology of Mediterranean cyclones using the ERA-40 dataset, *Int. J. Climatol.*, 31(11), 1596–1614, doi:10.1002/joc.2183, 2011.

795

Cao, J. and Xu, Q.: Computing Hydrostatic Potential Vorticity in Terrain-Following Coordinates, *Mon. Wea. Rev.*, 139(9), 2955–2961, doi:10.1175/MWR-D-11-00083.1, 2011.

800 Carrió, D. S., Homar, V., Jansa, A., Romero, R. and Picornell, M. A.: Tropicalization process of the 7 November 2014 Mediterranean cyclone: Numerical sensitivity study, *Atmos. Res.*, 197, 300–312, doi:10.1016/j.atmosres.2017.07.018, 2017.

Carrió, D. S., Homar, V., Jansà, A., Picornell, M. A. and Campins, J.: Diagnosis of a high-impact secondary cyclone during HyMeX-SOP1 IOP18, *Atmos. Res.*, 242, 104983, doi:10.1016/j.atmosres.2020.104983, 2020.

805

Cavicchia, L., von Storch, H. and Gualdi, S.: Mediterranean Tropical-Like Cyclones in Present and Future Climate, *J. Climate*, 27(19), 7493–7501, doi:10.1175/JCLI-D-14-00339.1, 2014.

810 Chaboureau, J.-P., Pantillon, F., Lambert, D., Richard, E. and Claud, C.: Tropical transition of a Mediterranean storm by jet crossing, *Q.J.R. Meteorol. Soc.*, 138(664), 596–611, doi:10.1002/qj.960, 2012.

Chagnon, J. M., Gray, S. L. and Methven, J.: Diabatic processes modifying potential vorticity in a North Atlantic cyclone, *Q.J.R. Meteorol. Soc.*, 139(674), 1270–1282, doi:10.1002/qj.2037, 2013.

815 [Chagnon, J. M. and Gray, S. L.: A Diabatically Generated Potential Vorticity Structure near the Extratropical Tropopause in Three Simulated Extratropical Cyclones, *Monthly Weather Review*, 143\(6\), 2337–2347, https://doi.org/10.1175/MWR-D-14-00092.1, 2015.](https://doi.org/10.1175/MWR-D-14-00092.1)

820 Charney, J.: The Use of the Primitive Equations of Motion in Numerical Prediction, *Tellus*, 7(1), 22–26, doi:10.1111/j.2153-3490.1955.tb01138.x, 1955.

[Chan, J. C.-L., Ko, F. M. F., and Lei, Y. M.: Relationship between potential vorticity tendency and tropical cyclone motion, *J. Atmos. Sci.*, 59, 1317–1336, 2002.](https://doi.org/10.1175/JAS4113)

825 Claud, C., Alhammoud, B., Funatsu, B. M. and Chaboureau, J.-P.: Mediterranean hurricanes: large-scale environment and convective and precipitating areas from satellite microwave observations, *Nat. Hazards Earth Syst. Sci.*, 10(10), 2199–2213, doi:10.5194/nhess-10-2199-2010, 2010.

830 Dafis, S., Rysman, J.-F., Claud, C. and Flaounas, E.: Remote sensing of deep convection within a tropical-like cyclone over the Mediterranean Sea, *Atmos. Sci. Lett.*, 19(6), e823, doi:10.1002/asl.823, 2018.

Davis, C. A. and Emanuel, K. A.: Potential vorticity diagnostics of cyclogenesis, *Mon. Wea. Rev.*, 119(8), 1929–1953, 1991.

Davis, C.A.: Piecewise potential vorticity inversion. *J. Atmos. Sci.*, 49, 1397–1411, 1992.

835

Davis, C.A. and Bosart, L.F.: The TT problem: Forecasting the tropical transition of cyclones. *B. Am. Meteorol. Soc.*, 85(11): 1657., 2004.

Decker, S. G.: Nonlinear Balance in Terrain-Following Coordinates, *Mon. Wea. Rev.*, 138(2), 605–624, 2010.

840

[Deveson, A. C. L., Browning, K. A. and Hewson, T. D.: A classification of FASTEX cyclones using a height-attributable quasi-geostrophic vertical-motion diagnostic, *Q. J. R. Meteorol. Soc.*, 128\(579\), 93–117, <https://doi.org/10.1256/00359000260498806>, 2002.](https://doi.org/10.1256/00359000260498806)

845 Di Muzio, E., Riemer, M., Fink, A. H. and Maier-Gerber, M.: Assessing the predictability of Medicanes in ECMWF ensemble forecasts using an object-based approach, *Q.J.R. Meteorol. Soc.*, 145(720), 1202–1217, doi:10.1002/qj.3489, 2019.

Dudhia, J.: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model. *J. Atmos. Sci.*, 46:3077–3107, 1989.

850

Emanuel, K.: Genesis and maintenance of "Mediterranean hurricanes", *Adv. Geosci.*, 2, 217–220, doi:10.5194/adgeo-2-217-2005, 2005.

855 Epifanio, C.C. and Durran, D.R.: Lee-vortex formation in free-slip stratified flow over ridges. Part II: Mechanisms of vorticity and PV production in nonlinear viscous wakes. *J. Atmos. Sci.*, 59, 1166–1181, 2002.

Fita, L. and Flaounas, E.: Medicanes as subtropical cyclones: the December 2005 case from the perspective of surface pressure tendency diagnostics and atmospheric water budget, *Q.J.R. Meteorol. Soc.*, 144(713), 1028–1044, doi:10.1002/qj.3273, 2018.

860

Fita, L., Romero, R., Luque, A., Emanuel, K. and Ramis, C.: Analysis of the environments of seven Mediterranean tropical-like storms using an axisymmetric, nonhydrostatic, cloud resolving model, *Nat. Hazards Earth Syst. Sci.*, 7(1), 41–56, doi:10.5194/nhess-7-41-2007, 2007.

865 Fita, Ll., Romero, R. and Ramis, C.: Intercomparison of intense cyclogenesis events over the Mediterranean basin based on baroclinic and diabatic influences, *Adv. Geosci.*, 7, 333–342, doi:10.5194/adgeo-7-333-2006, 2006.

870 Flamant, C., Richard, E., Schär, C., Rotunno, R., Nance, L., Sprenger, M. and Benoit, R.: The wake south of the Alps: Dynamics and structure of the lee-side flow and secondary potential vorticity banners, *Q.J.R. Meteorol. Soc.*, 130(599), 1275–1303, doi:10.1256/qj.03.17, 2004.

Flaounas, E., Kotroni, V., Lagouvardos, K. and Flaounas, I.: CycloTRACK (v1.0) – tracking winter extratropical cyclones based on relative vorticity: sensitivity to data filtering and other relevant parameters, *Geosci. Model Dev.*, 7(4), 1841–1853, doi:10.5194/gmd-7-1841-2014, 2014.

875

Flaounas, E., Raveh-Rubin, S., Wernli, H., Drobinski, P. and Bastin, S.: The dynamical structure of intense Mediterranean cyclones, *Clim. Dyn.*, 44(9–10), 2411–2427, doi:10.1007/s00382-014-2330-2, 2015.

880 Flaounas, E., Kelemen, F. D., Wernli, H., Gaertner, M. A., Reale, M., Sanchez-Gomez, E., Lionello, P., Calmanti, S., Podrascanin, Z., Somot, S., Akhtar, N., Romera, R. and Conte, D.: Assessment of an ensemble of ocean–atmosphere coupled and uncoupled regional climate models to reproduce the climatology of Mediterranean cyclones, *Clim. Dyn.*, 51(3), 1023–1040, doi:10.1007/s00382-016-3398-7, 2018a.

885 Flaounas, E., Kotroni, V., Lagouvardos, K., Gray, S. L., Rysman, J.-F. and Claud, C.: Heavy rainfall in Mediterranean cyclones. Part I: contribution of deep convection and warm conveyor belt, *Clim. Dyn.*, 50(7–8), 2935–2949, doi:10.1007/s00382-017-3783-x, 2018b.

890 Flaounas, E., Fita, L., Lagouvardos, K. and Kotroni, V.: Heavy rainfall in Mediterranean cyclones, Part II: Water budget, precipitation efficiency and remote water sources, *Clim. Dyn.*, 53(5–6), 2539–2555, doi:10.1007/s00382-019-04639-x, 2019.

Galanaki, E., Flaounas, E., Kotroni, V., Lagouvardos, K. and Argiriou, A.: Lightning activity in the Mediterranean: quantification of cyclones contribution and relation to their intensity: Mediterranean cyclones associated with lightning activity, *Atmos. Sci. Lett.*, 17(9), 510–516, doi:10.1002/asl.685, 2016.

895 Gray, S. L.: Mechanisms of midlatitude cross-tropopause transport using a potential vorticity budget approach, *J. Geophys. Res.*, 111(D17), D17113, doi:10.1029/2005JD006259, 2006.

[Gray, S. L. and Dacre, H. F.: Classifying dynamical forcing mechanisms using a climatology of extratropical cyclones, Q. J. R. Meteorol. Soc., 132\(617\), 1119–1137, https://doi.org/10.1256/qj.05.69, 2006.](https://doi.org/10.1256/qj.05.69)

900

Hanley, J. and Caballero, R.: Objective identification and tracking of multicentre cyclones in the ERA-Interim reanalysis dataset, *Q.J.R. Meteorol. Soc.*, 138(664), 612–625, doi:10.1002/qj.948, 2012.

905 Hong, S.Y., Juang H.M.H. and Zhao, Q.: Implementation of prognostic cloud scheme for a regional spectral model, *Mon. Wea. Rev.*, 126: 26212639, 1998.

Hong, S.-Y.: A new stable boundary-layer mixing scheme and its impact on the simulated East Asian summer monsoon, Q.J.R. Meteorol. Soc., 136(651), 1481–1496, doi:10.1002/qj.665, 2010.

910 Hong, S.-Y., Noh, Y. and Dudhia, J.: A New Vertical Diffusion Package with an Explicit Treatment of Entrainment Processes, Mon. Wea. Rev., 134(9), 2318–2341, doi:10.1175/MWR3199.1, 2006a.

Hong, S.-Y., Noh, Y. and Dudhia, J.: A New Vertical Diffusion Package with an Explicit Treatment of Entrainment Processes, Mon. Wea. Rev., 134(9), 2318–2341, doi:10.1175/MWR3199.1, 2006b.

915

Huo, Z., Zhang, D. L. and Gyakum, J. R.: Interaction of potential vorticity anomalies in extratropical cyclogenesis. Part I: Static piecewise inversion. Mon. Wea. Rev., 127(11), 2546-2562, 1999.

Kain, J.S.: The Kain–Fritsch convective parameterization: an update, J. Appl. Meteorol., 43(1):170-181, 2004.

920

Kouroutzoglou, J., Flocas, H. A., Keay, K., Simmonds, I. and Hatzaki, M.: Climatological aspects of explosive cyclones in the Mediterranean, Int. J. Climatol., 31(12), 1785–1802, doi:10.1002/joc.2203, 2011.

Lackmann, G. M.: Cold-frontal potential vorticity maxima, the low-level jet, and moisture transport in extratropical cyclones, Mon. Weather Rev., 130, 59–74, [https://doi.org/10.1175/1520-0493\(2002\)130<0059:cfpvmt>2.0.co;2](https://doi.org/10.1175/1520-0493(2002)130<0059:cfpvmt>2.0.co;2), 2002.

925

Lagouvardos, K., Kotroni, V. and Defer, E.: The 21-22 January 2004 explosive cyclogenesis over the Aegean Sea: Observations and model analysis, Q.J.R. Meteorol. Soc., 133(627), 1519–1531, doi:10.1002/qj.121, 2007.

930 Lamarque, J.-F., and Hess, P. G.: Cross-tropopause mass exchange and potential vorticity budget in a simulated tropopause folding, J. Atmos. Sci., 51, 2246–2269, 1994.

Maheras, P., Flocas, H. A., Patrikas, I. and Anagnostopoulou, Chr.: A 40 year objective climatology of surface cyclones in the Mediterranean region: spatial and temporal distribution, Int. J. Climatol., 21(1), 109–130, doi:10.1002/joc.599, 2001.

935

Martínez-Alvarado, O., Gray, S. L. and Methven, J.: Diabatic Processes and the Evolution of Two Contrasting Summer Extratropical Cyclones, Mon. Wea. Rev., 144(9), 3251–3276, doi:10.1175/MWR-D-15-0395.1, 2016.

McTaggart-Cowan, R., Galarneau, T. J., Bosart, L. F. and Milbrandt, J. A.: Development and Tropical Transition of an
940 Alpine Lee Cyclone. Part I: Case Analysis and Evaluation of Numerical Guidance, *Mon. Wea. Rev.*, 138(6), 2281–2307,
doi:10.1175/2009MWR3147.1, 2010a.

McTaggart-Cowan, R., Galarneau, T. J., Bosart, L. F. and Milbrandt, J. A.: Development and Tropical Transition of an
Alpine Lee Cyclone. Part II: Orographic Influence on the Development Pathway, *Mon. Wea. Rev.*, 138(6), 2308–2326,
945 doi:10.1175/2009MWR3148.1, 2010b.

[Miglietta, M. M., Moscatello, A., Conte, D., Mannarini, G., Lacorata, G. and Rotunno, R.: Numerical analysis of a
Mediterranean ‘hurricane’ over south-eastern Italy: Sensitivity experiments to sea surface temperature, *Atmospheric
Research*, 101\(1–2\), 412–426, <https://doi.org/10.1016/j.atmosres.2011.04.006>, 2011.](#)

950 | [Miglietta, M. M. and Rotunno, R.: Development mechanisms for Mediterranean tropical-like cyclones \(medicanes\), *Q.J.R.
Meteorol. Soc.*, 145\(721\), 1444–1460, doi:10.1002/qj.3503, 2019.](#)

Miglietta, M. M., Laviola, S., Malvaldi, A., Conte, D., Levizzani, V. and Price, C.: Analysis of tropical-like cyclones over
955 the Mediterranean Sea through a combined modeling and satellite approach, *Geophys. Res. Lett.*, 40(10), 2400–2405,
doi:10.1002/grl.50432, 2013.

[Miglietta, M. M., Mastrangelo, D. and Conte, D.: Influence of physics parameterization schemes on the simulation of a
tropical-like cyclone in the Mediterranean Sea, *Atmospheric Research*, 153, 360–375,
960 <https://doi.org/10.1016/j.atmosres.2014.09.008>, 2015.](#)

Miglietta, M. M., Cerrai, D., Laviola, S., Cattani, E. and Levizzani, V.: Potential vorticity patterns in Mediterranean
“hurricanes,” *Geophys. Res. Lett.*, 44(5), 2537–2545, doi:10.1002/2017GL072670, 2017.

965 | [Miglietta, M. M. and Rotunno, R.: Development mechanisms for Mediterranean tropical-like cyclones \(medicanes\), *Q.J.R.
Meteorol. Soc.*, 145\(721\), 1444–1460, doi:10.1002/qj.3503, 2019.](#)

Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J. and Clough, S. A.: Radiative transfer for inhomogeneous
atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, 102(D14), 16663–16682,
970 doi:10.1029/97JD00237, 1997.

Möller, J. D. and Montgomery, M. T.: Tropical cyclone evolution via potential vorticity anomalies in a three-dimensional balance model, *J. Atmos. Sci.*, 57, 3366–3387, 2000.

975 Nastos, P. T., Karavana Papadimou, K. and Matsangouras, I. T.: Mediterranean tropical-like cyclones: Impacts and composite daily means and anomalies of synoptic patterns, *Atmos. Res.*, 208, 156–166, doi:10.1016/j.atmosres.2017.10.023, 2018.

980 Neu, U., Akperov, M. G., Bellenbaum, N., Benestad, R., Blender, R., Caballero, R., Coccozza, A., Dacre, H. F., Feng, Y., Fraedrich, K., Grieger, J., Gulev, S., Hanley, J., Hewson, T., Inatsu, M., Keay, K., Kew, S. F., Kindem, I., Leckebusch, G. C., Liberato, M. L. R., Lionello, P., Mokhov, I. I., Pinto, J. G., Raible, C. C., Reale, M., Rudeva, I., Schuster, M., Simmonds, I., Sinclair, M., Sprenger, M., Tilinina, N. D., Trigo, I. F., Ulbrich, S., Ulbrich, U., Wang, X. L. and Wernli, H.: IMILAST: A community effort to intercompare extratropical cyclone detection and tracking algorithms, *B. Am. Meteorol. Soc.*, 94(4), 529–547, doi:10.1175/BAMS-D-11-00154.1, 2013.

985

Nissen, K. M., Leckebusch, G. C., Pinto, J. G., Renggli, D., Ulbrich, S. and Ulbrich, U.: Cyclones causing wind storms in the Mediterranean: characteristics, trends and links to large-scale patterns, *Nat. Hazards Earth Syst. Sci.*, 10(7), 1379–1391, doi:10.5194/nhess-10-1379-2010, 2010.

990 Pettersen S.: *Weather analysis and forecasting*. McGraw-Hill, New York 428, 1956.

Piaget, N., Froidevaux, P., Giannakaki, P., Gierth, F., Martius, O., Riemer, M., Wolf, G. and Grams, C. M.: Dynamics of a local Alpine flooding event in October 2011: moisture source and large-scale circulation: Dynamics of a Local Alpine Flooding Event, *Q.J.R. Meteorol. Soc.*, 141(690), 1922–1937, doi:10.1002/qj.2496, 2015.

995

Plant, R. S., Craig, G. C. and Gray, S. L.: On a threefold classification of extratropical cyclogenesis, *Q.J.R. Meteorol. Soc.*, 129(594), 2989–3012, <https://doi.org/10.1256/qj.02.174>, 2003.

1000 Prezerakos, N. G., Flocas, H. A. and Brikas, D.: The role of the interaction between polar and subtropical jet in a case of depression rejuvenation over the Eastern Mediterranean, *Meteorol. Atmos. Phys.*, 92(1–2), 139–151, doi:10.1007/s00703-005-0142-y, 2006.

Price, C., Asfur, M. and Yair, Y.: Maximum hurricane intensity preceded by increase in lightning frequency, *Nat. Geosci.*, 2(5), 329–332, doi:10.1038/ngeo477, 2009.

1005

[Pytharoulis, I., Kartsios, S., Tegoulas, I., Feidas, H., Miglietta, M., Matsangouras, I. and Karacostas, T.: Sensitivity of a Mediterranean Tropical-Like Cyclone to Physical Parameterizations, Atmosphere, 9\(11\), 436, <https://doi.org/10.3390/atmos9110436>, 2018a.](https://doi.org/10.3390/atmos9110436)

1010 [Pytharoulis, I.: Analysis of a Mediterranean tropical-like cyclone and its sensitivity to the sea surface temperatures, Atmospheric Research, 208, 167–179, <https://doi.org/10.1016/j.atmosres.2017.08.009>, 2018b.](https://doi.org/10.1016/j.atmosres.2017.08.009)

Raveh-Rubin, S. and Flaounas, E.: A dynamical link between deep Atlantic extratropical cyclones and intense Mediterranean cyclones: Dynamics upstream intense Mediterranean cyclones, Atmos. Sci. Lett, 18(5), 215–221, doi:10.1002/asl.745, 2017.

1015

Raveh-Rubin, S. and Wernli, H.: Large-scale wind and precipitation extremes in the Mediterranean: a climatological analysis for 1979–2012: Mediterranean Large-scale Wind and Precipitation Extremes, Q.J.R. Meteorol. Soc, 141(691), 2404–2417, doi:10.1002/qj.2531, 2015.

1020 Rotunno, R., Grubišić, V. and Smolarkiewicz, P.K.: Vorticity and Potential Vorticity in Mountain Wakes. J. Atmos. Sci., 56, 2796–2810, 1999.

Saffin, L., Methven, J. and Gray, S. L.: The non-conservation of potential vorticity by a dynamical core compared with the effects of parametrized physical processes: Dynamical Core PV, Q.J.R. Meteorol. Soc., 142(696), 1265–1275,

1025 doi:10.1002/qj.2729, 2016.

Schär, C. and Durran D. R.: Vortex Formation and Vortex Shedding in Continuously Stratified Flows past Isolated Topography. J. Atmos. Sci., 54, 534–554, 1997.

1030 Schär, C., Sprenger, M., Lüthi, D., Jiang, Q., Smith, R. B. and Benoit, R.: Structure and dynamics of an Alpine potential-vorticity banner, Q.J.R. Meteorol. Soc., 129(588), 825–855, doi:10.1256/qj.02.47, 2003.

Schlemmer, L., Martius, O., Sprenger, M., Schwierz, C. and Twitchett, A.: Disentangling the Forcing Mechanisms of a Heavy Precipitation Event along the Alpine South Side Using Potential Vorticity Inversion, Mon. Wea. Rev., 138(6), 2336–

1035 2353, doi:10.1175/2009MWR3202.1, 2010.

Schneiderreit, A., Peters, D. H. W., Grams, C. M., Quinting, J. F., Keller, J. H., Wolf, G., Teubler, F., Riemer, M. and Martius, O.: Enhanced tropospheric wave forcing of two anticyclones in the prephase of the January 2009 major stratospheric sudden warming event, Mon. Wea. Rev., 145(5), 1797–1815, doi:10.1175/MWR-D-16-0242.1, 2017.

- 1040 [Schubert, W. H., Hausman, S. A., Garcia, M., Ooyama, K. V., and Kuo, H.-C.: Potential vorticity in a moist atmosphere, J. Atmos. Sci., 58, 3148–3157, 2001](#)
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M., Huang, X. Y., Wang, W., and Powers, J. G.: A description of the advanced research WRF version3, 603NCAR, Tech. Note, Mesoscale and Microscale Meteorology Division, National Center for Atmospheric Research, Boulder, Colorado, USA, 2008
- Seiler, C.: A Climatological Assessment of Intense Extratropical Cyclones from the Potential Vorticity Perspective, J. Climate, 32(8), 2369–2380, doi:10.1175/JCLI-D-18-0461.1, 2019.
- 1050 Spreitzer, E., Attinger, R., Boettcher, M., Forbes, R., Wernli, H. and Joos, H.: Modification of Potential Vorticity near the Tropopause by Nonconservative Processes in the ECMWF Model, J. Atmos. Sci., 76(6), 1709–1726, doi:10.1175/JAS-D-18-0295.1, 2019.
- Stoelinga, M. T.: A potential vorticity-based study of the role of diabatic heating and friction in a numerically simulated baroclinic cyclone. Mon. Wea. Rev., 124(5), 849–874, 1996
- 1055 Teubler, F. and Riemer, M.: Dynamics of Rossby wave packets in a quantitative potential vorticity–potential temperature framework, J. Atmos. Sci., 73(3), 1063–1081, doi:10.1175/JAS-D-15-0162.1, 2016.
- 1060 Teubler, F. and Riemer, M.: Potential-Vorticity Dynamics of Troughs and Ridges within Rossby Wave Packets during a 40-year reanalysis period, WCD, submitted in August 2020
- Tous, M. and Romero, R.: Meteorological environments associated with medicane development, Int. J. Climatol., 33(1), 1–14, doi:10.1002/joc.3428, 2013.
- 1065 Tsidulko, M. and Alpert, P.: Synergism of upper-level potential vorticity and mountains in Genoa lee cyclogenesis - A numerical study, Meteorol. Atmos. Phys., 78(3–4), 261–285, doi:10.1007/s703-001-8178-8, 2001.
- 1070 [Vannière, B., Czaja, A., Dacre, H., Woollings, T. and Parfitt, R.: A potential vorticity signature for the cold sector of winter extratropical cyclones, Q.J.R. Meteorol. Soc., 142\(694\), 432–442, <https://doi.org/10.1002/qj.2662>, 2016.](#)
- Whittaker, I. C., Douma, E., Rodger, C. J. and Marshall, T. J. C. H.: A quantitative examination of lightning as a predictor of peak winds in tropical cyclones, J. Geophys. Res. Atmos., 120(9), 3789–3801, doi:10.1002/2014JD022868, 2015.


1075 Wu, C. C. and Emanuel, K. A.: Potential vorticity diagnostics of hurricane movement. Part 1: A case study of Hurricane Bob (1991). *Mon. Wea. Rev.*, 123(1), 69-92, 1995.

[Zhang, C. and Wang, Y.: Projected Future Changes of Tropical Cyclone Activity over the Western North and South Pacific in a 20-km-Mesh Regional Climate Model, *J. Climate*, 30\(15\), 5923–5941, <https://doi.org/10.1175/JCLI-D-16-0597.1>, 2017.](#)

1080

1085

1090



1095

Figure 1: Simulations domain and orography (colour). Dots depict the locations of the 100 cyclones at their mature stage. The size

of the dots is proportional to the cyclones' central mean sea level pressure (see legend upper left corner of the domain). Dot colours depict a certain season: *black* for winter, *magenta* for spring, *green* for summer and *blue* for autumn.

1100




Figure 2: (a) Monthly distribution of cyclone occurrence. (b) Cyclones sorted according to their central mean sea level pressure during their mature stage.

1105

1110




Figure 3: Composite for all 100 cyclones at the time of their mature stage of PV at 300 hPa (in grey contours with 0.5 PVU contour intervals, starting from 1.5 PVUs), mean sea level pressure (in black contours) and horizontal wind (arrows) and equivalent potential temperature at 850 hPa (colours) with (0,0) denoting the position of cyclone centres.

1115

1120




Figure 4: Vertical profiles of total PV and PV tracers for all 100 cyclones (in *grey lines*), averaged 100 km around cyclone centres at the time of their mature stage. Profile averages and zero PVU are shown in *solid* and *dashed* black lines, respectively. Note that range of values in x-axis is not common for all panels.







Figure 5: (a) Composite averages for all 100 cyclones at the time of their mature stage of q_{co} (in colour) and wind (in arrows) at 300 hPa. (b) as in a, but at 850 hPa. (c) as in (a), but for the sum of all non-conserved PV tracers. (d) as in (b) but for the sum of all non-conserved PV tracers. Note that the value ranges in the colourbars differ among panels. **As a guide to the location of fronts, the composite average of equivalent potential temperature (θ) at 850 hPa is also shown in panels (b) and (d) (black contours, every 2 K, bold for 308 K).**



1135 Figure 6: (a) Average vertical profiles of θ tracers for all 100 simulations, averaged within a 100 km radius around cyclone centres; whiskers length equals twice the standard deviation. (b) as in (a) but for instantaneous temperature tendencies.




1140 **Figure 7: Composite for all 100 cyclones at the time of their mature stage of horizontal wind (arrows) and of different PV tracers at different atmospheric levels (colours), with (0,0) denoting the position of the cyclone centres. As a guide to the location of fronts, the composite average of equivalent potential temperature (θ_e) at 850 hPa is also shown (black contours, every 2 K, bold for 308 K). Note that the range of the colourbar values changes among panels.**




1145

Figure 8: Average vertical profiles of PV tracers for all 100 simulations, averaged within a 100 km radius around the cyclone centres. Whiskers length equals twice the standard deviation.



1150 Figure 9: Scatterplot relating the sum of q_{bl} and q_{lw} to the sum of q_{cu} and q_{mp} for all 100 cyclones. Values of the PV tracers correspond to averages within a 100 km radius around the cyclone centres at the 850 hPa level. The size of the dot is proportional to the mean sea level pressure at the cyclone's centre and *dashed line* represents the $q_{cu} + q_{mp} = -(q_{bl} + q_{lw})$ equation.



1155

Figure 10: Scatterplot relating the balanced relative vorticity fields after inverting the conserved PV tracers (x-axis) to the sum of all non-conserved PV tracers (y-axis). Red circles depict the ten medicane cases in our dataset.

1160

1165

1170

1175





Figure 11: (a) Balanced relative vorticity (*in colour*), wind field (*in arrows*) and geopotential height (*in contour*) at 850 hPa after inverting the total atmospheric PV for cyclone case #5. The model outputs for this case are shown in the supplementary material. 1180 (b) as in (a), but for the case #78. (c) the field of q_{co} at 300 hPa (*in colour*) and the balanced wind (*in arrows*) and geopotential height (*in contour*) fields at 850 hPa after inverting q_{co} . (d) as in (c) but for case #78. (e) the field of the sum of all non-conserved PV tracers at 850 hPa (*in colour*), as well as the balanced wind field (*in arrows*) and geopotential height (*in contour*) at 850 hPa after inverting all non-conserved PV. (f) as in (e), but for case #78. In all panels, the *magenta dot* depicts the cyclones' centre.

1185



1190 Figure 12: As for Fig. 11 but for cases #30 and #75.

1195




Figure 13: (a) The field of $q_{mp}+q_{cu}+q_{bl}+q_{lw}$ at 850 hPa (in colour) and the balanced wind (in arrows) and geopotential height (in contours) fields at 850 hPa after inverting $q_{mp}+q_{cu}+q_{bl}+q_{lw}$ for the cyclone case #30. (b) as in (a), but for case #75.

1200





Figure 14: As for Fig. 11, but for cases #15 and #58.

1205

1210



1215

Figure 15: (a) The field of $q_{mp} + q_{cu} + q_{bl} + q_{lw}$ at 850 hPa (in colour) and the balanced wind (in arrows) and geopotential height (in contours) fields at 850 hPa after inverting $q_{mp} + q_{cu} + q_{bl} + q_{lw}$ for the case #15. (b) as in (a), but for case #58. (c) the field of q_{mo} at 850 hPa (in colour) and the balanced wind (in arrows) and geopotential height (in contours) fields at 850 hPa after inverting q_{mo} , for the case #15. (d) as in (c), but for case #58.





Figure 16: As for Fig. 11, but for cases #24 and #62.



1220

Figure 17: (a) fields of θ (in colour), wind (in arrows), q_{mo} (in black contour with 0.5 PVU contour intervals, starting from 1 PVU) and q_{mo1} (in magenta contour for -0.5 and -1 PVUs) at 850 hPa at the time of the mature stage of cyclone #62. (b) Vertical cross section along the green line of panel (a), showing in colours the field of F_x (momentum forcing from PBL parametrisation) and in black contours the field of baroclinic PV production (sum of two last terms in Eq. 8), starting from 0.05 PVU s^{-1} with an interval of 0.1 PVU s^{-1} .

1225