

~~SST fronts along the~~ The Gulf Stream and Kuroshio SST fronts affect the winter climatology primarily in the absence of cyclones

Leonidas Tsopouridis¹, Thomas Spengler¹, and Clemens Spensberger¹

¹Geophysical Institute, University of Bergen, and Bjerknes Centre for Climate Research, Bergen, Norway

Correspondence: leonidas.tsopouridis@uib.no

Abstract. The Gulf Stream and Kuroshio regions feature strong sea surface temperature (SST) gradients that influence cyclone development and the storm track. ~~Smoothing~~ Previous studies showed that smoothing the SSTs in either the North Atlantic or North Pacific ~~has been shown to yield~~ yields a reduction in cyclone activity, surface heat fluxes, and precipitation, as well as a southward shift of the storm track and the upper-level jet. To what extent these changes are attributable to changes in individual
5 cyclone behaviour, however, remains unclear. Comparing simulations with realistic and smoothed SSTs in the atmospheric general circulation model AFES, we find that the intensification of individual cyclones in the Gulf Stream or Kuroshio region is only marginally affected by reducing the SST gradient. In contrast, we observe considerable changes in the climatological mean state, with a reduced cyclone activity in the North Atlantic and North Pacific storm tracks that are also shifted equator-
10 ward in both basins. The upper-level jet in the Atlantic also shifts equator-ward, while the jet in the Pacific strengthens in its climatological position and extends further east. Surface heat fluxes, specific humidity, and precipitation also respond strongly to the smoothing of the SST, with a considerable decrease of their mean values on the warm side of the SST front. This decrease is more pronounced in the Gulf Stream than in the Kuroshio region, due to the amplified decrease in SST along the Gulf Stream
15 SST front. ~~Subdividing the winter climatology into dates with~~ Considering the differences of the different variables occurring within/without cyclones present in the Gulf Stream and Kuroshio regions outside of a 750 km-radius of any cyclone over their
entire lifetime, we find that cyclones play only a secondary role in explaining the mean ~~states~~ state differences between the smoothed and realistic SST experiments.

1 Introduction

The Gulf Stream and Kuroshio ~~region~~ regions with their strong sea surface temperature (SST) gradients are preferential locations for cyclogenesis (e.g., Hoskins and Hodges, 2002; Nakamura et al., 2004) and are found to determine the location
20 and structure of storm tracks (e.g., Chen et al., 2010; Ogawa et al., 2012; Ma et al., 2015; Yao et al., 2018). Sensitivity tests with smoothed SSTs and a weaker SST gradient yield a reduced cyclone activity. In addition, these experiments feature an equator-ward shift of both the storm track as well as the upper level jet (e.g., Ma et al., 2015; Zhang et al., 2020) and a decrease of surface heat fluxes as well as precipitation on the warm side of the SST front (e.g., Kuwano-Yoshida et al., 2010b; Kuwano-Yoshida and Minobe, 2017). However, as it remains unclear if the latter changes can be attributed to the changes

25 in cyclone activity, we quantify the contribution of cyclones to the documented differences in the Gulf Stream and Kuroshio ~~region~~regions.

SST gradients also influence individual cyclone intensification (e.g., Sanders, 1986; Wang and Rogers, 2001; Jacobs et al., 2008), where the intensification has been associated to low-level baroclinicity originating from sensible heat fluxes (e.g., Hotta and Nakamura, 2011) and latent heating (e.g., Papritz and Spengler, 2015) along the SST front. However, other studies related
30 the intensification of individual cyclones in the western Atlantic to the low-level baroclinicity associated with the pronounced land-sea contrast (e.g., Brayshaw et al., 2009; Tsopouridis et al., 2020a). Thus, while several studies highlighted the sensitivity of cyclogenesis and the storm track to a smoothing of the SST (e.g., Nakamura et al., 2008; Kuwano-Yoshida and Minobe, 2017; Ma et al., 2017; Zhang et al., 2020), the impact of a weaker SST gradient on the intensification of individual cyclones remains unclear.

35 Randomly selecting 24 individual cyclones that occurred in the Gulf Stream region, de Vries et al. (2019) highlighted the reduction of surface latent heat fluxes and low-level baroclinicity when smoothing the SST. They, however, emphasised that these changes vary based on the position of each storm relative to the SST front. Similarly, Tsopouridis et al. (2020a) found cyclones following different pathways with respect to the SST front position to be associated with different characteristics. They, however, attributed the structural changes primarily to the absolute SST over which the cyclone is propagating rather
40 than the SST front itself. ~~Along the same lines~~This is consistent with the idealised simulations of cyclone development using different SST and SST gradients by Bui and Spengler (2021), who identified a primary dependence on cyclone development to absolute SSTs and only minor dependence on the SST gradient. Similarly, Booth et al. (2012) found ~~storm strength~~strength of storms to increase in the Gulf Stream region with increased SSTs, even if only a weak SST gradient is present. ~~Given Overall,~~
45 ~~with respect to either absolute SST or SST fronts needs to be further clarified~~indicates that both influence the development of cyclones.

In addition to low-level baroclinicity, upper-level forcing by the jet stream is known to contribute to cyclogenesis (e.g., Sanders and Gyakum, 1980; Uccellini et al., 1984; Sinclair and Revell, 2000; Yoshida and Asuma, 2004) and can influence cyclone intensification (e.g., Evans et al., 1994; Schultz et al., 1998; Riviere and Joly, 2006; Tsopouridis et al., 2020b). At the
50 same time, the very existence of the extratropical jet depends on the cyclones maintaining the storm track (Hoskins and Valdes, 1990; Holton and Hakim, 2012; Papritz and Spengler, 2015). Accordingly, using experiments with realistic and smoothed SSTs, Kuwano-Yoshida and Minobe (2017) argued that the increased cyclone activity over the SST front influences the upper-level jet, causing its meandering over the North Pacific.

In the light of this tight coupling between the jet and the storm track, it is not surprising that a smoothing of the SST can
55 affect the upper-level flow. Indeed both the storm track (e.g., Small et al., 2014; Ma et al., 2015; Piazza et al., 2016; Zhang et al., 2020) and the upper-level jet (e.g., Ma et al., 2017; O'Reilly et al., 2017) were shown to shift equatorward in the North Atlantic and Pacific ocean when the SSTs were smoothed. Kuwano-Yoshida and Minobe (2017) showed that smoother SST in the Kuroshio region resulted in a more zonally oriented storm track and argued that a weaker SST front is not able to anchor the upper-level flow. A southward shift of both the storm track density and the upper-level jet when smoothing the SST has

60 also been documented in the North Atlantic region by Piazza et al. (2016), though their shift of the storm track was smaller compared to the one in the study of Small et al. (2014). They related this difference by Small et al. (2014), which they related to the stronger SST smoothing in the study of Small et al. (2014). Based on the aforementioned arguments, a smoothing of an already climatologically weaker SST front in the Kuroshio region (e.g., Nakamura et al., 2004; Tsopouridis et al., 2020b) should have a comparatively minor impact on the storm track and the upper-level wind speed compared to the Gulf Stream region.

65 Focusing on mesoscale aspects of the atmospheric response to a smoothing of the SSTs, Piazza et al. (2016) documented a considerable decrease in the surface heat fluxes (30-50%) and convective precipitation (up to 60%) over the warm side of the North-Atlantic SST front after they removed small-scale SST features. Consistently, Zhang et al. (2020) found a similar, yet significantly smaller, decrease of the surface heat fluxes (5%) and precipitation (7%) within the Kuroshio and Oyashio confluence region. Atmospheric general circulation model (AGCM) experiments with real and smoothed SSTs revealed that the SST front is important to maintain convective precipitation (in line with Minobe et al., 2008), with the atmospheric response of the SST smoothing being stronger in the Gulf Stream than in the Kuroshio region (Kuwano-Yoshida et al., 2010b). Indeed, comparing differences in precipitation between the original and smoothed SSTs as well as between the Atlantic (Minobe et al., 2008) and the Pacific (Kuwano-Yoshida and Minobe, 2017), the decrease of precipitation is more pronounced in the Gulf Stream region. Thus, surface heat fluxes and precipitation are considerably affected by the strength of the SST gradient, with the effect being stronger in the Gulf Stream than in the Kuroshio region. However, whether surface heat fluxes and precipitation the time-mean distributions of such atmospheric patterns are only altered by the SST gradients or to what extent changes in the occurrence or intensity of cyclones contribute to their distribution remains ambiguous.

70 While the spatial distribution of surface wind convergence into a narrow band has been linked to strong SST gradients (Small et al., 2008), recent studies associated the mean state's characteristics in the Gulf Stream and Kuroshio regions to synoptic features (e.g., O'Neill et al., 2017; Parfitt and Seo, 2018). In particular, O'Neill et al. (2017), associated the existence of the Gulf Stream convergence zone with intense cyclones propagating in the region and highlighted the overall role of storms in shaping the mean state of the atmosphere in the northwest Atlantic. Similarly, Parfitt and Seo (2018) indicated that the near-surface wind convergence over the Gulf Stream and Kuroshio regions is influenced by extratropical cyclones, in particular by their associated fronts. Masunaga et al. (2020a, b), on the other hand, showed that strong cyclones and atmospheric fronts only have a minor contribution to the climatological mean convergence in the Gulf Stream and Kuroshio regions and that the main contribution is associated with weaker storms and fronts. However, given the weakness of these systems, it could be questioned how significantly the climatological contribution is associated with fronts and storms in general.

85 The ability of extratropical cyclones to strongly modulate the moisture transport (e.g., Ruprecht et al., 2002; Chang and Song, 2006) and precipitation (e.g., Bjercknes, 1922; Pfahl and Wernli, 2012; Hawcroft et al., 2012) is commonly accepted, a similar association between cyclones and. While surface heat fluxes can have a direct and indirect effect on cyclone development (e.g., Haualand and Spengler, 2020), the role of cyclones shaping heat fluxes is under debate. Some studies suggest a close relationship between the surface heat fluxes and cyclones in the midlatitudes on both synoptic (e.g., Alexander and Scott, 1997; Schemm et al., 2015; Dacre et al., 2020) and longer time scales (e.g., Parfitt et al., 2016; Ogawa and Spengler, 2019).

95 Using a more statistical approach, Zolina and Gulev (2003) argued that the surface fluxes mainly occur on synoptic time scales. However, based on a composite analysis Rudeva and Gulev (2011) ~~actually~~ indicated that cyclones in the North Atlantic do not contribute significantly to the climatological surface heat fluxes in this region. Furthermore, Tanimoto et al. (2003) noted that in regions with active ocean dynamics, such as along the western boundary currents, the SST anomalies mainly regulate the surface heat fluxes and not the cyclones.

100 To shed light to the outlined uncertainties, we assess the effect of a weak or strong SST gradient using an atmospheric general circulation model (AFES 3) based on simulations with realistic and smoothed SSTs in the Gulf Stream and Kuroshio ~~region~~regions. Our analysis of these simulations is twofold. Firstly, we follow the approach of Tsopouridis et al. (2020a, b, hereafter TSSa,TSSb) and investigate the effect of the smoothed SSTs on the structure of individual cyclones. Secondly, we consider pertinent variables within and outside a radius around cyclone centres in the Atlantic and Pacific basin throughout
105 their lifetime to examine the contributions of cyclones ~~on the different winter climatologies between to the winter climatologies~~ for the realistic and smoothed simulations (similar to Ma et al., 2015). This two-pronged approach allows us to establish a ~~link~~ between changes in the structure of connection between structural changes in individual cyclones and changes in the time-mean winter climatology. ~~As we will show, the impact on individual cyclones is markedly smaller compared to changes in the winter climatology, suggesting potential answers to the questions raised above.~~

110 2 Data and Methods

2.1 Data

We use data from version 3 of the ~~Atmospheric General Circulation Model (AGCM)~~ AGCM for the Earth Simulator (AFES) developed by the Earth Simulator Center of the Japan Agency for Marine Earth Science and Technology (~~Ohfuchi et al., 2004; Enomoto et al.~~ JAMSTEC, Ohfuchi et al., 2004; Enomoto et al., 2008; Kuwano-Yoshida et al., 2010a). This version of AFES has been first
115 used by Kuwano-Yoshida and Minobe (2017) and O'Reilly et al. (2017) ~~;~~ and has a horizontal resolution of T239 (approximately 50 km) and 48 sigma levels in the vertical. The model was integrated from 1 September 1981 to 31 August 2001, where we only focus on the winter periods December to February, hereafter DJF. ~~In the production of the AFES data~~ Throughout the time period, NOAA 0.25° daily SST data (Reynolds et al., 2007) were used as boundary conditions. For our analysis we use the AFES output on a 0.5° horizontal grid at 6 hourly intervals. More information about the model configuration can be found
120 in Kuwano-Yoshida and Minobe (2017).

~~Kuwano-Yoshida and Minobe (2017); using~~ Using AFES 3, Kuwano-Yoshida and Minobe (2017), produced two experiments for the North Pacific. Firstly, the control experiment (hereafter CNTL) that uses the original global SST data and secondly an experiment that uses smoothed SSTs over the ~~Northwestern Pacific~~ greater area around the Kuroshio Extension (hereafter SMTHK). They also composed an analogous experiment with spatially smoothed SSTs over the ~~Northwestern Atlantic~~ greater
125 area around the Gulf Stream (hereafter SMTHG). In both cases, the NOAA 0.25° daily SST were smoothed by applying a 1-2-1 running mean filter 200 times in the zonal and meridional direction. It is a three-point filter with the weights 0.25, 0.5 and 0.25, that has a sharp cutoff frequency, so that unwanted frequency components are effectively removed.

We use SST, latent and sensible heat fluxes, large-scale and convective precipitation, specific humidity at 850 hPa, and wind at 300 hPa for our analysis. We also compare the model simulations with the same variables from the ERA-Interim reanalysis that was created using a four-dimensional variational data assimilation scheme and a spectral truncation of T255 and 60 levels in the vertical (Dee et al., 2011).

2.2 SST front and jet stream detection

We identify the position of SST fronts using an objective frontal detection scheme based on the “thermal” method, as described in detail by ~~Tsopouridis et al. (2020a)~~ TSSa. This method has also been used to detect atmospheric fronts (Jenkner et al., 2010; Berry et al., 2011; Schemm et al., 2015). ~~To capture the most prominent SST fronts, we have chosen a SST gradient~~ We perform the detection using SST data filtered with a spectral truncation to T84 resolution and detect the SST fronts in the instantaneous SST field every 6-hours. We detect SST front lines in order to be able to define when cyclones cross the front. After thoroughly testing different thresholds for the two regions, we use a threshold of 2K/100km for the Atlantic and a smaller threshold of 1.25K/100km for the Pacific region. ~~The smaller threshold in the Pacific is due to the weaker SST gradient, as documented by Nakamura et al. (2004), and to capture the most prominent SST front lines. It was necessary to use different thresholds to accommodate the different SST gradients in the two ocean basins (Nakamura et al., 2004), which is also in line with the previous studies of TSSa~~ previous studies for the Atlantic ~~and TSSb for the Pacific (TSSa) and Pacific (TSSb).~~

To assess the potential impact of the SST smoothing on the upper levels, we detect the ~~jet stream position in both basins for the three experiments using the method and criteria of Spensberger et al. (2017), which identifies lines separating~~ position of jet following the algorithm of Spensberger et al. (2017). The algorithm detects jets by their axis, the line of maximum winds that separates cyclonic from anticyclonic wind shear ~~yielding automatically detected jet axes. The algorithm requires the wind maximum to be well-defined, but does not impose a strict minimum wind speed (Spensberger et al., 2017).~~

For the climatologies and composites we normalise the occurrence of both SST front lines and jet axis lines to account for the latitudinally varying area covered by grid cells. We achieve this by showing the average length of SST front line/jet axis line per unit area, hence the resulting unit of length per area. For details on the normalisation we refer to the jet climatology by Spensberger and Spengler (2020).

2.3 Cyclone detection and tracking

We employ the University of Melbourne cyclone detection and tracking algorithm (~~Murray and Simmonds, 1991a, b~~) (Murray and Simmonds, 1991a, b). The algorithm detects maxima in the Laplacian of the MSLP field and tracks them over time using a nearest-neighbour method together with the most likely direction of propagation (Murray and Simmonds, 1991a, b; Michel et al., 2018; TSSa; TSSb).

~~Analogous to TSSa; TSSb, we take into account only cyclones propagating for at least 12 hours (three consecutive time steps) in the Gulf Stream region (30–50°N and 290–310°E) and the Kuroshio Extension region (30–50°N and 145–170°E). We also consider cyclone tracks with at least five 6-hour time steps in the North Atlantic or the North Pacific and require the great circle distance between cyclogenesis and cyclolysis to be greater than 300 km in order to remove quasi-stationary systems.~~

160 The cyclone density pattern is in good spatial agreement with previous studies (Hanley and Caballero, 2012; Neu et al., 2013 ; TSSa; TSSb), successfully capturing the major regions of cyclone activity in both basins. Consistent with the results of TSSa and TSSb for ERA-Interim, we relate the small quantitative differences in the density climatology presented by both Neu et al. (2013) or Murray and Simmonds (1991a), who also used the Melbourne University algorithm, to the neglect of shallow and weak systems in our database.

165 **2.4 Classification of cyclone tracks based on their position relative to the SST front**

We categorise the identified cyclone tracks with a maximum intensification in either the Gulf Stream region (30-50°N and 290-310°E) or the Kuroshio region (30-50°N and 145-170°E) based on their propagation relative to the SST front. Analogous to TSSa and TSSb, for this classification we consider only cyclones with at least three 6-hour time steps in the Gulf Stream and Kuroshio regions. Firstly, we identify the shortest distance between each cyclone position and the SST front line for every
170 time-step along the cyclone track and define a vector pointing to the cyclone. Then, we follow TSSa and TSSb and focus on cyclones that always stay on the cold (C1) or warm (C2) side of the SST front, and those that cross the SST front from the warm to the cold side (C3).

We categorise the cyclones for the SMTHG and SMTHK experiments analogously to the CNTL experiment. However, as the SST gradient in the smoothed experiments is very homogeneous over a large region and thus too weak to qualify as a front,
175 we instead use the front positions from the CNTL experiment for the classification. We ~~thus categorise cyclone tracks based on similarities in their geographical location and propagation direction to achieve a consistent classification of the cyclone tracks across all AFES~~ use the same classification as in CNTL to be able to compare cyclones with geographically similar genesis locations and tracks across the experiments. For simplicity we still refer to C1-3 as the cold, warm, and crossing cases for the smooth (SMTH) experiments, even though, strictly speaking, no SST front is crossed.

180 **2.5 Decomposition of climatological differences**

In addition to the cyclone track classification, we ~~decompose the winter climatology into dates with and without cyclones present the Gulf Stream or the Kuroshio region, respectively. This decomposition allows us to assess how much of the climatological differences in the smoothed experiment arise while cyclones are present in the respective region, and thus how much of the climatological differences are directly or indirectly associated with cyclones. For this analysis, we consider~~
185 all cyclones that pass through either the Gulf Stream or the Kuroshio region during their evolution present a decomposition of the winter climatology for selected variables, where we conditioned the two composites on either occurring within or outside an area with a radius around a cyclone centre throughout the lifecycle of a cyclone. We performed this analysis for each ocean basin, irrespective of ~~their direction of propagation and the location of their~~ the direction of cyclone propagation and location of its maximum intensification. Consistent with the threshold on cyclone circumference in Wernli and Schwerz (2006) and the
190 analysis of Rudeva and Gulev (2011), we choose a radius of 750 km. We also obtained results for a radius of 500 and 1000 km, respectively (Fig. B1-Fig. B4). However, given that the results were not very sensitive to this choice, we focus on results with a radius of 750 km, for which most cyclone-related features, such as fronts, are included.

3 Results and Discussion

3.1 SST front and SST/SST gradient distribution between the experiments

195 Analysing the SST (Fig. 1a) and SST gradient (Fig. 2a) distribution in the Gulf Stream region (black box) for CNTL, we note a remarkable SST contrast across the Gulf Stream, which results in a strong SST gradient (Fig. 2a) and in locally well-confined SST front detections (Fig. 3a), consistent with an oceanographic viewpoint (Meinen and Luther, 2016). The SST front distribution also resembles the correspondent feature presented in TSSa for the same region, but based on a different period and dataset.

200 ~~Across the Kuroshio~~ In the Kuroshio (black box in Fig. 1d), we observe ~~an overall similar, but less spatially confined,~~ a similar but spatially less confined SST contrast compared to the Gulf Stream region (compare Fig. 1a with Fig. 1d), which results in a weaker SST gradient in the Kuroshio region (Fig. 2d). Consequently, the detected SST fronts are more ~~wide-spreadly distributed~~ wide-spread in the Pacific (Fig. 3b), but remain collocated with the region of the climatologically largest SST gradient (in line with, e.g., Tozuka et al., 2018; Wang et al., 2019). The less pronounced SST gradient in the Pacific compared
205 to the Atlantic is also evident in the ERA-Interim winter climatology (Fig. A1b,e), with the differences between the reanalysis and AFES simulations arising from the coarser SST resolution used in ERA-Interim prior to 2002 (e.g., Masunaga et al., 2015; Parfitt et al., 2017).

Compared to CNTL, the SSTs in SMTHG are smoother and their gradient is more widely distributed (compare Fig. 1a,b and Fig. ~~1a~~ 2a,b). In the western Atlantic, we also observe that the smoothing affects SSTs at a considerable distance from the Gulf
210 Stream SST front (Fig. 3a), for example reducing the SST to the east of the Florida Peninsula (Fig. 1a,b). At approximately 40°N, the SST differences exhibit a clear dipole, with increased SST to the north and decreased SST to the south, following the position of the SST front (Fig. 1c, Fig. 3a). The largest differences occur offshore off the central US East coast ($\Delta\text{SST} < -4\text{K}$) as well as off the coast of Newfoundland ($\Delta\text{SST} > 4\text{K}$; Fig. 1c).

The SST distribution in SMTHK (Fig. 1e) is similar to CNTL, though smoother, and the region with the strongest gradients
215 off the east coast of Japan is oriented slightly more zonally (Fig. 1d). Contrary to the Gulf Stream region, the SSTs outside the Kuroshio region remain largely unaffected. As in the Gulf Stream region, the SST differences between the two experiments follow a ~~bipolar~~ dipolar structure (Fig. 1f), but they are considerably weaker. The smoothing results in a maximum decrease (increase) of ~~2K~~ ~~(3K~~ 2 K (3 K) south (north) of the SST front (Fig. 1f), with the SST differences being more pronounced in the western part of the domain close to Japan (Fig. 1f).

220 3.2 Cyclone Density and Upper-level Wind

~~To evaluate the potential impact of the upper-level forcing on cyclones, we present the climatological position of the jet stream over the two ocean basins.~~ The position of the North Atlantic jet coincides with the location of the SST front (Fig. 3a), whereas the Pacific jet is located south of the Kuroshio SST front (Fig. 3b). The Pacific jet is stronger, meridionally more confined, and located over lower latitudes compared to the Atlantic jet, which is consistent with previous studies (e.g., Nakamura, 1992;
225 Spensberger and Spengler, 2020). Both the strength and the position of the jet axes distribution in the North Atlantic are




Figure 1. (a) Climatological SSTs in SST for DJF for (a) CNTL, (b) SMTHG, and (c) the difference between SMTHG and CNTL ($\text{SMTHG}-\text{CNTL}$ [K]). (d-f) as As (a-c), but for the North Pacific. The Gulf Stream and Kuroshio regions are marked with a black box, respectively.

analogous to the study of TSSa, while for the Pacific we observe fewer jets in the region to the south of Japan compared to TSSb.

The changes in the SST field between CNTL and SMTHG/SMTHK introduce differences in both cyclone density and the climatological jet stream position. We observe an equatorward shift in the maximum cyclone density in both the North Atlantic and North Pacific, particularly in the central and eastern part of the basins (Fig. 4a,c). We also notice an analogous shift in the upper-level jet in the North Atlantic (Fig. 4b), with negative anomalies in the northern part of the basin and positive anomalies mainly to the east of the Gulf Stream region. However, a similar shift of the upper-level jet is not observed in the North Pacific, where with the position of the jet remains remaining rather unchanged, with a more confined and zonal distribution of the jet and the jet being more confined and zonally distributed in SMTHK (Fig. 4d). We relate the lack of a shift in the upper-level jet in the North Pacific to the small SST difference after the SST smoothing ($\Delta\text{SST}=1\text{K}$; Fig. 1f), whereas in the Atlantic the jet is located near the region with the most pronounced SST changes ($\Delta\text{SST}>4\text{K}$; Fig. 1e).

Analogously, the sharper SST gradient in the Gulf Stream region (Fig. 2c,f) leads to larger changes in storm-track activity in the Northwestern Atlantic, with a more pronounced decrease in cyclone density in the Gulf Stream region compared to the Kuroshio region (Fig. 4a,c). This is consistent with Kuwano-Yoshida et al. (2010b) and in line with Nakamura et al. (2008), who found the storm-track activity in the Southern Ocean to be substantially weaker after the removal of the strong SST front.




Figure 2. (a) Climatological SST gradient in-for DJF for (a) CNTL [$K (100 \text{ km})^{-1}$], (b) SMTHG [$0.5 K (100 \text{ km})^{-1}$], and (c) the-difference between-SMTHG and-CNTL- $SMTHG-CNTL [K (100 \text{ km})^{-1}]$. (d-f) as-As (a-c), but for the North Pacific. The Gulf Stream and Kuroshio regions are marked with a black box, respectively.




Figure 3. (a) SST front distributions-distribution (blue shading, $\text{km of line } (100 \text{ km})^{-2}$) and jet axis distributions-distribution (orange-pale red shading, $\text{km of jet axis line } (1000 \text{ km})^{-2}$) for the North Atlantic. (b) as-As (a), but for the North Pacific

Thus, while differences in-While cyclone density (Fig. 4a) and jet occurrence (Fig. 4b) are generally collocated both shift equatorward in the Atlantic, they are displaced from each other in the SMTHG experiment, there is no clear relation between these different fields in the Pacific region (Fig. 4c,d), where a considerable increase in- For example, cyclone density in




Figure 4. (a) Difference in the climatological cyclone density between SMTHG and CNTL (SMTHG-CNTL, shading, 10^{-6} km^{-2}) and climatological cyclone density for CNTL (contours, first contour: $6 \times 10^{-6} \text{ km}^{-2}$, interval $3 \times 10^{-6} \text{ km}^{-2}$) for the CNTL experiment. (b) Difference in the climatological jet stream density (SMTHG-CNTL, shading, km of jet axis line (1000 km^{-2})) and climatological jet stream density for CNTL (contours, first contour: 400 km of jet axis line (1000 km^{-2}), interval: 200 km of jet axis line (1000 km^{-2})). (c,d) As (a,b), but for the Kuroshio region. The Gulf Stream and Kuroshio regions are marked with a black box, respectively North Pacific.

the Northeast Pacific (Gulf of Alaska, Fig. 4c) is contrasted by a decrease in the increases, whereas upper-level wind speed decreases in this region (Fig. 4d). Zhang et al. (2020) found a similar decrease in the upper-level response in the northeastern Pacific, expressed by differences of meridional eddy wind variance and eddy kinetic energy at 300 hPa. However, they showed that the upper-level decrease was not accompanied by a reciprocal negative anomaly in the low-level storm track and thus documented a different response of the SST smoothing in the lower and upper levels in the Pacific.

3.3 Categorisation Classification and Intensification of Cyclones

To assess the effect of smoothing the SSTs on the evolution of individual cyclones, we now restrict our focus to cyclones with maximum intensification in the Gulf Stream or Kuroshio region (details in section 2.4). For the Atlantic CNTL, 57 cyclones consistently stay on the cold side of the Gulf Stream SST front (C1), 27 cyclones stay on the warm side (C2), and 40 cyclones cross the SST front from the warm to the cold side (C3). In SMTHG, 62, 30, and 25 cyclones belong to C1, C2, and C3,




Figure 5. Pressure tendency (hPa h^{-1}) for the three categories relative to the time of maximum intensification for (a) CNTL and (b) SMTHG. Lines indicate the median and the shading the interquartile range. (c) Medians of the pressure tendency (hPa h^{-1}) for the three categories relative to the time of maximum intensification for CNTL (solid lines) and SMTHG (dashed lines). (d-f) As (a-c), but for the North Pacific.

respectively. Comparing these numbers, we notice that the number of cyclones staying on either side of the SST front is
 255 nearly unaffected by the smoothing, whereas the number of crossing cyclones is substantially reduced. This implies an overall
 reduction in the number of cyclones, which is in line with the decreased cyclone density along the Gulf Stream SST front in
 SMTHG (Fig. 3a, Fig. 4a).

In the Kuroshio region, the number of cyclones in C1 (86/81) and C3 (59/60) is more or less unchanged between CNTL/
 SMTHK, whereas cyclones in C2 (24/14) decrease slightly in number. The ~~restricted~~-small number of cyclones in C2, particu-
 260 larly in SMTHK, implies some uncertainty for the corresponding results. Note that in contrast to the Atlantic, there is no
 reduction in cyclones crossing the SST front, potentially because the SST gradient in the Pacific is already comparatively weak
 in CNTL (compare Figs. 2a, d).

The more pronounced reduction of cyclones crossing the SST front in SMTHG compared to SMTHK highlights the sig-
 nificance of a strong SST gradient to anchor the position of the storm track, as discussed in previous studies (e.g., Nakamura
 265 et al., 2004, 2008; Sampe et al., 2010). Our results confirm these studies not only based on the number of cyclones, but also
 based on the location of cyclones during their maximum intensification. In CNTL, most cyclones undergo their most rapid
 intensification close to the SST front (Fig. A3b), whereas the location of most rapid intensification is distributed over a wider
 region in SMTHG (Fig. A3e).

Among the three categories, Atlantic cyclones of C3 and C1 are deepening the most in CNTL, with a maximum 6-hourly intensification corresponding to approximately 28 and 25 hPa/day, respectively. Conversely, cyclones of C2 are characterised by a weaker intensification throughout their evolution (Fig. 5a). The results for C2 are statistically significant around the time of maximum intensification (from -6h to 6h), as the 50th percentile of C2 coincides with the 75th percentile of C1 and with the 75-100th percentile (not shown) of C3. Moreover, we notice that the median (50th percentile) of C2 is always above the ones for C1 and C3 during the 48 hour time period in both experiments for the Gulf Stream region, indicating a clear tendency for higher intensification in C1 and C3, compared to C2. In SMTHG, cyclones intensify similarly fast as in CNTL (Fig. 5a,b). In particular, cyclones of C3 experience only a slightly weaker intensification in SMTHG, although the SST front that they cross is barely existing in SMTHG. These results support the findings of TSSa, who related the intensification of cyclones in the Gulf Stream region to the low-level baroclinicity associated with the land-sea contrast, hypothesising that a strong SST gradient only weakly modifies the deepening of the cyclones. Nevertheless, the distinction between ~~the these~~ categories becomes less clear in SMTHG, where the interquartile ranges of the three categories overlap considerably.

In the Kuroshio region, cyclones of C3 are ~~on average~~ deepening the fastest (approx. ~~30~~ 30 hPa/day), followed by cyclones of C1 (approx. 25 hPa/day; Fig. 5d). There is a considerable overlap in the interquartile range for the 3 categories and a larger variability compared to the Gulf Stream region. Nevertheless, the median of C2 is also above the ones for C1 and C3 before the time of maximum intensification, indicating a clear tendency for weaker intensification in C2, compared to C1 and C3. In line with TSSb, C2 becomes the category with the larger deepening rate among the three categories after the time of maximum intensification. ~~In~~ However, in contrast to TSSb, cyclones of C2 ~~, however,~~ deepen the least before their maximum intensification in both CNTL and SMTHK (Fig. 5d,e). We relate this difference to cyclones in the AFES simulations being located further away from an upper-level jet stream than in the ERA-Interim data used in TSSb (not shown), where the upper levels ~~proved appear~~ to substantially influence the intensification of cyclones in the Kuroshio region. Moreover, the limited number of cyclones in the AFES simulations lowers the statistical robustness of these results (24 cyclones in CNTL compared with 97 cyclones in TSSb) ~~lowers the statistical robustness of these results.~~

In SMTHK, the cyclones in the three categories have similar pressure tendencies as in CNTL (Fig. 5e), which becomes even more apparent when comparing the median of their pressure tendencies (Fig. 5f). In particular, C3 ~~on average~~ intensifies virtually identically in the two experiments. In the light of the corresponding findings for the Gulf Stream region, this result is not surprising. Even for the more focused Gulf Stream SST front, the smoothing had only a minor impact on the evolution of C3 cyclones (Fig. ~~25~~ c,f). A more pronounced effect is evident for C2 cyclones in the Kuroshio region, with cyclones intensifying slightly less after the smoothing (Fig. 5f). However, the number of cyclones is even smaller in SMTHK (14 cyclones) than in CNTL, making it difficult to draw conclusions from this difference.

Considering the evolution of the SST underneath the cyclone core, cyclones of C1 propagate over comparatively low SSTs, because they remain on the cold side of the SST front in both regions (Fig. 6a,d). In contrast, cyclones of C2 propagate over approximately 16 K higher SSTs than cyclones of C1 in the North Atlantic (Fig. 6a) and over 12K higher SSTs in the North Pacific (Fig. 6d) during maximum intensification. However, during their evolution, cyclones gradually propagate towards lower SSTs. Cyclones of C3 propagate over higher SSTs at an early stage of their development and over lower SSTs after crossing




Figure 6. SSTs (K) for the three categories relative to the time of maximum intensification for (a) CNTL and (b) SMTHG. Lines indicate the median and the shading the interquartile range. (c) Medians of the SSTs (SST [K]) underneath the cyclone for the three categories relative to the time of maximum intensification for CNTL (solid lines) and SMTHG (dashed lines). (d-f) As (a-c), but for the North Pacific.

the SST front. The latter is associated with the cross-frontal SST difference, which is more pronounced in the Atlantic (Fig. 305 6a) than in the Pacific (Fig. 6d) due to the sharper Gulf Stream SST front (consistent with Nakamura et al., 2004; Joyce et al., 2009; TSSb).

After the SST smoothing, cyclones of C1 propagate over approximately 1 K higher SSTs in both regions and cyclones of C2 over 1-2 K lower SSTs. Further, cyclones of C3 in the smoothed experiments experience a less sharp decrease in SST across the SST front compared to the CNTL experiment (Fig. 6b,c,e,f). The SST differences introduced by the smoothing are 310 more pronounced before maximum intensification, because at this stage cyclones typically propagate in the western part of the regions of interest where the SST differences associated with the smoothing tend to be larger (Fig. 1c,f).

Overall, the considerable reduction in the number of cyclones of C3 after the smoothing of the SST in the Atlantic highlights the anchoring effect of a strong Gulf Stream SST front on the storm track. On the other hand, the already weak SST gradient in the Kuroshio prior to the smoothing leads to minor SST differences between CNTL and SMTHK and ~~thus~~ indicate indicates 315 of cyclones that cross the SST front. The rather similar cyclone intensification between the experiments indicate indicates that the SST gradient is not particularly important for the intensification of individual cyclones (consistent with TSSa; TSSb).

Our results thus suggest that the SST smoothing does not result in significant differences in the characteristics of individual cyclones, which is consistent with Bui and Spengler (2021), who found that cyclone development is more sensitive to absolute SST than SST gradients. This result is supported by cyclone-relative composites of, for example, surface heat fluxes and pre-

320 cipitation, which exhibit only minor differences between the ~~smooth-SMTH~~ and CNTL experiments for all cyclone categories (not shown). Nevertheless, we observe a latitudinal shift in the storm track and the jet stream climatologies associated with the SST smoothing (Fig. 4) and previous studies documented a significant reduction in the climatological surface heat fluxes and precipitation (e.g., Kuwano-Yoshida et al., 2010b; Kuwano-Yoshida and Minobe, 2017). This discrepancy raises the questions how the largely unaffected characteristics of individual cyclones relate to the evident changes in the climatological mean state
325 of the storm track. In the following, we turn to this question by ~~decomposing the winter climatology into dates with and without cyclones propagating through either the Gulf Stream or Kuroshio region~~estimating the contribution of cyclones to the observed climatological differences between the experiments.

3.4 Contribution of cyclones to the climatological differences introduced by smoothing the SST

3.4.1 Surface Heat Fluxes

330 In the CNTL climatology, we observe a maximum of latent and sensible surface heat fluxes on the warm side of both the Gulf Stream and the Kuroshio SST front (Fig. 7a,e). Peak latent heat fluxes are slightly offset to the south of the peak sensible heat fluxes. TSSb associated this offset with the increase of sea surface saturation mixing ratio with increasing SSTs, following the Clausius-Clapeyron relation. In the North Atlantic, latent and sensible heat fluxes exceed 350 W m^{-2} and 80 W m^{-2} , respectively (Fig. 7a). Sensible heat fluxes are slightly larger in the North Pacific, whereas latent heat fluxes remain slightly
335 weaker (Fig. 7e).

The surface heat fluxes are similarly distributed in the ERA-Interim dataset, though latent heat fluxes in CNTL are considerably larger compared to ERA-Interim (compare Fig. 7a with Fig. A1c, and Fig. 7e with Fig. A1f). This discrepancy most likely arises from the difference of the SST resolution between the AFES and ERA-Interim, with the latter having a lower resolution prior to 2002. Comparing the CNTL fluxes with the ERA-Interim winter climatology after 2002, the differences are
340 significantly reduced (not shown).

The SST smoothing affects the amount of surface heat fluxes in both regions, though to a different extent. In the Gulf Stream region, we observe considerably weaker surface heat fluxes (Fig. 7b) along the weaker SST gradient (Fig. 2c). The maximum decrease of the latent heat fluxes is of the order of 120 W m^{-2} , while the reduction of sensible heat fluxes exceeds 30 W m^{-2} (in line with Small et al. (2014)). A slight increase of surface heat fluxes is observed mainly to the northeast and less in
345 the southern parts of the Gulf Stream region (Fig. 7b), consistent with the increase of SSTs due to the smoothing (Fig. 1c). This dipole of positive and negative anomalies of the surface heat fluxes is less pronounced in the Kuroshio region (Fig. 7f), reaching only about half the amplitude compared to the Gulf Stream region. We attribute the reduced amplitude to the smaller impact of the SST smoothing on the SST distribution in the Kuroshio region (Fig. 1f).

To estimate the role of cyclones ~~being present~~ for the differences when smoothing the SST, we ~~subdivide the differences between SMTHG/SMTHK and CNTL into time-steps with or without cyclones in the Gulf Stream or Kuroshio region, respectively~~decompose the winter climatology considering the surface heat fluxes occurring within and outside of a radius of 750 km around the cyclones' center over their entire lifetime (Fig. 7c,d,g,h). ~~In the Gulf Stream region, cyclones are present~~

about a third of the time-steps in both CNTL and SMTHG. In the Kuroshio region, cyclones are slightly more prevalent, and a cyclone is present in about 40% of the time-steps. This decomposition allows us to assess how much of the climatological differences are associated with cyclones.

The division into cyclone/no-cyclone time-steps shows that the climatological differences between CNTL and SMTHG/SMTHK predominantly arise when no cyclones are present in the respective region. We do not consider heat fluxes associated with cyclones (Fig. 7c,d,g,h). The contribution from no-cyclone time-steps is larger. On the other hand, the contribution from cyclones to the climatological heat flux differences is smaller in amplitude and the distribution of the flux anomalies more less closely resembles the climatological differences between the experiments (Fig. 7b,f). The largest differences in Fig. 7d arise close to the climatological SST front position and are clearly connected to the presence or absence of a sharp SST front. This can be explained by a cold air mass transitioning over a SST front experiencing less surface heating when the smoothing reduced the SSTs on the warm side of the SST front in SMTHG and SMTHK (see Fig. 1c,f; Zolina and Gulev, 2003 and Vanniere et al., 2017). The differences are more pronounced in the Gulf Stream region (Fig. 7c,d) compared to Kuroshio (Fig. 7g,h) due to the larger total mean differences of surface heat fluxes after the SST smoothing in the Atlantic (Fig. 7b,f).

Our results indicate that the smoothing of the SST front has only a minor effect when cyclones are present. We consider cyclones, which is confirmed by a cyclone-relative composite analysis, where surface heat fluxes are only moderately reduced by the smoothing of the SST front (not shown). Our findings are also in line with Rudeva and Gulev (2011), who indicated that cyclones in the North Atlantic sector do not contribute significantly to the climatological surface heat fluxes in the region.

3.4.2 Precipitation

The precipitation distribution in the Atlantic in CNTL is characterised by a maximum of large-scale ($> 6 \text{ mm day}^{-1}$) and convective precipitation ($> 4 \text{ mm day}^{-1}$) along the Gulf Stream SST front extending eastward (Fig. 8a). The maximum values of precipitation are located along the SST front (Fig. 3a), with a well-defined convective rain-band displaced towards the warm side of the Gulf Stream SST front, consistent with the findings of Kuwano-Yoshida et al. (2010b).

In the Pacific, we observe an analogous spatial distribution of the precipitation pattern, but the amplitude is somewhat larger than in the Gulf Stream region (compare Fig. 8a with Fig. 8e). Further, the peaks of large-scale and convective precipitation are more collocated in the Kuroshio region, likely related to the weaker SST gradient (Fig. 2a,d) western North Pacific compared to the North Atlantic. When compared to ERA-Interim (Fig. A2a,d), there is a good resemblance of the spatial distribution of precipitation, but we notice higher large-scale and convective precipitation in AFES, consistent with the larger latent heat flux.

Analogous to the surface heat fluxes, the smoothing of the SSTs affects precipitation in both regions. In the North Atlantic, the smoothing leads to a remarkable decrease of precipitation (Fig. 8b), in line with the study of Kuwano-Yoshida et al. (2010b). In the North Pacific, however, precipitation is reduced considerably less, with both large-scale and convective precipitation being reduced by less than 1 mm day^{-1} (consistent with Zhang et al. (2020)). We relate the more pronounced reduction of precipitation in the Gulf Stream to the originally sharper SST gradient (Fig. 2). We also observe a similar dipole pattern with reduced precipitation over the Gulf stream and Kuroshio core as well as increased precipitation in the southeast of both regions




Figure 7. (a) Climatological latent (shading, W m^{-2}) and sensible heat fluxes (contours, W m^{-2}) for the North Atlantic for CNTL. (b) Difference-Statistically significant (>95%) difference of the heat fluxes-flux climatologies between SMTHG and CNTL (SMTHG-CNTL) for latent (shading) and sensible heat fluxes (grey contours, interval: 10 W m^{-2} , zero contour omitted). (c,d) As (b) ,but only considering the dates with separated for fluxes within (c) or without and outside (d) cyclones present in the Gulf Stream region a radius of 750 km around a cyclone centre. (e-h) As (a-d), but for the Kuroshio region. The Gulf Stream and Kuroshio regions are marked with a black box, respectively. Panels (c,d,g,h) are scaled such that their sum yields panels (b) and (f), respectivelyNorth Pacific.

(Fig. 8b,f). This equatorward shift in precipitation is consistent with the equatorward shift of the storm track in the smoothed experiments (Fig. 4).

Among the two types of precipitation, convective precipitation is more sensitive to the SST smoothing. In SMTHG, the mean convective precipitation is reduced by half compared to CNTL and the narrow convective precipitation band observed in CNTL largely disappears in SMTHG (not shown). This finding is in line with Minobe et al. (2008) and Kuwano-Yoshida et al. (2010b), who showed that the Gulf Stream SST front is crucial for the distribution and amount of convective precipitation and found convective precipitation to be significantly reduced after heavily smoothing the SST distribution in their simulations with the same model.

Compared to the surface heat fluxes, the precipitation associated with cyclones is more influenced by the SST smoothing, in particular in SMTHG. There is a noticeable reduction in convective precipitation in the Gulf Stream region just south of the climatological position of the SST front (Fig. 8c), the region featuring the strongest decrease of SST due to the smoothing (Fig. 1c). This finding supports TSSa, who found convective precipitation to be closely related to the SSTs underneath the cyclone core. When no cyclone is present in the Gulf Stream region Taking into account the precipitation, which is not associated with cyclones, the SST smoothing results in a reduction of both large-scale and convective precipitation almost everywhere in the Gulf Stream region western North Atlantic (Fig. 8d).




Figure 8. (a) Climatological large-scale precipitation (shading, mm day^{-1}) and convective precipitation (grey contours, interval: 1, mm day^{-1} , zero contour omitted) for the North Atlantic for CNTL. (b) Difference-Statistically significant (>95%) difference of the precipitation climatologies between SMTHG and CNTL (SMTHG-CNTL) for the large-scale (shading) and convective precipitation (grey contours, interval: 1, mm day^{-1} , zero contour omitted). (c,d) As (b) ~~-, but only considering the dates with separated for large-scale and convective precipitation within~~ (c) ~~or without and outside~~ (d) ~~eyelones present in the Gulf Stream region~~ a radius of 750 km around a cyclone centre. (e-h) As (a-d), but for the Kuroshio region. ~~The Gulf Stream and Kuroshio regions are marked with a black box, respectively. Panels (c,d,g,h) are scaled such that their sum yields panels (b) and (f), respectively~~North Pacific.

Overall, cyclones account for a larger fraction of the precipitation differences than they did for the difference in surface heat fluxes when comparing CNTL and SMTHG. This result is also in line with Hawcroft et al. (2012), who found the winter precipitation in the Northern Hemisphere to be associated with extratropical cyclones. They also showed that the contribution of cyclones in the Gulf Stream region accounts for 55-80% of the total DJF precipitation. This considerable fraction suggests that precipitation should significantly shift equatorward along with the cyclone track density, which is consistent with our analysis (Figs. 4a, 8c).

In the Kuroshio region we note a rather equal influence of the SST smoothing on precipitation when cyclones are present or absent when we consider precipitation related or unrelated with cyclones. These differences mainly concern the large-scale precipitation and are more evident to the east of the Kuroshio region in the no-cyclone case in the central North Pacific (Fig. 8h), forming a dipole of reduced precipitation to the north and increased precipitation to the south, similar to the Atlantic (Fig. 8d). Interestingly, we note a slightly higher decrease of large-scale precipitation (approx. 0.25-0.5 mm day^{-1}) when eyelones are present in the Kuroshio region, which is most likely related to a diminished cyclone density in SMTHG compared to CNTL precipitation is not associated with cyclones in the vicinity of the Kuroshio, becoming less pronounced though when increasing the radius from 750 km to 1000 km and thus including precipitation from features that are more distant from the cyclone center (Fig. 4eB4d,f).




Figure 9. (a) Climatological specific humidity at 850 hPa (shading, g kg^{-1}) for the North Atlantic for CNTL. (b) Difference-Statistically significant (>95%) difference of climatological specific humidity between SMTHG and CNTL (SMTHG-CNTL). (c,d) As (b) ~~but only considering the dates with separated for specific humidity within (c) or without and outside (d) cyclones being present in the Gulf Stream region~~ a radius of 750 km around a cyclone centre. (e-h) As (a-d), but for the Kuroshio region. ~~The Gulf Stream and Kuroshio regions are marked with a black box, respectively. Panels (c,d,g,h) are scaled such that their sum yields panels (b) and (f), respectively~~ North Pacific.

3.4.3 Specific Humidity at 850 hPa

Higher values of specific humidity are observed to the south of the Gulf Stream and Kuroshio ~~region~~ regions (Fig. 9a,e) due to the Clausius-Clapeyron relation with higher SSTs (Fig. 1a,d). The specific humidity maximum exceeds 6 g kg^{-1} in the Gulf Stream region, while lower maximum values (5 g kg^{-1}) are found in the southeastern part of the Kuroshio region (Fig. 9a,e).
 420 The values are comparable to ERA-Interim (Fig. A2b,e).

Analogous to the surface heat fluxes (Fig. 7b,f), smoothing the SST causes a noticeable decrease in specific humidity to the south of the SST front (Fig. 9b,f), where the SSTs are lower than in CNTL (Fig. 1c,f). The decrease in specific humidity is more pronounced in the Gulf Stream region, following the larger SST decrease in the Atlantic when the SSTs are smoothed. Our findings are consistent with the results of Zhang et al. (2020), who analysed the meridional eddy specific humidity flux
 425 instead of specific humidity.

For specific humidity, cyclones account only for a small part of the climatological differences between CNTL and SMTHG (Fig. 9c,d). In addition to the larger amplitudes of the differences when ~~no cyclones are present in the Gulf Stream, also specific humidity is not associated with cyclones (Fig. 9d)~~, the structure is slightly different, with the negative anomalies being more zonally oriented than when ~~eyclones propagate in the region~~ considering the contribution from cyclones (Fig. 9c). In the
 430 presence of eyclones latter case, specific humidity is generally reduced in SMTHG the Gulf Stream region, whereas a slight increase is observed in the central part of the North Atlantic, most likely related to the southeastward shift of the storm track

after the SST smoothing (Fig. 9c). ~~In contrast, in the absence of cyclones, and Fig. 4a). Regarding the specific humidity which is not related to cyclones, we in contrast observe~~ a slight increase ~~in specific humidity is evident~~ to the north of the SST front, where the smoothing leads to an SST increase (Fig. 9d). ~~The largest differences~~ Consistently, the largest decrease evident in Fig. 9d ~~arise~~ towards the warm flank of the climatological SST front ~~and are is~~ clearly related to the largest decrease in the SST between SMTHG and CNTL (Fig. 1c). ~~The, and the~~ more pronounced decrease of specific humidity ~~in the absence of cyclones in the Gulf Stream region~~ resembles the decrease in surface heat fluxes (Fig. 7e,d), ~~which was also larger in the absence of cyclones when neglecting the contribution of cyclones for both variables.~~

~~In the Kuroshio region, the difference~~ Consistent with the results in the Atlantic, in the North Pacific we acquire larger differences in specific humidity (Fig. 9f) ~~is distributed more equally between time-steps with and without cyclones present (Fig. 9g,h). Indeed, cyclones even contribute more to the reduction in specific humidity in the southern half of the Kuroshio, when the latter is not associated with cyclones propagating in the region (Fig. 9g) than no-cyclone time-steps (Fig. 9h). The reduction in specific humidity likely explains the reduction of location of a maximum decrease of approximately 0.3 g kg^{-1} coincides with the region of the largest SST decrease (2K, Fig. 1f), located to the south of Japan. Apart from this reduction in specific humidity to the western part of the basin, there is an equivalent increase in specific humidity to the east, over the central North Pacific ocean (Fig. 9h), most likely triggering the increase of the large-scale precipitation in the same location (Fig. 8g,h). Nonetheless, in absolute terms these differences remain small.~~

~~We hypothesise that the different distribution of specific humidity between the dates with and without cyclones in the two regions arises from the different ratios of cyclones being present or absent in the Gulf Stream and the Kuroshio region, respectively. Cyclones are present in the Gulf Stream region in about 30% of the time-steps, whereas the respective percentage is higher in the Kuroshio region (about %40 of the time-steps). Therefore, the less pronounced decrease in humidity in the cyclone days in the Atlantic is most likely related to the smaller amount of time-steps, whereas the more equal split in decrease of specific humidity in the Pacific is attributable to the more equal split into cyclone/no-cyclone days~~ precipitation observed in the exact region (Fig. 8h).

Apart from the well-established Clausius-Clapeyron relationship between SSTs and moisture, several studies indicate the leading role of cyclones on the poleward transport of moisture (e.g., Peixoto and Oort, 1992; Nakamura et al., 2004; Chang and Song, 2006; Newman et al., 2012). However, ~~based on the subdivision of the winter climatology into cyclone/no-cyclone dates,~~ our results support instead that the SST is the dominating factor determining the distribution of specific humidity, with cyclones playing only a modulating role.

460 3.4.4 Upper-level Wind Speed at 300 hPa

In CNTL, the strongest climatological winds reach 40 m s^{-1} in the Gulf Stream region and occur in a southwest (SW) to northeast (NE) tilted band (Fig. 10a) that is located close to the position of the SST front (Fig. 3a). In the Kuroshio region, the climatological jet is more zonal than in the North Atlantic, exceeds 60 m s^{-1} (Fig. 10e), and is located somewhat to the south of the SST front (Fig. 3b). As for almost all the other fields, there is a good agreement between ERA-Interim and the AFES simulations, with a slightly reduced maximum wind speed in the Gulf Stream region (approx. 5 m s^{-1}) and the maximum wind




Figure 10. (a) Climatological wind speed at 300 hPa (shading, m s^{-1}) for the North Atlantic for CNTL. (b) Difference Statistically significant (>95%) difference of the wind speed climatology between SMTHG and CNTL (SMTHG-CNTL). (c,d) As (b), but only considering the dates with separated wind speed within (c) and without outside (d) cyclones being present in the Gulf Stream region a radius of 750 km around a cyclone centre. (e-h) As (a-d), but for the Kuroshio region. The Gulf Stream and Kuroshio regions are marked with a black box, respectively. Panels (c,d,g,h) are scaled such that their sum yields panels (b) and (f), respectively North Pacific.

speed being geographically less extended over the North Pacific in the AFES climatology compared to ERA-Interim (compare Fig. 10a,e and Fig. A2c,f).

In both basins, we observe decreasing (increasing) wind speeds to the north (south) of the climatological jet position after the SST smoothing (Fig. 10b,f). This dipole is more pronounced downstream, in the central and eastern North Atlantic and Pacific, respectively. Kuwano-Yoshida and Minobe (2017) documented a similar difference pattern for the North Pacific jet.

The presence (Fig. 10c) or absence (Fig. 10d) of cyclones in the Gulf Stream region does not have a considerable imprint on the distribution of the observed upper-level wind speed differences, with dates with cyclones present accounting for less of the climatological wind differences. In the Pacific, however, the contribution from cyclones determines the overall dipolar structure of the In the North Atlantic, the wind speed differences at 300 hPa that is not associated with cyclones (Fig. 10d) explains more of the difference in the wind speed climatology (Fig. 10f-h). In contrast, in the absence of cyclones, wind speeds are reduced throughout the Kuroshio region as well as in most parts of the North Pacific b), when compared to the wind speed associated with cyclones in the region (Fig. 10h).

This discrepancy between the contributions of cyclones c). This becomes more apparent in the North Atlantic versus North Pacific response to the smoothed SSTs might arise from the different relative shifts of cyclones and jets in these sectors Pacific, where cyclones contribute even less (Fig. 10g) to the climatological differences between SMTHK and CNTL (Fig. 4). Whereas cyclones and jets shift southward in tandem in the North Atlantic, it is mainly the cyclones that shift southward in the North Pacific, while the North Pacific jet intensifies in-place. Thus, the smoothed SSTs in the Kuroshio region allow cyclones

to propagate closer to the climatological position of the Pacific jet, thereby intensifying its eddy-driven component, while reducing the occurrence of a separate eddy-driven jet further north. This interpretation explains both the observed focusing and strengthening of 10f) compared to the respective results for the North Atlantic (Fig. 10c).

The higher contribution of cyclones to the observed differences in upper-level climatological wind speed in the Atlantic compared to the Pacific is consistent with previous studies indicating that the Pacific jet (Figs. 4d, 10f), and the larger contribution of cyclones to the upper-level response to the smoothed SSTs (Fig. 10g, h) is externally (more thermally) driven, as opposed to the Atlantic jet, which is more eddy-driven (e.g., Lee and Kim, 2003; Li and Wettstein, 2012).

490 4 Conclusions

We quantified and attributed differences in the atmospheric response when using realistic (CNTL) and smooth SSTs for the North Atlantic (SMTHG) and North Pacific (SMTHK), respectively, based on simulations with the AFES 3 model. The CNTL simulation compares well to ERA-Interim, except for considerably larger latent heat fluxes in CNTL, but these are most likely associated with the lower SST resolution in ERA-Interim prior to 2002. Overall, the AFES model successfully captured the distribution of the different variables in both ocean basins. Given the stronger SST gradient in the Atlantic, the effect of the smoothing on the SST front yields stronger SST differences between the CNTL and the respective smooth experiments (see Fig. 1c,f) as well as a distinctly stronger reduction of the SST gradient in the Atlantic compared to the Pacific (see Fig. 2c,f).

We first examined the impact of the smoothing of the SST on the intensification of individual cyclones. Considering only cyclones with a maximum intensification in the Gulf Stream or the Kuroshio region, we classified them into 3 categories based on their propagation relative to the SST front, where cyclones either always stay on the cold (C1) or warm (C2) side of the SST front or they cross the SST front from the warmer to the colder side (C3). Similar deepening rates for all these cyclone-categories across all experiments reveal the rather minor role of the SST gradient on the intensification of individual cyclones. This result is valid for both ocean basins, though it is particularly relevant for the Gulf Stream region where the SST smoothing dramatically weakens the strong SST gradient.

Considering all cyclones propagating in either the North Atlantic or the North Pacific, irrespective of their direction of propagation, stage of evolution, and their location of maximum intensification, we found the cyclone density in the storm track to decrease when the SSTs are smoothed in the Kuroshio and even more so in the Gulf Stream region. We relate the different response of the cyclone densities for the two regions to the more pronounced reduction of the SST gradient in SMTHG for the Atlantic compared to SMTHK for the Pacific. Overall, we observe an equator-ward shift in cyclone density in both regions, which is more pronounced over the central and eastern parts of the two ocean basins. Both cyclone density differences have a distinct SW-NE tilt, basically following the storm track. An analogous southward shift is observed in the upper-level jet for the North Atlantic, whereas for the North Pacific such a shift is absent and the difference between the experiments instead reveals a more meridionally focused and zonally extended jet for smoother SST.

We found a considerable decrease of both latent and sensible heat fluxes along the SST front when smoothing the SSTs, which was more pronounced across the SST front in the Gulf Stream region compared to the Kuroshio. Analogous to the surface

heat fluxes, precipitation in the Gulf Stream region is strongly reduced when smoothing the SST front, which is particularly evident for convective precipitation on the warm side of the Gulf Stream SST front. However, both types of precipitation are only slightly affected by the SST smoothing in the Kuroshio region. Differences in specific humidity at 850 hPa feature a similar reduction after smoothing the SST. The weaker reduction of moisture and precipitation in the Kuroshio region is related to the smaller differences in the SST and SST front between CNTL and the smoothed fields in the Pacific compared to the Atlantic.

To clarify whether the differences between the ~~realistic and smooth~~ CNTL and SMTH experiments stem directly from ~~differences in the SST or from the occurrence of cyclones~~, we subdivided the winter climatology into days with and without ~~cyclones present in the Gulf Stream or Kuroshio region~~ cyclones interacting with the SST and SST gradient, we considered selected variables within and outside an area with a radius of 750 km around cyclone centres propagating in either the North Atlantic or the North Pacific. We found ~~the contribution from time-steps without cyclones in the Gulf Stream or Kuroshio region to more closely resemble~~ that the surface heat fluxes that are associated with cyclones in both basins do not considerably contribute to the climatological differences in surface heat fluxes between the CNTL and SMTH experiments. Differences in precipitation, however, were more closely associated with cyclones being present in the Gulf Stream or Kuroshio region propagating in either the North Atlantic or the North Pacific.

~~Differences~~ For specific humidity, cyclones have only a minor contribution to the climatological differences between CNTL and SMTHG/ SMTHK, with a more evident decrease in specific humidity in the Gulf Stream region between CNTL and SMTHG are mainly attributable to days when cyclones are not present in the region. For the Kuroshio, however, the differences are almost equally attributed between time-steps with and without cyclones present in the region. Likewise, while cyclones in the Gulf Stream region do not seem to have a considerable imprint on the differences in Atlantic, arising from a considerable decrease in SST in the vicinity of the Gulf Stream. In contrast, both humidity and SST are not changed as significantly in the Pacific sector. Our results support that the SST is the dominating factor determining the distribution of specific humidity, with cyclones playing only a modulating role.

Similar to the surface heat fluxes and the specific humidity at 850 hPa, we found cyclones to only play a secondary role in explaining the upper-level wind, differences in the Kuroshio region are mainly attributable to time-steps when cyclones are present. We associate this discrepancy between the two basins to cyclones in the Pacific propagating more closely (300 hPa) wind speed differences arising from the SST smoothing. Notwithstanding this secondary role, Atlantic cyclones contribute more to the climatological jet position in SMTHK, thereby intensifying its differences than Pacific cyclones, which is consistent with previous studies indicating that the Atlantic jet is more eddy-driven component than the Pacific jet.

Overall, our analysis highlights that SST fronts only have a minor impact on the characteristics and intensification of individual cyclones propagating in the Gulf Stream or Kuroshio region. Nevertheless, we found a large-scale response with decreased cyclone density as well as an equator-ward shift of cyclone densities associated with the SST smoothing. Analogously, the upper-level jet shifts equator-ward in the North Atlantic, while the Pacific jet strengthens and extends more zonally eastward along its climatological location, which can be associated with the more collocated storm track. The effect of the SST smoothing was stronger in the Gulf Stream region, where the smoothing has a larger impact on the climatologically sharper SST front.

555 ~~Subdividing time-steps into those with and without cyclones~~ Considering pertinent variables within or outside an area with a radius of 750 km around cyclone centres in the two regions ~~underlined the leading role of the SST to explain~~, we demonstrate that the differences in the surface heat fluxes, specific humidity, ~~and~~ convective precipitation, ~~with cyclones generally playing a secondary modulating role~~ and upper-level wind speed, between the CNTL and SMTH experiments, largely arise in the absence of cyclones.

Code availability. The codes to construct the figures in this study is available upon request.

Data availability. ERA-Interim data are provided by European Centre for Medium-Range Weather Forecasts (ECMWF) available online at <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim> (Dee et al., 2011)

560 *Author contributions.* LT carried out the bulk of the data analysis and writing. TS contributed to detailed discussion about the methods and interpretation of the findings as well as to the writing process. CS contributed to both data analysis and writing.

Competing interests. The authors declare that they have no conflict of interest.

Appendix A: ERA-INTERIM CLIMATOLOGIES AND CYCLONES LOCATION




Figure A1. (a) DJF climatology over ERA-Interim period 1982-2000 for (a) the SSTs (K), (b) the SST gradient ($K (100 km)^{-1}$), (c) the latent (shading, $W m^{-2}$) and sensible heat fluxes (contours, $W m^{-2}$) for the North Atlantic. (d-f) As (a-c), but for the North Pacific. The Gulf Stream and Kuroshio regions are marked with a black box, respectively.




Figure A2. (a) DJF climatology over ERA-Interim period 1982-2000 for (a) ~~the~~ large-scale precipitation (shading, mm day^{-1}) and ~~the~~ convective precipitation (contours, mm day^{-1}), (b) ~~the~~ specific humidity at 850 hPa (g kg^{-1}) and (c) ~~the~~ wind speed at 300 hPa (m s^{-1}) for the North Atlantic. (d-f) As (a-c), but for the North Pacific. The Gulf Stream and Kuroshio regions are marked with a black box, respectively.




Figure A3. Locations of cyclones for C3 in the North Atlantic, (a) 12 hours prior to maximum intensification (blue crosses), (b) at the time of maximum intensification (dots colored depending on their pressure tendency, (hPa h^{-1})) and (c) 12 hours past maximum intensification (green crosses) in the CNTL experiment. (d-f) As (a-c), but for the SMTHG experiment. The Gulf Stream region is marked with a black box.




Figure A4. Locations of cyclones for C3 in the North Pacific, (a) 12 hours prior to maximum intensification (blue crosses), (b) at the time of maximum intensification (dots colored depending on their pressure tendency, hPa h^{-1}) and (c) 12 hours past-after maximum intensification (green crosses) in the CNTL experiment. (d-f) As (a-c), but for the SMTHK experiment. The Kuroshio region is marked with a black box.

Appendix B: CLIMATOLOGICAL DIFFERENCES FOR THE PERTINENT VARIABLES OCCURRING WITHIN/OUTSIDE OF DIFFERENT RADII OF ANY CYCLONE IN THE NORTH ATLANTIC AND THE NORTH PACIFIC

565




Figure B1. Difference (SMTHG-CNTL) of latent (shading) and sensible heat fluxes (grey contours, interval: 10 W m^{-2} , zero contour omitted) within a radius of 500 km, 750 km, and 1000 km (a,c,e, respectively) and outside of this radius (b,d,f, respectively) for the North Atlantic.




Figure B2. Difference (SMTHK-CNTL) of latent (shading) and sensible heat fluxes (grey contours, interval: 10 W m^{-2} , zero contour omitted) within a radius of 500 km, 750 km, and 1000 km (a,c,e, respectively) and outside of this radius (b,d,f, respectively) for the North Pacific.




Figure B3. Difference (SMTHG-CNTL) of large-scale precipitation (shading, mm day^{-1}) and convective precipitation (grey contours, interval: $0.5, \text{mm day}^{-1}$, zero contour omitted) within a radius of 500 km, 750 km, and 1000 km (a,c,e, respectively) and outside of this radius (b,d,f, respectively) for the North Atlantic.




Figure B4. Difference (SMTHK-CNTL) of large-scale precipitation (shading, mm day⁻¹) and convective precipitation (grey contours, interval: 0.5, mm day⁻¹, zero contour omitted) within a radius of 500 km, 750 km, and 1000 km (a,c,e, respectively) and outside of this radius (b,d,f, respectively) for the North Pacific.

Acknowledgements. We thank ECMWF for providing the ERA-Interim data and Akira Kuwano-Yoshida and Akira Yamazaki, who provided the AFES 3 data. [We also want to thank Irina Rudeva and an anonymous referee for their constructive feedback.](#) The AFES 3 data ~~was~~ [were](#) supported by the Grant-in-Aid for Scientific Research on Innovative Areas 22106008 ("A 'hot spot' in the climate system: Extra-tropical air-sea interaction under the East Asian monsoon system") from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and calculated using the Earth Simulator under JAMSTEC support. [The work was conducted as part of the UNPACC project, funded by the Research Council of Norway \(project number 262220\).](#)

References

- Alexander, M. A. and Scott, J. D.: Surface flux variability over the North Pacific and North Atlantic oceans, *Journal of climate*, 10, 2963–2978, 1997.
- 575 Berry, G., Reeder, M. J., and Jakob, C.: A global climatology of atmospheric fronts, *Geophys. Res. Lett.*, 38, L04 809, <https://doi.org/10.1029/2010GL046451>, 2011.
- Bjerknes, J.: Life cycle of cyclones and the polar front theory of atmospheric circulation, *Geophys. Publik.*, 3, 1–18, 1922.
- Booth, J. F., Thompson, L., Patoux, J., and Kelly, K. A.: Sensitivity of Midlatitude Storm Intensification to Perturbations in the Sea Surface Temperature near the Gulf Stream, *Mon. Wea. Rev.*, 140, 1241–1256, <https://doi.org/10.1175/MWR-D-11-00195.1>, 2012.
- 580 Brayshaw, D. J., Hoskins, B., and Blackburn, M.: The basic ingredients of the North Atlantic storm track. Part I: Land–sea contrast and orography., *J. Atmos. Sci.*, 66, 2539–2558, <https://doi.org/10.1175/2009JAS3078.1>, 2009.
- Bui, H. and Spengler, T.: On the Influence of Sea Surface Temperature distributions on the Development of Extratropical Cyclones, *Journal of the Atmospheric Sciences*, <https://doi.org/https://doi.org/10.1175/JAS-D-20-0137.1>, 2021.
- Chang, E. K. and Song, S.: The seasonal cycles in the distribution of precipitation around cyclones in the western North Pacific and Atlantic, *Journal of the atmospheric sciences*, 63, 815–839, 2006.
- 585 Chen, G., Plumb, R. A., and Lu, J.: Sensitivities of zonal mean atmospheric circulation to SST warming in an aqua-planet model, *Geophysical Research Letters*, 37, 2010.
- Dacre, H. F., Josey, S. A., and Grant, A. L.: Extratropical cyclone induced sea surface temperature anomalies in the 2013/14 winter, *Weather Clim. Dyn.*, 1, 27–44, <https://doi.org/10.5194/wcd-1-27-2020>, 2020.
- 590 de Vries, H., Scher, S., Haarsma, R., Drijfhout, S., and Van Delden, A.: How Gulf-Stream SST-fronts influence Atlantic winter storms, *Climate Dyn.*, 52, 5899–5909, <https://doi.org/10.1007/s00382-018-4486-7>, 2019.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Bergd, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kållberg, P., M. Köhler, M. M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quart. J. Roy. Meteor. Soc.*, 137, 553–587, <https://doi.org/10.1002/qj.828>, 2011.
- 595 Enomoto, T., Kuwano-Yoshida, A., Komori, N., and Ohfuchi, W.: Description of AFES 2: Improvements for high-resolution and coupled simulations, in: *High Resolution Numerical Modelling of the Atmosphere and Ocean*, pp. 77–97, Springer, 2008.
- Evans, M. S., Keyser, D., Bosart, L. F., and Lackmann, G. M.: A satellite-derived classification scheme for rapid maritime cyclogenesis, *Monthly weather review*, 122, 1381–1416, 1994.
- 600 Hanley, J. and Caballero, R.: The role of large-scale atmospheric flow and Rossby wave breaking in the evolution of extreme windstorms over Europe, *Geophys. Res. Lett.*, 39, L21 708, <https://doi.org/10.1029/2012GL053408>, 2012.
- Hualand, K. F. and Spengler, T.: Direct and Indirect Effects of Surface Fluxes on Moist Baroclinic Development in an Idealized Framework, *Journal of the Atmospheric Sciences*, 77, 3211–3225, <https://doi.org/https://doi.org/10.1175/JAS-D-19-0328.1>, 2020.
- 605 Hawcroft, M., Shaffrey, L., Hodges, K., and Dacre, H.: How much Northern Hemisphere precipitation is associated with extratropical cyclones?, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053866>, 2012.
- Holton, J. R. and Hakim, G. J.: *An Introduction to Dynamic Meteorology*, Academic Press, 2012.

- Hoskins, B. J. and Hodges, K. I.: New perspectives on the Northern Hemisphere winter storm tracks, *J. Atmos. Sci.*, 59, 1041–1061, [https://doi.org/10.1175/1520-0469\(2002\)059<1041:NPOTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPOTNH>2.0.CO;2), 2002.
- 610 Hoskins, B. J. and Valdes, P. J.: On the existence of storm-tracks, *Journal of the atmospheric sciences*, 47, 1854–1864, 1990.
- Hotta, D. and Nakamura, H.: On the significance of sensible heat supply from the ocean in the maintenance of mean baroclinicity along storm tracks, *J. Climate*, 24, 3377–3401, <https://doi.org/10.1175/2010JCLI3910.1>, 2011.
- Jacobs, N., Raman, S., Lackmann, G., and Childs Jr, P.: The influence of the Gulf Stream induced SST gradients on the US East Coast winter storm of 24–25 January 2000, *Int. J. Remote Sens.*, 29, 6145–6174, <https://doi.org/10.1080/01431160802175561>, 2008.
- 615 Jenkner, J., Sprenger, M., Schwenk, I., Schwierz, C., Dierer, S., and Leuenberger, D.: Detection and climatology of fronts in a high-resolution model reanalysis over the Alps, *Meteorol. Appl.*, 17, 1–18, <https://doi.org/10.1002/met.142>, 2010.
- Joyce, T. M., Kwon, Y.-O., and Yu, L.: On the relationship between synoptic wintertime atmospheric variability and path shifts in the Gulf Stream and the Kuroshio Extension, *Journal of Climate*, 22, 3177–3192, 2009.
- Kuwano-Yoshida, A. and Minobe, S.: Storm-track response to SST fronts in the northwestern Pacific region in an AGCM, *Journal of Climate*, 620 30, 1081–1102, 2017.
- Kuwano-Yoshida, A., Enomoto, T., and Ohfuchi, W.: An improved PDF cloud scheme for climate simulations, *Quarterly Journal of the Royal Meteorological Society*, 136, 1583–1597, 2010a.
- Kuwano-Yoshida, A., Minobe, S., and Xie, S.-P.: Precipitation response to the Gulf Stream in an atmospheric GCM, *Journal of Climate*, 23, 3676–3698, 2010b.
- 625 Lee, S. and Kim, H.-k.: The dynamical relationship between subtropical and eddy-driven jets, *Journal of the atmospheric sciences*, 60, 1490–1503, 2003.
- Li, C. and Wettstein, J. J.: Thermally driven and eddy-driven jet variability in reanalysis, *Journal of Climate*, 25, 1587–1596, 2012.
- Ma, X., Chang, P., Saravanan, R., Montuoro, R., Hsieh, J.-S., Wu, D., Lin, X., Wu, L., and Jing, Z.: Distant influence of Kuroshio eddies on North Pacific weather patterns?, *Scientific reports*, 5, 17 785, 2015.
- 630 Ma, X., Chang, P., Saravanan, R., Montuoro, R., Nakamura, H., Wu, D., Lin, X., and Wu, L.: Importance of resolving Kuroshio front and eddy influence in simulating the North Pacific storm track, *Journal of Climate*, 30, 1861–1880, 2017.
- Masunaga, R., Nakamura, H., Miyasaka, T., Nishii, K., and Tanimoto, Y.: Separation of climatological imprints of the Kuroshio Extension and Oyashio fronts on the wintertime atmospheric boundary layer: Their sensitivity to SST resolution prescribed for atmospheric reanalysis, *Journal of Climate*, 28, 1764–1787, 2015.
- 635 Masunaga, R., Nakamura, H., Taguchi, B., and Miyasaka, T.: Processes shaping the frontal-scale time-mean surface wind convergence patterns around the Gulf Stream and Agulhas Return Current in winter, *Journal of Climate*, 33, 9083–9101, 2020a.
- Masunaga, R., Nakamura, H., Taguchi, B., and Miyasaka, T.: Processes shaping the frontal-scale time-mean surface wind convergence patterns around the Kuroshio Extension in winter, *Journal of Climate*, 33, 3–25, 2020b.
- Meinen, C. S. and Luther, D. S.: Structure, transport, and vertical coherence of the Gulf Stream from the Straits of Florida to the Southeast 640 Newfoundland Ridge, *Deep-Sea Res. Pt. I*, 112, 137–154, <https://doi.org/10.1016/j.dsr.2016.03.002>, 2016.
- Michel, C., Terpstra, A., and Spengler, T.: Polar mesoscale cyclone climatology for the Nordic Seas based on ERA-Interim, *J. Climate*, 31, 2511–2532, <https://doi.org/10.1175/JCLI-D-16-0890.1>, 2018.
- Minobe, S., Kuwano-Yoshida, A., Komori, N., Xie, S.-P., and Small, R. J.: Influence of the Gulf Stream on the troposphere, *Nature*, 452, 206–210, <https://doi.org/10.1038/nature06690>, 2008.

- 645 Murray, R. J. and Simmonds, I.: A numerical scheme for tracking cyclone centres from digital data. Part I: Development and operation of the scheme, *Aust. Met. Mag.*, 39, 155–166, 1991a.
- Murray, R. J. and Simmonds, I.: A numerical scheme for tracking cyclone centres from digital data. Part II: application to January and July general circulation model simulations, *Aust. Met. Mag.*, 39, 167–180, 1991b.
- Nakamura, H.: Midwinter suppression of baroclinic wave activity in the Pacific, *Journal of the atmospheric sciences*, 49, 1629–1642, 1992.
- 650 Nakamura, H., Sampe, T., Tanimoto, Y., and Shimpo, A.: Observed associations among storm tracks, jet streams and midlatitude oceanic fronts, *Geoph. Monog. Series*, 147, 329–345, <https://doi.org/10.1029/147GM18>, 2004.
- Nakamura, H., Sampe, T., Goto, A., Ohfuchi, W., and Xie, S.-P.: On the importance of midlatitude oceanic frontal zones for the mean state and dominant variability in the tropospheric circulation, *Geophysical Research Letters*, 35, 2008.
- Neu, U., Akperov, M. G., Bellenbaum, N., Benestad, R., Blender, R., Caballero, R., Cocozza, A., Dacre, H. F., Feng, Y., Fraedrich, K., et al.:
- 655 IMILAST: A community effort to intercompare extratropical cyclone detection and tracking algorithms, *Bull. Amer. Meteor. Soc.*, 94, 529–547, <https://doi.org/10.1175/BAMS-D-11-00154.1>, 2013.
- Newman, M., Kiladis, G. N., Weickmann, K. M., Ralph, F. M., and Sardeshmukh, P. D.: Relative contributions of synoptic and low-frequency eddies to time-mean atmospheric moisture transport, including the role of atmospheric rivers, *Journal of climate*, 25, 7341–7361, 2012.
- Ogawa, F. and Spengler, T.: Prevailing Surface Wind Direction during Air–Sea Heat Exchange, *J. Climate*, 32, 5601–5617,
- 660 <https://doi.org/10.1175/JCLI-D-18-0752.1>, 2019.
- Ogawa, F., Nakamura, H., Nishii, K., Miyasaka, T., and Kuwano-Yoshida, A.: Dependence of the climatological axial latitudes of the tropospheric westerlies and storm tracks on the latitude of an extratropical oceanic front, *Geophysical research letters*, 39, 2012.
- Ohfuchi, W., Nakamura, H., Yoshioka, M. K., Enomoto, T., Takaya, K., Peng, X., Yamane, S., Nishimura, T., Kurihara, Y., Ninomiya, K., et al.: 10-km mesh meso-scale resolving simulations of the global atmosphere on the Earth Simulator: Preliminary outcomes of AFES
- 665 (AGCM for the Earth Simulator), *J. Earth Simulator*, 1, 8–34, 2004.
- O’Reilly, C. H., Minobe, S., Kuwano-Yoshida, A., and Woollings, T.: The Gulf Stream influence on wintertime North Atlantic jet variability, *Quarterly Journal of the Royal Meteorological Society*, 143, 173–183, 2017.
- O’Neill, L. W., Haack, T., Chelton, D. B., and Skillingstad, E.: The Gulf Stream convergence zone in the time-mean winds, *Journal of the Atmospheric Sciences*, 74, 2383–2412, 2017.
- 670 Papritz, L. and Spengler, T.: Analysis of the slope of isentropic surfaces and its tendencies over the North Atlantic, *Quart. J. Roy. Meteor. Soc.*, 141, 3226–3238, <https://doi.org/10.1002/qj.260>, 2015.
- Parfitt, R. and Seo, H.: A new framework for near-surface wind convergence over the Kuroshio extension and Gulf Stream in wintertime: The role of atmospheric fronts, *Geophysical Research Letters*, 45, 9909–9918, 2018.
- Parfitt, R., Czaja, A., Minobe, S., and Kuwano-Yoshida, A.: The atmospheric frontal response to SST perturbations in the Gulf Stream region,
- 675 *Geophys. Res. Lett.*, 43, 2299–2306, <https://doi.org/10.1002/2016GL067723>, 2016.
- Parfitt, R., Czaja, A., and Kwon, Y.-O.: The impact of SST resolution change in the ERA-Interim reanalysis on wintertime Gulf Stream frontal air-sea interaction, *Geophysical Research Letters*, 44, 3246–3254, 2017.
- Peixoto, J. P. and Oort, A. H.: *Physics of climate*, 1992.
- Pfahl, S. and Wernli, H.: Quantifying the relevance of cyclones for precipitation extremes, *Journal of Climate*, 25, 6770–6780, 2012.
- 680 Piazza, M., Terray, L., Boé, J., Maisonnave, E., and Sanchez-Gomez, E.: Influence of small-scale North Atlantic sea surface temperature patterns on the marine boundary layer and free troposphere: A study using the atmospheric ARPEGE model, *Climate dynamics*, 46, 1699–1717, 2016.

- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily high-resolution-blended analyses for sea surface temperature, *Journal of Climate*, 20, 5473–5496, 2007.
- 685 Riviere, G. and Joly, A.: Role of the low-frequency deformation field on the explosive growth of extratropical cyclones at the jet exit. Part II: Baroclinic critical region, *Journal of the atmospheric sciences*, 63, 1982–1995, 2006.
- Rudeva, I. and Gulev, S. K.: Composite analysis of North Atlantic extratropical cyclones in NCEP–NCAR reanalysis data, *Mon. Wea. Rev.*, 139, 1419–1446, <https://doi.org/10.1175/2010MWR3294.1>, 2011.
- Ruprecht, E., Schröder, S. S., and Ubl, S.: On the relation between NAO and water vapour transport towards Europe, *Meteorologische Zeitschrift*, 11, 395–401, 2002.
- 690 Sampe, T., Nakamura, H., Goto, A., and Ohfuchi, W.: Significance of a midlatitude SST frontal zone in the formation of a storm track and an eddy-driven westerly jet, *Journal of Climate*, 23, 1793–1814, 2010.
- Sanders, F.: Explosive cyclogenesis in the west-central North Atlantic Ocean, 1981–84. Part I: Composite structure and mean behavior, *Mon. Wea. Rev.*, 114, 1781–1794, [https://doi.org/10.1175/1520-0493\(1986\)114<1781:ECITWC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1986)114<1781:ECITWC>2.0.CO;2), 1986.
- 695 Sanders, F. and Gyakum, J. R.: Synoptic-dynamic climatology of the “bomb”, *Mon. Wea. Rev.*, 108, 1589–1606, [https://doi.org/10.1175/1520-0493\(1980\)108<1589:SDCOT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<1589:SDCOT>2.0.CO;2), 1980.
- Schemm, S., Rudeva, I., and Simmonds, I.: Extratropical fronts in the lower troposphere—global perspectives obtained from two automated methods, *Quart. J. Roy. Meteor. Soc.*, 141, 1686–1698, <https://doi.org/10.1002/qj.2471>, 2015.
- Schultz, D. M., Keyser, D., and Bosart, L. F.: The effect of large-scale flow on low-level frontal structure and evolution in midlatitude cyclones, *Monthly weather review*, 126, 1767–1791, 1998.
- 700 Sinclair, M. R. and Revell, M. J.: Classification and composite diagnosis of extratropical cyclogenesis events in the southwest Pacific, *Mon. Wea. Rev.*, 128, 1089–1105, [https://doi.org/10.1175/1520-0493\(2000\)128<1089:CACDOE>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<1089:CACDOE>2.0.CO;2), 2000.
- Small, R. d., deSzoeko, S. P., Xie, S., O’neill, L., Seo, H., Song, Q., Cornillon, P., Spall, M., and Minobe, S.: Air–sea interaction over ocean fronts and eddies, *Dynamics of Atmospheres and Oceans*, 45, 274–319, 2008.
- 705 Small, R. J., Tomas, R. A., and Bryan, F. O.: Storm track response to ocean fronts in a global high-resolution climate model, *Climate dynamics*, 43, 805–828, 2014.
- Spensberger, C. and Spengler, T.: Feature-based Jet Variability in the Upper Troposphere, *Journal of Climate*, p. submitted, 2020.
- Spensberger, C., Spengler, T., and Li, C.: Upper-tropospheric jet axis detection and application to the boreal winter 2013/14, *Monthly Weather Review*, 145, 2363–2374, 2017.
- 710 Tanimoto, Y., Nakamura, H., Kagimoto, T., and Yamane, S.: An active role of extratropical sea surface temperature anomalies in determining anomalous turbulent heat flux, *Journal of Geophysical Research: Oceans*, 108, 2003.
- Tozuka, T., Ohishi, S., and Cronin, M. F.: A metric for surface heat flux effect on horizontal sea surface temperature gradients, *Climate Dynamics*, 51, 547–561, 2018.
- Tsopouridis, L., Spensberger, C., and Spengler, T.: Characteristics of Cyclones Following Different Pathways in the Gulf Stream Region, *Quarterly Journal of the Royal Meteorological Society*;1-16, <https://doi.org/10.1002/qj.3924>, 2020a.
- 715 Tsopouridis, L., Spensberger, C., and Spengler, T.: Cyclone Intensification in the Kuroshio Region and its relation to the Sea Surface Temperature Front and Upper-Level Forcing, *Quarterly Journal of the Royal Meteorological Society*, <https://doi.org/10.1002/qj.3929>, 2020b.
- Uccellini, L. W., Kocin, P. J., Petersen, R. A., Wash, C. H., and Brill, K. F.: The Presidents’ Day cyclone of 18–19 February 1979: Synoptic overview and analysis of the subtropical jet streak influencing the pre-cyclogenetic period, *Monthly weather review*, 112, 31–55, 1984.

- 720 Vanniere, B., Czaja, A., Dacre, H. F., and Woollings, T.: A “Cold Path” for the Gulf Stream–Troposphere Connection, *J. Climate*, 30, 1363–1379, <https://doi.org/10.1175/JCLI-D-15-0749.1>, 2017.
- Wang, C.-C. and Rogers, J. C.: A composite study of explosive cyclogenesis in different sectors of the North Atlantic. Part I: Cyclone structure and evolution, *Mon. Wea. Rev.*, 129, 1481–1499, [https://doi.org/10.1175/1520-0493\(2001\)129<1481:ACSOEC>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<1481:ACSOEC>2.0.CO;2), 2001.
- Wang, L., Hu, H., and Yang, X.: The atmospheric responses to the intensity variability of subtropical front in the wintertime North Pacific, *Climate Dynamics*, 52, 5623–5639, 2019.
- 725 Wernli, H. and Schwierz, C.: Surface cyclones in the ERA-40 dataset (1958–2001). Part I: Novel identification method and global climatology, *Journal of the atmospheric sciences*, 63, 2486–2507, 2006.
- Yao, Y., Zhong, Z., and Yang, X.-Q.: Impacts of the subarctic frontal zone on the North Pacific storm track in the cold season: an observational study, *International Journal of Climatology*, 38, 2554–2564, 2018.
- 730 Yoshida, A. and Asuma, Y.: Structures and environment of explosively developing extratropical cyclones in the northwestern Pacific region, *Monthly Weather Review*, 132, 1121–1142, 2004.
- Zhang, C., Liu, H., Xie, J., Lin, P., Li, C., Yang, Q., and Song, J.: North Pacific storm track response to the mesoscale SST in a global high-resolution atmospheric model, *Climate Dynamics*, 55, 1597–1611, 2020.
- Zolina, O. and Gulev, S. K.: Synoptic variability of ocean–atmosphere turbulent fluxes associated with atmospheric cyclones, *Journal of*
735 *climate*, 16, 2717–2734, 2003.