

# Impact of climate change on wintertime European atmospheric blocking

Sara Bacer<sup>1</sup>, Fatima Jomaa<sup>1</sup>, Julien Beaumet<sup>2</sup>, Hubert Gallée<sup>2</sup>, Enzo Le Bouëdec<sup>1</sup>, Martin Ménégoz<sup>2</sup>, and Chantal Staquet<sup>1</sup>

<sup>1</sup>Univ. Grenoble Alpes, CNRS, Grenoble INP, LEGI, 38000 Grenoble, France

<sup>2</sup>Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

**Correspondence:** Sara Bacer (sara.bacer@univ-grenoble-alpes.fr)

**Abstract.** We study the impact of climate change on wintertime atmospheric blocking over Europe focusing on the frequency, duration, and size of blocking events. These events are identified via the weather type decomposition (WTD) methodology applied on the output of climate models of the Coupled Model Intercomparison Project phase 6 (CMIP6). Historical simulations as well as two future scenarios, SSP2-4.5 and SSP5-8.5, are considered. The models are evaluated against the reanalysis and

5 only a subset of climate models, which better represent the blocking weather regime in the recent-past climate, is considered for the analysis. We show that the spatio-temporal characteristics of recent-past atmospheric blocking are in agreement with previous studies that define blocking events with blocking indexes. We find that frequency and duration of blocking events remain relatively stationary over the 21st century. We define a methodology that relies on the WTD for the blocking event identification in order to quantify the size of the blocking events, and we find that the blocking size is basically unchanged in

10 the future. We show that our results are in agreement with previous studies that define blocking events with blocking indexes.

## 1 Introduction

Atmospheric blocking is a persistent and quasi-stationary phenomenon which highly impacts the mid-latitude circulation. By obstructing the usual westerly winds, atmospheric blocking can promote cut-off cyclones (Munoz et al., 2020) and enhance cooling in winter and warming in summer. Its long duration (from days to weeks) affects surface weather and climate and fosters

15 regional extreme events, such as heatwaves, droughts, and severe cold weather in winter (Barriopedro et al., 2010; Woollings et al., 2018, and references therein). Blocking events are generally associated with high-pressure systems. During anticyclonic periods, solar radiation and high temperatures in summer promote ozone formation, while thermal inversions with subsidence conditions in winter promote the accumulation of particulate matter (e.g. Largeron and Staquet, 2016; Hou and Wu, 2016).

20 Simulating blocking is a challenging task for atmospheric models as it requires an accurate description of the topography, a fine resolution both vertically and horizontally, appropriate physical parameterizations, and a correct description of internal dynamics (Davini et al., 2017). It has been shown that general circulation models (GCMs) are able to reproduce the blocking regime and its variability, although they tend to underestimate frequency and persistence of blocking events (Dunn-Sigouin

et al., 2013; Masato et al., 2013, 2014; Woollings et al., 2018; Davini and D'Andrea, 2020). Increasing model resolution can 25 improve the blocking occurrence, as the transient eddies and orography are better described (Berckmans et al., 2013; Schiemann et al., 2017). Since atmospheric blocking is related to stratospheric variability (e.g. the stratospheric sudden warming, Davini et al., 2014), a good representation of the stratosphere can also improve blocking simulations.

The identification of blocking events in numerical simulations is complicated by the fact that blocking is determined by 30 various dynamical mechanisms and presents different patterns. Several blocking indexes have been proposed in the literature, based on meteorological fields, usually the geopotential height at 500 hPa (e.g. Tibaldi and Molteni, 1990), or anomalies of meteorological fields (e.g. Dole and Gordon, 1983). Blocking indexes focus on different characteristics of blocking, so the choice of the index depends on the purpose of the study. Additionally, index definitions depend on various (user-dependent) 35 parameters, like latitude band limits, latitude references, and anomaly thresholds (a review of the blocking indexes can be found in Barriopedro et al. (2010), while a recent discussion about their differences is in Pinheiro et al. (2019)). Given the variety of blocking indexes, the comparison across studies is not straightforward.

Atmospheric blocking can also be identified via the so-called weather type decomposition (WTD) methodology, which classifies the atmospheric circulation into discrete weather regimes (Michelangeli et al., 1995). The WTD methodology, referred to as *the WTD* hereafter for brevity, relies on a partitioning algorithm that groups data of a meteorological variable (usually geopotential height or sea level pressure) into clusters so that the variance between clusters is maximized and the variance 40 within a given cluster is minimized. In this way, the clusters (weather regimes or weather types) are the result of a mathematical algorithm. The results of the WTD depend on certain user choices, such as the sector size, the clustering algorithm, and the initialization of this algorithm. Despite the fact that the clusters may not be well separated, *the WTD* has proved to be very useful in the literature. In fact, ~~WTD allows the WTD can be used~~ to explain most of the atmospheric variability and has largely been used to define weather regimes especially in the Northern Hemisphere (e.g. Michelangeli et al., 1995; Cassou et al., 2004; 45 Barriopedro et al., 2006; Ullmann et al., 2014; Fabiano et al., 2020). In the European-Atlantic sector, for example, four winter weather types have been recognized: positive North Atlantic Oscillation (NAO), negative NAO, Atlantic ridge, and European blocking. The WTD has also been used to ~~analyze analyse~~ weather types in relation to other quantities like temperature (e.g. Cassou et al., 2005), ~~precipitations precipitation~~ (e.g. Ullmann et al., 2014), winds (e.g. Jiménez et al., 2009), and pollutants (e.g. Russo et al., 2014). In this study, the WTD is used to identify blocking events in the European-Atlantic sector.

50 The impact of blocking events ~~on weather and climate~~ is related to their spatio-temporal characteristics, such as occurrence, duration, and size. Many studies investigated frequency and duration of blocking events in the past climate using reanalysis data (e.g. Wiedenmann et al., 2002; Barriopedro et al., 2006; Mokhov et al., 2013; Cheung et al., 2013; Drouard and Woollings, 2018; Lupo et al., 2019). Understanding the impact of climate change on atmospheric blocking is of fundamental importance to estimate future climate and extreme events, thus, blocking has also been investigated in the future in response to global 55 warming. For example, ~~it has been shown the Arctic amplification has been studied in relation to the intensity and frequency of blocking events; although some studies suggest that the Arctic amplification, which has a strong influence on mid-latitude atmospheric circulation, modulates the frequency and the intensity of blocking events (e.g. Hassanzadeh et al., 2014). Climate models suggest warming yields to an increment of these two quantities (e.g. Francis and Vavrus, 2012), further investigations~~

are necessary to define the Arctic amplification response to blocking (Hassanzadeh et al., 2014; Barnes et al., 2014). So far, 60 studies have mainly focused on frequency and duration of future blocking events. Some of these studies found that blocking frequency may will decrease in the Northern Hemisphere in the future (e.g. Dunn-Sigouin and Son, 2013; Fabiano et al., 2020), and blocking activity could shift eastwards (e.g. Masato et al., 2013, 2014; Woollings et al., 2018), while there is no clear tendency for changes in blocking duration.

So far, studies have mainly focused on (e.g. Dunn-Sigouin and Son, 2013; Matsueda and Endo, 2017; Fabiano et al., 2020; Davini and D'Andrea, 2020), while blocking duration may either increase (Sillmann and Croci-Maspoli, 2009) or decrease (Fabiano et al., 2020). Other 65 studies showed that blocking frequency and duration changes of future blocking events (e.g. Barriopedro et al., 2006; Patterson et al., 2019; Huguenin et al., 2020) will not change notably in warming climate (Dunn-Sigouin and Son, 2013; Huguenin et al., 2020). Future changes in blocking size have received less attention (Nabizadeh et al., 2019). These works (Hassanzadeh et al., 2014; Nabizadeh et al., 2019).

Most of the studies mentioned above determined blocking events via blocking indexes and considered one or more (Sillmann and Croci-Maspoli, 2009; Hassanzadeh et al., 2014) GCMs participating in the Coupled Model Intercomparison Project phase 5 (CMIP5) (CMIP5, Dunn-Sigouin and Son, 2013) or idealized GCMs (Hassanzadeh et al., 2014). To our knowledge, only Fabiano et al. (2020) employed CMIP6. Fabiano et al. (2020) 70 applied the WTD on CMIP6 models in order to project future weather types and analyse their analyse the changes in frequency and duration of the blocking weather type during the 21st century.

In this study, we investigate the impact of climate change on European atmospheric blocking in terms of frequency, duration, 75 and especially size. Several GCMs of the latest model intercomparison CMIP6 are considered for this purpose under two different future scenarios (SSP2-4.5 and SSP5-8.5). In order to identify blocking events, the WTD is applied. We focus on wintertime blocking as it is more frequent, longer, and stronger than blocking in summer in the European-Atlantic sector (Barriopedro et al., 2006; Cheung et al., 2013; Lupo et al., 2019). Moreover, winter blocking events are often associated with severe particulate matter pollution episodes. We define a method, referred to as the composite WTD method, to quantify the size 80 of blocking events that are identified via the WTD. We compare the results obtained with this method with the results obtained for the blocking events identified via the index of Dole and Gordon (1983). Besides using GCMs of the latest CMIP phase, investigating frequency, duration, and size of blocking events that are determined via the WTD instead of blocking indexes makes this work an original study.

## 2 Data

85 Daily means of geopotential height at 500 hPa (Z500) are used for the WTD. More precisely, the WTD is applied on winter anomalies of Z500, where the winter season is defined from 1 November to 31 March (NDJFM, like in Cassou (2008), for instance). The numerical domain of Z500 covers the European-Atlantic sector whose boundaries are 80°W, 50°E, 20°N, and 80°N.

In this study, GCMs of the CMIP6 (Eyring et al., 2016) are considered. It has been shown that the weather regimes are 90 reproduced better in CMIP6 models than in CMIP5 models, especially over the European-Atlantic sector (Fabiano et al., 2020; Davini and D'Andrea, 2020). We use historical runs to analyse blocking conditions in recent-past climate and two future

Climate center	GCM	Acronym	Lon x Lat
Beijing Climate Center (China)	BCC-CSM2-MR	BCC	1.1°x1.1°
Canadian Centre for Climate Modelling and Analysis (Canada)	CanESM5	CanESM	2.8°x2.8°
Institute of Atmospheric Physics (China)	FGOALS-g3	FGOALS	2.0°x2.0°
National Oceanic and Atmospheric Administration (USA)	GFDL-CM4	GFDL	2.5°x2.0°
Institute of Numerical Mathematics (Russia)	INM-CM5-0	INM	2.0°x1.5°
Institut Pierre-Simon Laplace (France)	IPSL-CM6A-LR	IPSL	2.5°x1.3°
Atmospheric and Ocean Research Institute (Japan)	MIROC6	MIROC	1.4°x1.4°
Max Planck Institute for Meteorology (Germany)	MPI-ESM1-2-HR	MPI	0.9°x0.9°
Meteorological Research Institute (Japan)	MRI-ESM2-0	MRI	1.1°x1.1°

**Table 1.** The CMIP6-GCMs used in this study. The columns contain, respectively, the name of the research center developer of the GCM, the name of the GCM, the acronym used in this study, and the resolution of the Z500 output. All data were provided by the Mésocentre ESPRI.

projections, SSP2-4.5 and SSP5-8.5 (Riahi et al., 2017), to investigate their changes in future climate. SSP2-4.5 assumes that social, economic, and technological trends broadly follow their historical patterns and is considered as a likely scenario given the current policies. In contrast, SSP5-8.5 projects strong increments of emissions without mitigation policies; it is the worst-case scenario and is considered unlikely (Hausfather and Peters, 2020). We also use the ERA5 reanalysis of the European Centre for Medium-Range Weather Forecasts with a resolution of 31 km (Hersbach et al., 2020) to evaluate the GCM ability in reproducing the blocking weather regime.

The Z500 outputs considered in this study are archived in the Mésocentre ESPRI. We selected the nine CMIP6-GCMs presented in Table 1 according to the following criteria: one GCM per each climate research centre, as different versions of the same model could present model-dependent similarities (Ullmann et al., 2014); GCMs having both SSP2-4.5 and SSP5-8.5 scenarios available; GCMs with the “r1i1p1f1” run available (where “r1”: initial conditions, “i1”: initialization method, “p1”: physical scheme, and “f1”: forcing configuration), as this is the most frequently accessible simulation. The analysed periods are 30-year long: 1980-2009 (HIST hereafter) and 2070-2099 (SSP2 or SSP5 hereafter, according to the scenario).

### 3 Methods

#### 105 3.1 Detection of the blocking weather regime

The following procedure is carried out for each GCM of Table 1 and ERA5 and for each period (HIST, SSP2, and SSP5). First, daily anomalies of Z500, noted  $\Delta Z500$ , are computed as difference between the daily means of Z500 and the annual cycle (including the mean) of the 30-year period used as a climatology reference (see subsection 3.2); only the winter season (NDJFM) is retained. Second, the anomalies are weighted (multiplied) by the square root of the cosine of the latitude (Chung and Nigam, 1999) in order to account for the convergence of the meridians and to decrease the impact of high-latitude grid

boxes that represent a small area of the globe (like in Cassou, 2008; Ullmann et al., 2014; Cortesi et al., 2019). Since the GCMs have different resolutions (Table 1), the anomalies are linearly interpolated onto a common grid of resolution  $1^\circ \times 1^\circ$ .

Blocking events are identified through the application of the WTD. This weather classification has largely been used in order to infer the recurrent atmospheric features at mid-latitudes (Michelangeli et al., 1995; Philipp et al., 2016). It can be divided into 115 two steps: dimensional reduction of the data set and clustering. Similarly to other studies (e.g. Boé and Terray, 2008; Hertig and Jacobbeit, 2014; Sáenz and Durán-Quesada, 2015), we apply the Principal Component Analysis (PCA) for the first step and the k-means algorithm for the second step. Therefore, the PCA is applied to the resulting anomalies, and the eigenvectors necessary to explain 95% of the total variance (24 eigenvectors on average) are retained to define the reduced data set. Then, 120 k-means is applied to this data set by imposing that the number of clusters (k) is equal to four, i.e. the four well-known weather types of the European-Atlantic sector (positive and negative NAO, Atlantic ridge, and European blocking), as done in Cassou (2008), Ullmann et al. (2014), and Fabiano et al. (2020), and each day of HIST, SSP2, and SSP5 is assigned to one of the four weather types. Only the weather regime corresponding to the European atmospheric blocking is analysed in this study. On the whole, while identifying blocking via blocking indexes implies making several choices, identifying blocking via the WTD can be considered as a standard procedure. This motivated us to apply the WTD and to explore this methodology for identifying 125 blocking events and then studying their main characteristics (frequency, duration, size).

### 3.2 Z500 anomalies

Climate change causes an overall increase of Z500 due to the warming of air masses. In order to study the changes of the spatio-temporal characteristics of blocking in the future, we compute the anomalies in all periods (HIST, SSP2, SSP5) by subtracting from Z500 the annual cycle of that period; these anomalies are noted  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$ . 130 Being the blocking events identified with the departure (anomaly) from the atmospheric mean state, the comparison between recent-past and future results will allow to quantify the dynamical climate signal ignoring the thermodynamical signal related to the anthropogenic warming. ~~Additionally, we~~ We also compute the future anomalies by subtracting from Z500 of the future periods the HIST annual cycle; these anomalies are noted  $\Delta Z500_{SSP2-HIST}$  and  $\Delta Z500_{SSP5-HIST}$ . In this case, the comparison between recent-past and future anomalies will show the ~~net~~ gross impact on blocking of the total climate change signal, 135 governed by greenhouse gases increase, global warming, and associated regional circulation changes.

### 3.3 Definition of blocking events

Consecutive days that belong to the blocking weather type can form a blocking event. The blocking events considered in this study ~~satisfy the following conditions: they must be longer than five days (like in Barriopedro et al., 2006; Matsueda et al., 2009; Mokhov et al., 2009) and separated by at least two non-blocking days. They must be at least five days long (like in Barriopedro et al., 2006; Matsueda et al., 2009; Mokhov et al., 2009)~~. A single non-blocking day (“hole”) ~~is assumed to~~ might represent a failure of k-means in that day, like in Matsueda et al. (2009). ~~Therefore~~ Concretely, the k-means result is processed in such a way that 1) two blocking events equal to/longer than two days separated by a hole form one blocking event and 2) one blocking event equal to/longer than three days and one block-

ing day separated by a hole (and then vice versa) form one blocking event. Overall, the number of holes that are converted into blocking days is very small, about 0.3% of the number of winter days (4530).

145 We clarify here the meaning of some specific terms. We call *centroids* the four centres of mass defined by the k-means algorithm in the reduced space, i.e. the space whose coordinates are the eigenvectors. Weather regime (or weather type) refers to the centroid transformed into the original latitude-longitude coordinate space. From now on, we call *blocking days* those days which belong to a blocking event. Finally, we refer to the temporal mean of  $\Delta Z500$  over the blocking days of a blocking event as the *composite* of that event.


150 **3.4 Computation of blocking area**

We quantify the size of a blocking event by its area. Two distinct methods are used to compute the blocking area: the so-called composite-WTD method, introduced in this study, and the *DG method*, used by Nabizadeh et al. (2019).

155 **Composite-WTD method.** We introduce this method to compute the area of the composites of the blocking events inferred from the WTD. The composite-WTD method starts from the detection of the center of each composite. In this study, we define as *center* of the European atmospheric blocking the location of the maximum positive anomaly of the composite between 30°W and 50°E (similarly to Barriopedro et al., 2006), in order to discard blocking with positive anomaly on the westernmost part of the sector. The blocking size is quantified by the area enclosed within the contour line equal to a certain threshold. In order to get non-zero areas, we define a threshold of  $\Delta Z500$  that Such threshold must be lower than the minimum value among the centers over all periods and all GCMs in order to get non-zero areas and compare past and future results among the GCMs. In 160 this study, the threshold is 75 m considering  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$ , and  $\Delta Z500_{SSP5}$  ~~and 100 m considering the  $\Delta Z500$  with respect to the recent-past climatology (i.e.  $\Delta Z500_{SSP2-HIST}$  and  $\Delta Z500_{SSP5-HIST}$ )~~. ~~For the purpose of this study, which requires a comparison between past and future results among several GCMs, we keep the thresholds constant for the entire analysis.~~ Technical details about the composite-WTD method are reported in the Supplement.

165 **DG method.** This method follows the work of Nabizadeh et al. (2019), who determined the size of the atmospheric blocking events identified with the index of Dole and Gordon (1983) (DG index hereafter). Thus, we also compute the daily DG index for each grid box of the domain and identify as blocking events those grid boxes where the DG index is higher than 1.5 for at least five consecutive days. We will refer to these days as *DG-blocking days* and to the blocking events as *DG-blocking events*. The DG method consists in computing, for each DG-blocking day, the area enclosed by the contour line where the DG index is equal to one, i.e. the contour line equal to a certain threshold of  $\Delta Z500$ : 112 m for  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$ , and  $\Delta Z500_{SSP5}$  ~~and 116 m for the  $\Delta Z500$  with respect to the recent-past climatology~~. Then, daily areas are averaged along the event duration to get the area of the DG-blocking event (more details about the DG method are in the Supplement). In the present paper, the DG-blocking days are identified within the blocking events inferred from the WTD (and not during the entire winter, as considered by Dole and Gordon (1983)).

175 Therefore, these methods compute the area of blocking events that are identified via two different approaches: the WTD and the DG index. Although the algorithm to compute the blocking area within a certain contour line is the same, it is applied on



**Figure 1.** Taylor diagram for the mean composites over all blocking events for ERA5 and all GCMs for the winter HIST period (1980-2009). The diagram allows to quantify standard deviation (black), correlation coefficient (light blue), and root-mean-square difference (green) between the mean GCM composites and the mean ERA5 composite. SSP2 and SSP5 results are obtained with  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$ , respectively.

blocking composites in the composite WTD method and on daily  $\Delta Z500$  in the DG method. Another difference between these two methods is the definition of the  $\Delta Z500$  values of the contour lines (i.e. the thresholds).

## 4 Results and Discussion

### 4.1 Evaluation of the GCMs

180 Before analysing the impact of climate change on European atmospheric blocking events, the ability of the GCMs (Table 1) in reproducing atmospheric blocking is evaluated with respect to the reanalysis with a Taylor diagram (Figure 1). This diagram compares the blocking composites of each GCM during HIST with the ERA5 composite. The deviation is quantified in terms of pattern correlation ( $R$ ), standard deviation ( $\sigma$ ), and root-mean-square difference (RMSD). All GCMs are able to represent blocking variability (i.e.  $\sigma$ ) quite close to the variability obtained with the reanalysis ( $\sigma_{ERA5} \approx 61$  m). More precisely, the 185 variability of all GCMs is within the range  $\sigma_{ERA5} \pm 6$  m, apart from INM and IPSL. Six models (MPI, BCC, MRI, GFDL, INM, and FGOALS) show a high correlation ( $R \geq 0.79$ ) with ERA5, while three models (CanESM, MIROC, and IPSL) present

a lower correlation ( $R < 0.6$ ) and a high RMSD. Hertig and Jacobbeit (2014) also found that historical runs of CanESM cannot well reproduce the blocking pattern, getting a correlation with reanalysis lower than 0.4. In this study, CanESM is the GCM with the coarsest resolution (Table 1), and it has been shown that a low resolution hinders a good description of the atmospheric variability patterns (Berckmans et al., 2013).


This analysis points out that MIROC, CanESM, and IPSL are less accurate in capturing the blocking pattern in recent-past climate (as also observable in Figure 2, e-f-g), and we expect these models to be less reliable in future projections of blocking. Previous studies (e.g. Chhin and Yoden, 2018; Mokhov and Timazhev, 2019; Khan et al., 2020) suggest to use a subset of GCMs selected according to their ability in simulating the quantity of interest (atmospheric blocking in this study) in the past in order to reduce the uncertainties associated to the future projections of that quantity. Therefore, we exclude MIROC, CanESM, and IPSL from the next analysis and focus on the results obtained by the other six GCMs: MPI, BCC, MRI, GFDL, INM, and FGOALS. In the same Taylor diagram, blocking projected ~~for in~~ future climates (both ~~SPP2 and SSP5~~<sup>SPP2-4.5 and SSP5-8.5</sup>) by these six GCMs is also shown. Overall, correlation coefficients, standard deviations, and RMSDs vary in a non-systematic way, so we do not find any regularity in the reproducibility of future blocking by the GCMs.

The spatial patterns of blocking during recent-past climate are shown in Figure 2 (a-i). All GCMs are considered during HIST, and the dissimilarity of CanESM, MIROC, and IPSL with respect to ERA5 (Figure 2, k) is evident. According to the reanalysis, the European blocking is centred over the Scandinavian peninsula and extends over northern Europe. Blocking occurrence is about 27% (Table S1 in the Supplement) in accordance with previous studies that considered, for example, NCEP/NCAR reanalysis (27%, Cassou, 2008) and ERA-interim reanalysis (26%, Ullmann et al., 2014). MPI, BCC, and MRI reproduce an occurrence similar to ERA5, while GFDL, INM, and FGOALS simulate less frequent blocking with an occurrence of about 23%. We observe that the first three models have the highest resolution (see Table 1), so we also find that the underestimation of atmospheric blocking occurrence is reduced in higher resolution GCMs. We compute the multi-model (MM) mean as the average of the composites over all blocking events of the six selected GCMs. The spatial pattern of the MM mean during HIST (Figure 2, j) is very close to the ERA5 blocking, as also demonstrated by the statistics:  $R \cong 0.98$ ,  $\text{RMSD} \cong 13 \text{ m}$ , and  $\sigma_{MM} \cong 56 \text{ m}$ . In future climate (Figure 2, l-m), the spatial characteristics of the blocking composites are very similar to the ones of the HIST period (these results will be confirmed in subsections 4.3 and 4.4). Considering the  $\Delta Z500_{\text{SSP2-HIST}}$  and  $\Delta Z500_{\text{SSP5-HIST}}$  results (Figure 2, n-o), blocking events get wider in SSP2 and especially in SSP5 and their centers are characterised by higher values. (The blocking composites computed for each GCM in the future are in Figures S2 and S3 in the Supplement.) Thus, we find that atmospheric blocking presents a dynamical component whose pattern is relatively stationary over the 21st century and a thermodynamical component that, as expected, is broadly driven by the overall warming of air masses in relation with the anthropogenic signal.

## 4.2 Frequency and duration of blocking events

Blocking events are identified for each GCM following the definition in subsection 3.3. The number of blocking days and blocking events per winter averaged over all winters of the 30-year periods and the duration of blocking events averaged over these periods are graphically represented in Figure 3 to facilitate the comparison of the HIST results against the future

**Taylor diagram for the mean composites over all blocking events for ERA5 and all GCMs for the winter HIST period (1980–2009). The diagram allows to quantify standard deviation (black), correlation coefficient (light blue), and root-mean-square difference (green) between the mean GCM composites and the mean ERA5 composite. SSP2 and SSP5 results are obtained with  $\Delta Z_{500\text{SSP2}}$  and  $\Delta Z_{500\text{SSP5}}$ , respectively.**




**Figure 2.** Blocking composites averaged over all blocking events for all GCMs (a-i) and ERA5 (k) during the winter HIST period (1980–2009). The multi-model means are computed over all blocking events of the six selected GCMs for the HIST period (1980–2009) using  $\Delta Z_{500\text{HIST}}$  (j) and for the future period (2070–2099) using  $\Delta Z_{500\text{SSP2}}$  (l),  $\Delta Z_{500\text{SSP5}}$  (m),  $\Delta Z_{500\text{SSP2-HIST}}$  (n), and  $\Delta Z_{500\text{SSP5-HIST}}$  (o).

(SSP2 and SSP5) results. We find that, during recent-past conditions, the MM mean number of blocking days per winter is about 30 and the MM mean number of events per winter is about 3. Our results are slightly lower than the findings of Mokhov et al. (2014), 35.8 days and 4.7 events, who detected blocking events in an Euro-Atlantic sector using a Z500-based blocking index applied on one GCM (IPSL). The MM mean of blocking duration is  $9.9 \pm 0.9$  days and is close to the mean duration of blocking events of  $10.2 \pm 5.3$  days obtained with the reanalysis. These results are in agreement with mean blocking durations found in the literature, e.g. 10.5 days for winter blocking in the European-Atlantic sector by Lupo et al. (2019), using reanalysis (NCEP/NCAR) of Z500, and 7.6 days by Mokhov et al. (2014). In summary, the mean temporal characteristics of blocking events during HIST are well reproduced by the GCMs: the MM means of number of blocking days and duration during HIST are close to the results obtained with the reanalysis, although most of the models tend to underestimate these quantities. When analysing the impact of climate change, no significant impact is found on blocking frequency and duration. With respect to the HIST results, MPI, BCC, and MRI simulate less frequent blocking events in both future scenarios, while the other GCMs present a higher blocking frequency (Figure 3). However, the uncertainty of the results is large (Table S2), so the differences between the periods are not statistically significant. Additionally, results for SSP2-4.5 and SSP5-8.5 are not in agreement among the various models (sometimes estimates are higher in SSP2 and sometimes in SSP5). Actually, a clear long-term change in blocking frequency in the past has not emerged so far (Barnes et al., 2014; Woollings et al., 2018), and there is no general consensus on the tendency of blocking frequency in future climate (Woollings et al., 2018). For example, Matsueda and Endo (2017) found a significant decrease in blocking frequency in the European-Atlantic sector involving all durations of blocking events simulated with six CMIP5-GCMs, Mokhov et al. (2014) found a general increase in the blocking frequency, while Masato et al. (2014) found that European blocking frequency remains unchanged using four CMIP5-GCMs.


The analysis of the occurrence of blocking events as a function of duration also indicates that the GCM projections agree well with the reanalysis (Figure 4 and Figure S4). Occurrence of blocking events decreases exponentially with duration, consistent with the findings of Wiedenmann et al. (2002); Barriopedro et al. (2006); Matsueda et al. (2009); Dunn-Sigouin and Son (2013); Mokhov et al. (2014). The distributions of all periods show long tails up to 30 days, but some isolated events can be even longer. In future climate, we find that the occurrence of short (5–8 days) blocking events slightly increases under the SSP5-8.5 scenario, while the occurrence of long (more than 10 days) events tends to decrease, as indicated by the mean lifetime ( $\tau$ ) of the exponential fit, which is lower for SSP5 ( $\tau \approx 9$  days) than for SSP2 and HIST ( $\tau \approx 10$  days).

### 4.3 Centers of blocking events

We now analyse the blocking centers (as defined in subsection 3.4) of the composites of blocking events in terms of their location and intensity, i.e. the value of  $\Delta Z500$  at that location. The geographical distribution of the center locations averaged over all blocking events of a given 30-year period is shown in Figure 225 (a-b). The ERA5-center is located over Sweden. The GCM-centers during HIST are over and close to the Scandinavian peninsula. In the future, we observe a general eastward shift of the center locations. In particular, four out of six models during SSP2 and SSP5 show blocking centers that are eastward with respect to the centers in HIST. The SSP2- and SSP5-MM means of the center locations are located about  $4^\circ$  and  $5^\circ$  eastward to the HIST-MM mean, respectively. An eastward shift of European blocking would lead to an increase of blocking over




**Figure 3.** Number of blocking days (*left*) and blocking events (*center*) averaged over all winters of the 30-year periods, and mean duration (in days) of blocking events occurred in 30 winters (*right*) for recent-past climate and future scenarios (SSP2-4.5 and SSP5-8.5) considering  $\Delta Z500_{HIST}$  (x-axis) and  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$  (y-axis). The black dashed line is the ERA5 mean. (The values are taken from Table S2.)



**Figure 4.** Occurrence of blocking events as a function of blocking duration for ERA5 and MM means during HIST, SSP2, and SSP5, considering  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$ , and  $\Delta Z500_{SSP5}$ . Exponential fits are drawn for ERA5 and MM means.

255 Western Russia (Dunn-Sigouin and Son, 2013). More uncertain is the meridional shift of the centers in the future. An eastward and northeastward shift of European blocking was also found by Masato et al. (2013, 2014) and Sillmann and Croci-Maspoli (2009), respectively. However, it must be stressed that there is a large variability associated to the blocking center locations on both meridional and zonal directions (as attested by the error bars in Figure ??5), and the shift of the centers is not significant.



**Figure 5.** (a-b) Locations of the blocking composite centers averaged over all blocking events for ERA5 and the GCMs during HIST considering  $\Delta Z500_{HIST}$  (top) and during SSP2 and SSP5 considering  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$  (bottom). The error bars indicate the standard deviations of latitudinal and longitudinal coordinates of the blocking centers. (c) Intensities (in m) of the blocking composite centers averaged over all blocking events during HIST, SSP2, and SSP5 considering  $\Delta Z500_{HIST}$  on the x-axis and  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$  on the y-axis. The black dashed line is the ERA5 mean. (The values are taken from Table S3.)

The MM mean of the blocking center intensities during HIST is  $248 \pm 18$  m, very close to the ERA5-intensity,  $251 \pm 48$  m (Table S3). The minimum intensity is simulated by INM,  $219 \pm 50$  m, the maximum one by MPI,  $273 \pm 61$  m. Under the SSP2-4.5 and SSP5-8.5 scenarios, the center intensities are very similar, on average, to the intensities of the blocking events in recent-past conditions (Figure 5, left). Also the variability of the centers, in terms of standard deviations and minimum-maximum intervals of the intensities (Table S3), do not change, implying that the future blocking intensities will be not affected by atmospheric dynamical changes. Additionally, we observe that differences between SSP2 and SSP5 periods are smaller than inter-model differences. ~~On the contrary, the center intensities of the future blocking events identified using  $\Delta Z500_{SSP2-HIST}$  and  $\Delta Z500_{SSP5-HIST}$  (i.e. the anomalies obtained using the recent-past climatology as reference) increase with respect to the recent-past conditions in both future scenarios, especially in the worst-case scenario (Figure 5, right). In particular, their MM means increase up to 306 m in SSP2 and 344 m in SSP5. The increments of the center intensities (i.e. of the geopotential~~

height) in this case are mainly explained by the general warming related to the anthropogenic greenhouse gases emissions

270 occurring under the considered scenarios.

Intensities (in m) of the blocking composite centers averaged over all blocking events during HIST, SSP2, and SSP5 considering  $\Delta Z500_{HIST}$  on the x-axis and  $\Delta Z500_{SSP2}$  and  $\Delta Z500_{SSP5}$  (left) and  $\Delta Z500_{SSP2-HIST}$  and  $\Delta Z500_{SSP5-HIST}$  (right) on the y-axis. The black dashed line is the ERA5 mean. (The values are taken from Table S3.)

#### 4.4 Size of blocking events


##### 275 4.4.1 Composite WTD method

Blocking area is computed for each composite of blocking event by using the composite WTD method described in subsection 3.4. This method takes into account events whose center is between 30°W and 50°E to focus on European blocking, although some events can extend westwards in the European-Atlantic sector. This is due to the fact that the events are identified via a partitioning algorithm (k-means) and not via blocking indexes designed for geopotential fields that are typical during atmospheric blocking. We could verify that, on average, only four events per GCM (i.e. ~ 4%) are of this type during HIST; we preferred not to disregard them in order not to introduce any subjectivity into the analysis, and the results are considered as an overestimation of the blocking size.

280 The MM mean size in HIST is  $9.1 \cdot 10^6$  km<sup>2</sup>, very close to the value obtained for ERA5 (Table S4). As expected from Figure 2 (j-m), the future blocking size is comparable to the recent-past blocking size. Actually, the MM mean size decreases by  $0.3 \cdot 10^6$  km<sup>2</sup> (i.e. about 3%) during SSP2 and SSP5, although not in a statistically significant way. Different results are obtained for the future blocking size computed with respect to the recent-past climatology (as anticipated in Figure 2, n-o). In this case, there is a clear tendency of blocking size to increase in the future; the size is nearly twice (1.7) and three times (2.7) larger during SSP2 and SSP5, respectively. This is mainly due to the global warming projected with the SSP2-4.5 and SSP5-8.5 scenarios, as already noted in subsection 4.1.

290 We further We also analyse the blocking size results in relation to the center intensity. We find a linear relation between size of blocking events and intensity of blocking centers (Figure ??6, left). The correlation is significant and higher than 0.7 during HIST and SSP2. The linear relation is in agreement with Barriopedro et al. (2010). Again, our results are in line with previous studies that followed a different approach for the blocking detection, based on the use of blocking indexes instead of the WTD.

295 Figure ?? (left) Moreover, Figure 6 (right) and Figure S5 show that blocking size is characterised by a normal distribution (e.g. Whiteman, 1982; Barriopedro et al., 2006). This is valid in all periods. In particular, the MM means of blocking area during SSP2 and SSP5 are very similar and close to the MM mean of the past blocking area. Different results are obtained for future blocking size computed with  $\Delta Z500_{SSP2-HIST}$  and  $\Delta Z500_{SSP5-HIST}$  (Figure ??, right). In this case, more blocking events with larger size will occur, as proved by the shift of the distribution towards higher values and its increasing width. Moreover, we notice that the blocking size in the future can be only roughly approximated by a gaussian law, especially in SSP5. It must 300 be reminded that the threshold chosen for  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$ , and  $\Delta Z500_{SSP5}$  (75 m) is lower than the threshold used with  $\Delta Z500$  with respect to the recent-past climatology (100 m, see subsection 3.4).



**Figure 6.** (Left) Mean area versus mean center intensity of blocking events computed for ERA5 and the GCMs during HIST, SSP2, and SSP5. (Right) Mean-area-Occurrence of blocking events computed with the DG method versus mean-area-as a function of blocking events computed with area for ERA5 and the composite-method GCMs during HIST, SSP2, and SSP5.  $R$  is the correlation;  $a$  is the slope of the regression lines (found by the least-squares fit) Gaussian fits are drawn when the correlation is statistically significant (at the 90% confidence level) for ERA5 and MM means. Both plots show the results for  $\Delta Z500_{HIST}$ ,  $\Delta Z500_{SSP2}$ , and  $\Delta Z500_{SSP5}$ .


#### 4.4.2 Comparison with the DG method

In order to check the reliability of the composite WTD method in estimating the area of blocking events, we compute that area by another approach relying on the DG index to identify blocking events, as done by Nabizadeh et al. (2019). As indicated in 305 subsection 3.4, the latter events will be denoted DG-blocking events; for clarity, in this section, the blocking events identified by the WTD will be denoted WTD-blocking events. We apply the DG method to compute the area of those DG-blocking days that belong to the WTD-blocking events (see subsection 3.4). Despite the number of these DG-blocking days may not match with the duration of the respective WTD-blocking events, we find that the two quantities generally agree (Figure S6).

The blocking areas resulted from the composite-WTD method and the DG method are compared in Figure ?? (right). 310 These areas are linearly correlated with statistical significance in all periods, the slopes of the linear regression being around 0.70 or larger (in SSP5). Therefore, the size of the blocking events identified via the WTD is in agreement with the size of the blocking events identified via the DG index.

## 5 Conclusions and discussion

We identify wintertime European blocking events by applying the weather type decomposition methodology on the European- 315 Atlantic sector. Our aim is to quantify the impact of climate change on the frequency, duration, and size of blocking events. For



**Figure 7.** Occurrence–Mean area of blocking events as a function of computed with the DG method versus mean area for ERA5 and of blocking events computed with the GCMs during HIST, SSP2, and SSP5–WTD method considering  $\Delta Z_{500}^{\text{HIST}}$ ,  $\Delta Z_{500}^{\text{SSP2}}$ , and  $\Delta Z_{500}^{\text{SSP5}}$ .  $R$  is the correlation;  $a$  is the slope of the linear regression; regression lines (left) and  $\Delta Z_{500}^{\text{SSP2-HIST}}$  and  $\Delta Z_{500}^{\text{SSP5-HIST}}$  (right) found by the least-squares fit. Gaussian fits are drawn for ERA5 and MM means. (GFDL when the correlation is excluded from statistically significant (at the MM means of  $\Delta Z_{500}^{\text{SSP5-HIST}}$ , 90% confidence level).

this purpose, we consider 30 years of historical runs and two future scenarios (SSP2-4.5 and SSP5-8.5) of nine CMIP6-GCMs. We show that the GCMs considered in this study capture well the spatio-temporal characteristics of atmospheric blocking in the recent-past climate, nevertheless, only those representing blocking patterns and variability closer to the reanalysis are used to investigate future blocking changes.

320 We find that the impact of climate change on blocking frequency and duration is not statistically significant, consistent with the literature. This is in line with the recent study of Huguenin et al. (2020), who found that changes of frequency and duration of weather types over central Europe in a warming climate are small and within the internal climate variability. By contrast, Fabiano et al. (2020), by applying the WTD as well, and Davini and D'Andrea (2020), by using blocking indexes, showed that frequency and duration of blocking events decrease significantly with climate change. It must be mentioned that 325 the methodology in Fabiano et al. (2020) is a bit different in terms of, for example, the data to be processed (the geopotential height anomalies are detrended and the seasonal cycle removal is different than here) and the number of retained eigenvectors (only four against twenty-four here). Overall, a clear long-term change in blocking frequency in the past has not emerged so far (Barnes et al., 2014; Woollings et al., 2018), and Woollings et al. (2018) highlighted the fact that there is no general consensus on the tendency of blocking frequency in the future (Woollings et al., 2018). future climate. In fact, Dunn-Sigouin and Son (2013) 330 and Matsueda and Endo (2017) found a significant decrease in blocking frequency in the European-Atlantic sector involving all durations of blocking events simulated with a subset of CMIP5-GCMs. Mokhov et al. (2014) found a general increase in the blocking frequency with four CMIP5-GCMs, while Masato et al. (2014) found that European blocking frequency remains

unchanged using four CMIP5-GCMs as well. Clearly, further investigations are needed, focusing on the last generation of models (i.e. CMIP6).

335 We introduce the ~~composite-WTD~~ method to quantify the size of blocking composites. We find that blocking area and center intensity are linearly correlated. Blocking patterns and size in the future are similar to the results obtained for the recent-past climate. This means that the spatial characteristics of blocking events will not change at the end of the century with respect to the future atmospheric mean state. ~~Instead, if we take into account the increasing geopotential height caused by a warmer climate, we find that the size of blocking events will significantly increase in the future, especially in the worst-case scenario.~~

340 We also apply another method, the DG method, to compute the blocking size, and we obtain similar results. ~~Our findings are in agreement with Hassanzadeh et al. (2014), who found smaller blocking area in a scenario where the midlatitude to pole surface temperature difference is reduced, as expected in warming climate; this result is in contrast to Nabizadeh et al. (2019), who showed that blocking size is projected to increase with climate change using a two-layer quasi-geostrophic model (and derived a scaling law for the blocking-event size). As only a few studies investigated blocking size, by applying different methodology and data sets (e.g idealized GCMs or GCMs from different CMIPs), more efforts should be devoted to this topic.~~

345 In general, we observe that the differences between SSP2- and SSP5-results are smaller than differences among the various ~~GCM-results~~GCM results, suggesting that there is no clear signal of climate change on blocking frequency, duration, and size. To the best of our knowledge, this is the first study investigating all these characteristics of blocking events that are identified via the WTD. Moreover, there are still few studies addressing this topic using GCMs of the CMIP6. Our results ~~for past climate~~ are in agreement with previous findings where blocking events are defined with blocking indexes. This confirms that the application of the WTD is also a good strategy to analyse blocking event characteristics.

350 ~~This study could be improved by analysing more GCMs, although other studies that considered many GCMs initially used only the best few GCMs for the analysis later; for example, Lee and Ahn (2017) selected five GCMs among twenty-two CMIP5-GCMs to study atmospheric blocking over the Pacific Ocean. Before comparing blocking event areas with other studies, it must be reminded that the results depend on the defined threshold. Finally~~Moreover, it must be pointed out that the four weather types imposed in the k-means algorithm allow to recover the ones usually obtained with the reanalysis. However, a different number of weather types may need to be computed in some models where the variability is different from the reanalysis (e.g. five regimes are considered in the CNRM model by Ménégoz et al. (2018)). ~~The optimal number of clusters also depends on the data to be processed; for instance, by applying the clustering on the full field data, Falkena et al. (2020) found that k=6 is an optimal choice.~~

360 ~~This study could be improved by analysing more GCMs, although other studies that considered many GCMs initially used only the best few GCMs for the analysis later; for example, Lee and Ahn (2017) selected five GCMs among twenty-two CMIP5-GCMs to study atmospheric blocking over the Pacific Ocean. Given the decadal variability of weather regimes (Dorrington and Stro, longer past and future periods could be considered (e.g. periods of 50 years, like in Fabiano et al. (2020)) so as to better smooth the dependency of the results on this decadal variability. Moreover, those days for which the geopotential height anomaly field does not resemble the blocking weather regime pattern could be classified as “neutral days”, like in Dorrington et al. (2021), and excluded from the analysis. Finally, a sensitivity analysis of the results to the clear-cut character of the clusters could be~~

conducted, for instance by removing the influence of the jet speed from the geopotential height field, like in Dorrington and Strommen (2020).

~

370 *Author contributions.* All authors contributed to designing the study. SB and FJ analysed the data, together with ELB, and produced the figures. All the authors discussed the results. SB wrote the paper; all the authors provided assistance in finalizing the article.

*Competing interests.* The authors declare that they have no conflict of interest.

375 *Acknowledgements.* The authors thank Dr. E. Nabizadeh for the useful discussion. The authors acknowledge the World Climate Research Programme (WCRP), which coordinated and promoted CMIP6, the climate modeling groups for producing and making available their model output, and the Mésocentre ESPRI (Ensemble de services pour la recherche à l'IPSL, <https://mesocentre.ipsl.fr/>) for archiving the data and providing the access to the CMIP6 data. SB thanks the LEGI for the postdoctoral support; ELB and FJ thank the IDEX program MobilAir for the PhD and MSc support, respectively.

## References

Barnes, E. A., Dunn-Sigouin, E., Masato, G., and Woollings, T.: Exploring recent trends in Northern Hemisphere blocking, *Geophysical Research Letters*, 41, 638–644, <https://doi.org/10.1002/2013GL058745>, 2014.

380 Barriopedro, D., García-Herrera, R., Lupo, A. R., and Hernández, E.: A Climatology of Northern Hemisphere Blocking, *Journal of Climate*, 19, 1042–1063, <https://doi.org/10.1175/JCLI3678.1>, 2006.

Barriopedro, D., García-Herrera, R., and Trigo, R. M.: Application of blocking diagnosis methods to General Circulation Models. Part I: a novel detection scheme, *Climate Dynamics*, 35, 1373–1391, <https://doi.org/10.1007/s00382-010-0767-5>, 2010.

385 Berckmans, J., Woollings, T., Demory, M.-E., Vidale, P.-L., and Roberts, M.: Atmospheric blocking in a high resolution climate model: influences of mean state, orography and eddy forcing, *Atmospheric Science Letters*, 14, 34–40, <https://doi.org/10.1002/asl2.412>, 2013.

Boé, J. and Terray, L.: A Weather-Type Approach to Analyzing Winter Precipitation in France: Twentieth-Century Trends and the Role of Anthropogenic Forcing, *Journal of Climate*, 21, 3118–3133, <https://doi.org/10.1175/2007JCLI1796.1>, 2008.

390 Cassou, C.: Intraseasonal interaction between the Madden–Julian Oscillation and the North Atlantic Oscillation, *Nature*, 455, 523–527, <https://doi.org/10.1038/nature07286>, 2008.

Cassou, C., Terray, L., Hurrell, J. W., and Deser, C.: North Atlantic Winter Climate Regimes: Spatial Asymmetry, Stationarity with Time, and Oceanic Forcing, *Journal of Climate*, 17, 1055–1068, [https://doi.org/10.1175/1520-0442\(2004\)017<1055:NAWCRS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1055:NAWCRS>2.0.CO;2), 2004.

Cassou, C., Terray, L., and Phillips, A.: Tropical Atlantic Influence on European Heat Waves, *Journal of Climate*, 18, 2805–2811, <https://doi.org/10.1175/JCLI3506.1>, 2005.

395 Cheung, H., Zhou, W., Mok, H., Wu, M., and Shao, Y.: Revisiting the climatology of atmospheric blocking in the Northern Hemisphere, *Adv. Atmos. Sci.*, 30, 397–410, <https://doi.org/10.1007/s00376-012-2006-y>, 2013.

Chhin, R. and Yoden, S.: Ranking CMIP5 GCMs for Model Ensemble Selection on Regional Scale: Case Study of the Indochina Region, *Journal of Geophysical Research: Atmospheres*, 123, 8949–8974, <https://doi.org/10.1029/2017JD028026>, 2018.

Chung, C. and Nigam, S.: Weighting of geophysical data in Principal Component Analysis, *Journal of Geophysical Research: Atmospheres*, 104, 16 925–16 928, <https://doi.org/10.1029/1999JD900234>, 1999.

400 Cortesi, N., Torralba, V., González-Reviriego, N., Soret, A., and J.Doblas-Reyes, F.: Characterization of European wind speed variability using weather regimes, *Climate Dynamics*, 53, 4961–4976, <https://doi.org/10.1007/s00382-019-04839-5>, 2019.

Davini, D. and D’Andrea, F.: From CMIP3 to CMIP6: Northern Hemisphere Atmospheric Blocking Simulation in Present and Future Climate, *Journal of Climate*, 33, 10 021–10 038, <https://doi.org/10.1175/JCLI-D-19-0862.1>, 2020.

405 Davini, P., Cagnazzo, C., and Anstey, J. A.: A blocking view of the stratosphere-troposphere coupling, *Journal of Geophysical Research: Atmospheres*, 119, 11,100–11,115, <https://doi.org/10.1002/2014JD021703>, 2014.

Davini, P., Corti, S., D’Andrea, F., Rivière, G., and von Hardenberg, J.: Improved Winter European Atmospheric Blocking Frequencies in High-Resolution Global Climate Simulations, *Journal of Advances in Modeling Earth Systems*, 9, 2615–2634, <https://doi.org/10.1002/2017MS001082>, 2017.

410 Dole, R. M. and Gordon, N. D.: Persistent Anomalies of the Extratropical Northern Hemisphere Wintertime Circulation: Geographical Distribution and Regional Persistence Characteristics, *Monthly Weather Review*, 111, 1567–1586, [https://doi.org/10.1175/1520-0493\(1983\)111<1567:PAOTEN>2.0.CO;2](https://doi.org/10.1175/1520-0493(1983)111<1567:PAOTEN>2.0.CO;2), 1983.

Dorrrington, J. and Strommen, K. J.: Jet Speed Variability Obscures Euro-Atlantic Regime Structure, *Geophysical Research Letters*, 47, e2020GL087907, <https://doi.org/10.1029/2020GL087907>, 2020.

415 Dorrington, J., Strommen, K., and Fabiano, F.: How well does CMIP6 capture the dynamics of Euro-Atlantic weather regimes, and why?,  
Weather and Climate Dynamics Discussions, 2021, 1–41, <https://doi.org/10.5194/wcd-2021-71>, 2021.

420 Drouard, M. and Woollings, T.: Contrasting Mechanisms of Summer Blocking Over Western Eurasia, Geophysical Research Letters, 45,  
12,040–12,048, <https://doi.org/10.1029/2018GL079894>, 2018.

Dunn-Sigouin, E. and Son, S.-W.: Northern Hemisphere blocking frequency and duration in the CMIP5 models, Journal of Geophysical  
425 Research: Atmospheres, 118, 1179–1188, <https://doi.org/10.1002/jgrd.50143>, 2013.

Dunn-Sigouin, E., Son, S.-W., and Lin, H.: Evaluation of Northern Hemisphere Blocking Climatology in the Global Environment Multiscale  
Model, Monthly Weather Review, 141, 707–727, <https://doi.org/10.1175/MWR-D-12-00134.1>, 2013.

430 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model  
Intercomparison Project Phase 6 (CMIP6) experimental design and organization, Geoscientific Model Development, 9, 1937–1958,  
435 <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

Fabiano, F., Meccia, V., Davini, P., Ghinassi, P., and Corti, S.: A regime view of future atmospheric circulation changes in Northern mid-  
latitudes, Weather and Climate Dynamics Discussions, 2020, 1–24, <https://doi.org/10.5194/wcd-2020-37>, 2020.

Falkena, S. K., de Wiljes, J., Weisheimer, A., and Shepherd, T. G.: Revisiting the identification of wintertime atmospheric circulation regimes  
440 in the Euro-Atlantic sector, Quarterly Journal of the Royal Meteorological Society, 146, 2801–2814, <https://doi.org/10.1002/qj.3818>, 2020.

Francis, J. A. and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes, Geophysical Research Letters,  
445 39, <https://doi.org/10.1029/2012GL051000>, 2012.

Hassanzadeh, P., Kuang, Z., and Farrell, B. F.: Responses of midlatitude blocks and wave amplitude to changes in the meridional temperature  
gradient in an idealized dry GCM, Geophysical Research Letters, 41, 5223–5232, <https://doi.org/10.1002/2014GL060764>, 2014.

Hausfather, Z. and Peters, G.: Emissions – the 'business as usual' story is misleading, Nature, 577, 618–620, <https://doi.org/10.1038/d41586-020-00177-3>, 2020.

450 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Sim-  
mons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren,  
P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J.,  
Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Vil-  
laume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, Quarterly Journal of the Royal Meteorological Society, 146, 1999–2049,  
<https://doi.org/10.1002/qj.3803>, 2020.

Hertig, E. and Jacobbeit, J.: Variability of weather regimes in the North Atlantic-European area: past and future, Atmospheric Science Letters,  
15, 314–320, <https://doi.org/10.1002/asl2.505>, 2014.

Hou, P. and Wu, S.: Long-term Changes in Extreme Air Pollution Meteorology and the Implications for Air Quality, Scientific Reports, 6,  
455 23 792, <https://doi.org/10.1038/srep23792>, 2016.

Huguenin, M. F., Fischer, E. M., Kotlarski, S., Scherrer, S. C., Schwierz, C., and Knutti, R.: Lack of Change in the Projected  
Frequency and Persistence of Atmospheric Circulation Types Over Central Europe, Geophysical Research Letters, 47,  
<https://doi.org/10.1029/2019GL086132>, 2020.

Jiménez, P. A., González-Rouco, J., Montávez, J., García-Bustamante, E., and Navarro, J.: Climatology of wind patterns in the northeast of  
460 the Iberian Peninsula, International Journal of Climatology, 29, 501–525, <https://doi.org/10.1002/joc.1705>, 2009.

Kautz, L.-A., Martius, O., Pfahl, S., Pinto, J. G., Ramos, A. M., Sousa, P. M., and Woollings, T.: Atmospheric Blocking and Weather Extremes over the Euro-Atlantic Sector – A Review, *Weather and Climate Dynamics Discussions*, 2021, 1–43, <https://doi.org/10.5194/wcd-2021-56>, 2021.

Khan, N., Shahid, S., Ahmed, K., Wang, X., Ali, R., Ismail, T., and Nawaz, N.: Selection of GCMs for the projection of spatial distribution of heat waves in Pakistan, *Atmospheric Research*, 233, 104 688, <https://doi.org/10.1016/j.atmosres.2019.104688>, 2020.

455 Largeron, Y. and Staquet, C.: Persistent inversion dynamics and wintertime PM<sub>10</sub> air pollution in Alpine valleys, *Atmospheric Environment*, 135, 92–108, <https://doi.org/10.1016/j.atmosenv.2016.03.045>, 2016.

Lee, D. Y. and Ahn, J.-B.: Future change in the frequency and intensity of wintertime North Pacific blocking in CMIP5 models, *International Journal of Climatology*, 37, 2765–2781, <https://doi.org/10.1002/joc.4878>, 2017.

460 Lupo, A., Jensen, A., Mokhov, I., Timazhev, A., Eichler, T., and Efe, B.: Changes in Global Blocking Character in Recent Decades, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10020092>, 2019.

Masato, G., Hoskins, B. J., and Woollings, T.: Winter and Summer Northern Hemisphere Blocking in CMIP5 Models, *Journal of Climate*, 26, 7044–7059, <https://doi.org/10.1175/JCLI-D-12-00466.1>, 2013.

Masato, G., Woollings, T., and Hoskins, B. J.: Structure and impact of atmospheric blocking over the Euro-Atlantic region in present-day and future simulations, *Geophysical Research Letters*, 41, 1051–1058, <https://doi.org/10.1002/2013GL058570>, 2014.

465 Matsueda, M. and Endo, H.: The robustness of future changes in Northern Hemisphere blocking: A large ensemble projection with multiple sea surface temperature patterns, *Geophysical Research Letters*, 44, 5158–5166, <https://doi.org/10.1002/2017GL073336>, 2017.

Matsueda, M., Mizuta, R., and Kusunoki, S.: Future change in wintertime atmospheric blocking simulated using a 20-km-mesh atmospheric global circulation model, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2009JD011919>, 2009.

470 Ménégoz, M., Cassou, C., Swingedouw, D., Ruprich-Robert, Y., Bretonnière, P.-A., and Doblas-Reyes, D.: Role of the Atlantic Multidecadal Variability in modulating the climate response to a Pinatubo-like volcanic eruption, *Climate Dynamics*, 51, 1863–1883, <https://doi.org/10.1007/s00382-017-3986-1>, 2018.

Michelangeli, P.-A., Vautard, R., and Legras, B.: Weather Regimes: Recurrence and Quasi Stationarity, *Journal of the Atmospheric Sciences*, 52, 1237–1256, [https://doi.org/10.1175/1520-0469\(1995\)052<1237:WRRAQS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1237:WRRAQS>2.0.CO;2), 1995.

475 Mokhov, I. and Timazhev, A.: Atmospheric Blocking and Changes in its Frequency in the 21st Century Simulated with the Ensemble of Climate Models, *Russ. Meteorol. Hydrol.*, pp. 369–377, <https://doi.org/10.3103/S1068373919060013>, 2019.

Mokhov, I., Akperov, M., Prokofyeva, M., Timazhev, A., Lupo, A., and Le Treut, H.: Blockings in the Northern Hemisphere and EuroAtlantic Region: Estimates of Changes from Reanalysis Data and Model Simulations, *Doklady Earth Sciences*, 449, 430–433, <https://doi.org/10.1134/S1028334X13040144>, 2013.

480 Mokhov, I. I., Timazhev, A. V., and Lupo, A. R.: Changes in atmospheric blocking characteristics within Euro-Atlantic region and Northern Hemisphere as a whole in the 21st century from model simulations using RCP anthropogenic scenarios, *Global and Planetary Change*, 122, 265–270, <https://doi.org/10.1016/j.gloplacha.2014.09.004>, 2014.

Munoz, C., Schultz, D., and Vaughan, G.: A Midlatitude Climatology and Interannual Variability of 200- and 500-hPa Cut-Off Lows, *Journal of Climate*, 33, 2201–2222, <https://doi.org/10.1175/JCLI-D-19-0497.1>, 2020.

485 Nabizadeh, E., Hassanzadeh, P., Yang, D., and Barnes, E. A.: Size of the Atmospheric Blocking Events: Scaling Law and Response to Climate Change, *Geophysical Research Letters*, 46, 13 488–13 499, <https://doi.org/10.1029/2019GL084863>, 2019.

Patterson, M., Bracegirdle, T., and Woollings, T.: Southern Hemisphere Atmospheric Blocking in CMIP5 and Future Changes in the Australia-New Zealand Sector, *Geophysical Research Letters*, 46, 9281–9290, <https://doi.org/10.1029/2019GL083264>, 2019.

Philipp, A., Beck, C., Huth, R., and Jacobbeit, J.: Development and comparison of circulation type classifications using the COST 733 dataset  
490 and software, *International Journal of Climatology*, 36, 2673–2691, <https://doi.org/10.1002/joc.3920>, 2016.

Pinheiro, M., Ullrich, P., and Grotjahn, R.: Atmospheric blocking and intercomparison of objective detection methods: flow field characteristics, *Climate Dynamics*, 53, 4189–4216, <https://doi.org/10.1007/s00382-019-04782-5>, 2019.

Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz,  
495 W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P.,  
Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaaf, D., Masui, T., Rogelj, J., Strefler, J., Drouet,  
L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-  
Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse  
gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>,  
2017.

Russo, A., Trigo, R. M., Martins, H., and Mendes, M. T.: NO<sub>2</sub>, PM10 and O<sub>3</sub> urban concentrations and its association with circulation  
500 weather types in Portugal, *Atmospheric Environment*, 89, 768–785, <https://doi.org/10.1016/j.atmosenv.2014.02.010>, 2014.

Sáenz, F. and Durán-Quesada, A.: A climatology of low level windregimes over Central America using a weather type classification approach,  
Front. Earth Sci., 3, <https://doi.org/10.3389/feart.2015.00015>, 2015.

Schiemann, R., Demory, M., Shaffrey, L. C., Strachan, J., Vidale, P. L., Mizielski, M. S., Roberts, M. J., Matsueda, M., Wehner, M. F., and  
505 Jung, T.: The Resolution Sensitivity of Northern Hemisphere Blocking in Four 25-km Atmospheric Global Circulation Models, *Journal of Climate*, 30, 337–358, <https://doi.org/10.1175/JCLI-D-16-0100.1>, 2017.

Sillmann, J. and Croci-Maspoli, M.: Present and future atmospheric blocking and its impact on European mean and extreme climate, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL038259>, 2009.

Tibaldi, S. and Molteni, F.: On the operational predictability of blocking, *Tellus A*, 42, 343–365, [https://doi.org/10.1034/j.1600-0870.1990.t01-2-00003.x](https://doi.org/10.1034/j.1600-<br/>510 0870.1990.t01-2-00003.x), 1990.

Ullmann, A., Fontaine, B., and Roucou, P.: Euro-Atlantic weather regimes and Mediterranean rainfall patterns: present-day variability and  
expected changes under CMIP5 projections, *International Journal of Climatology*, 34, 2634–2650, <https://doi.org/10.1002/joc.3864>, 2014.

Whiteman, C. D.: Breakup of Temperature Inversions in Deep Mountain Valleys: Part I. Observations, *Journal of Applied Meteorology*, 21,  
270–289, [https://doi.org/10.1175/1520-0450\(1982\)021<0270:BOTIID>2.0.CO;2](https://doi.org/10.1175/1520-0450(1982)021<0270:BOTIID>2.0.CO;2), 1982.

515 Wiedenmann, J. M., Lupo, A. R., Mokhov, I. I., and Tikhonova, E. A.: The Climatology of Blocking Anticyclones for the Northern and Southern Hemispheres: Block Intensity as a Diagnostic, *Journal of Climate*, 15, 3459–3473, [https://doi.org/10.1175/1520-0442\(2002\)015<3459:TCOBAF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<3459:TCOBAF>2.0.CO;2), 2002.

Woollings, T., Barriopedro, D., Methven, J., Son, S.-W., Martius, O., Harvey, B., Sillmann, J., Lupo, A. R., and Seneviratne, S.: Blocking  
and its Response to Climate Change, *Current Climate Change Reports*, 4, 287–300, <https://doi.org/10.1007/s40641-018-0108-z>, 2018.