

1 General comments

The authors present a really nice study about the reconstruction of winter climate anomalies in the Euro-Atlantic sector using different measures to classify large scale flow anomalies. This is done with the broadly used global pattern index of the NAO, widely recognised as an important factor to influence winter anomalies over Europe. The authors further use blocking regions and jet classifications in the North Atlantic to investigate how well European winter anomalies, in temperature and precipitation, can be reconstructed by those groups (NAO, blocking jet). The authors do not do this solely by using simple correlations, but also another measure also accounting for the represented temporal variability (amplitude) of those groups. The manuscript is very well structured, everything is quite well explained and results are presented in a very coherent way, guiding the reader through the individual steps of the results and how they are connected. The authors also spent enough time to explain the methods and give the reader the necessary information (e.g. including Fig. 1) to understand the applied diagnostics without having to check the references for the essential informations. All in all, very nicely done.

I only have two major issues. The smaller issue concerns the application of the diagnostics in the result section. It wasn't always fully clear which of the specific representation of their diagnostics is applied, e.g. concerning the CE representation. While reading the result section I thought the authors always use the scaled reconstruction (with β), only after they explicitly mentioned that they introduce now the scaling did it become clear to me that they didn't use it before. Also, the authors could have clarified how Equ. (4) and Equ. (6) are used in this manuscript. If I'm not mistaken, only the simplification (Equ. 4) is used. And although highlighting the different Equations, the authors did not make a statement on the impact of the different choices, except for saying that the assumption of a zero climatological mean of the reconstructed anomalies should be fulfilled. More specific comments about this can be found in the "Specific comments" section.

My main issue with the manuscript is that the authors maybe did not the best job in highlighting the importance of their study and the usefulness of their presented results. The final discussion is full of very interesting aspects which are closely linked to this study, but which weren't analysed. This includes for example the extension of the regions to identify the classifications of a specific group (e.g. the limited eastward extension of the jet-region), the number of classifications within a group, the impact of extreme events for this kind of climate reconstruction. Also, the main conclusion I did get from this is that somehow all those groups are able to some extent to reconstruct climate anomalies, but it is also not clear which one is best. The thing is that it cannot be surprising that any kind of large scale feature which represents a large part of the variability of winter anomalies (variability in the same variable) is very likely able to identify large scale seasonal anomalies in other variables like temperature or precipitation, as those variables are linked to each other (no rain within a blocking high, storms moving along the jet, etc.). I think the authors could made clearer what there main message is, why it might be important to use different classes. How can those results

interpreted and maybe applied to other fields (e.g. if climate models are able to capture the general jet behaviour over the Atlantic this might be not enough as the exact representation of the established blocking periods decides on whether or not the model can have skill for specific parts of Europe in reconstructing the climate). Not sure the above examples make any sense at all, but I try to say with that that the authors could put some more effort into selling their results, why is this study so useful (I think this manuscript is slightly missing this). This is probably even more important because the authors restricted their analysis strongly to the choices they have done. As previously mentioned, and also raised by the authors, there are so many interesting modifications of the setup of the study and modifications of the choices which weren't done by the authors, although probably having important impact on the results. Also are there a few questions arising from the conclusions of the authors, e.g. can the authors quantify the relevance of remote regions (Pacific jet configurations, ENSO, North American or Pacific blocking) for the European winter climate reconstruction (as authors highlight the importance of jet classification region to European climate response). To summarize, I think if the authors could highlight more the important aspects of their results beyond the connection that large scale flow anomalies are connected with large scale climate response, the impact and relevance of this study could be improved.

One example for highlighting the results of this manuscript could be the connection between the different groups of large scale flow representation. Not sure how new and innovative this is, but I really appreciated how well the authors presented the connection between those large scale flow anomalies (link between NAO, blocking and jet). This was really nicely presented (e.g. Fig 2-4 and Fig. 5). Not sure if this was already done in a similar fashion in other studies, but I expected this to be highlighted at the end, because it nicely show the connections and links between those different groups (not only using only one group, like NAO, but compare it with different large scale flow realisations and show how those are connected). Maybe one could have also interpreted how this can be used to understand the similar usefulness to reconstruct seasonal anomalies or how specific aspects/regions cannot be as well reconstructed as others for a specific group.

2 Specific comments

2.1 Introduction

lines 29-30:

I wonder if “obstructs” is the appropriate word, as specific locations of a blocking high can also lead to a northward or southward shift of the jet - so changing the path of the jet but not really fully blocking it. So maybe it would be better to specify “obstruction” in terms of obstructing the climatological position of the westerly flow (which of course will have an impact on seasonal anomalies in temperature, etc.)?

line 36:

Not sure how well the term storminess is established. I think it would need a short ex-

planation. Therefore I would suggest to remove “with distinct patterns of storminess”, because the sentence works also without this term.

Table 1:

Maybe replace “NAO values above (below) 0.5” with “NAO values above 0.5 (below -0.5)”.

lines 83-87:

Reading this, one might wonder about the choice of the distance $d = 0.5$ if 46 % cannot be classified. This does sound like only the days very close to a centroid are chosen, which seems fine to create clear jet classifications. However, the explanation is somewhat surprising, namely that the large group of undefined cases is in particular occurring during the transition days, which would mean that either the the different jet classification are associated with frequent changes or that the transition phase between the jet classifications is rather long. Is the choice of $d=0.5$ the same as in Madonna et al (2017)?

reference also to section 2.2.2:

Also having read the part of A_rec, this jet classification seems relevant, as stricter classification choices (larger d) leads to less days recognized as jet classification. Since A_rec is normalized by the full number of days (90), this could lead to an increase in the amplitude error (for CE, Equ. 4). So introducing the beta factor (Equ. 7) is essential to account for this and therefore corrects possible “missing” members of specific classifications, or am I mistaken? But in lines 137-139 the authors only mention the problem of a non-gaussian distribution which “requires” a correction with this β -factor. Not sure I made myself clear, therefore I include a short example. If I could assume that the seasonal anomalies are mainly driven by a N-jet and a S-jet, the choice of a larger d can lead to a underrepresentation of the anomalies for seasons mainly associated with either a N-jet or S-jet, meaning that the associated a-value will become smaller and with it will affect CE. So the β -factor will boost the dominant anomaly. If this is true, maybe this should also be included as a reason in lines 137-139.

lines 95-97:

Why do the authors not apply the same anomaly calculation to all variables? The position of the jet does also follow a seasonal cycle, following the location of the temperature gradients. Therefore, what is the reason for not applying the same anomaly calculation? If there is a reason for doing so, e.g. signal otherwise too noisy because of the mentioned strong fluctuations, I think this is worth mentioning. Otherwise the reader will wonder about the different procedures for the different variables.

Equation (3):

This refers to Equation (3) and the whole section 2.2.1. According to Equ. (3), CE represents a 2d field and the given equations and text (CE values of 0, 1, etc) are for time series at one gridpoint, correct? While first scanning the manuscript, I expected CE to be a single-value metric to capture the overall skill of the 2d reconstruction (because

the examples were for one value, e.g. $CE = 1$), but in section 2.2.1 references are only made to time series, so CE must have values at each longitude and latitude. To make this fully clear and prevent confusion, CE could be written in the same way as A_{rec} , as $CE(\phi, \lambda)$. Alternatively a short sentence mentioning this would be helpful.

Section 2.2 Seasonal reconstructions:

The underlying assumption of this procedure is that the unclassified days more or less represent the climatological mean, otherwise the large undefined classifications (with representing partly more than 50 %) would account for large anomalies that this procedure will never be able to capture. This means for normally distributed quantities, as long as there is no mixing of different classifications (e.g. one jet classification and one blocking classification), that this method tests how well the undefined classifications represent the climatology (i.e. is there a missing classification with strongly impacts seasonal anomalies), right? Maybe this is worth mentioning (if true), because this could get lost in the text with all the description of weighting procedure and combination of classifications? However, non-normally distributed quantities would also lower the reconstruction skill. So probably these two aspects can lead to a reduced reconstruction skill, non-climatological rest-classifications and non-normal distributions. Maybe worth mentioning here, because both aspects are a necessary assumption for this method.

line 120:

Same as in last comment, the assumption hereby is that the undefined classifications are close to climatology, otherwise one cannot assume that the average of p is zero. As this is now part of the definition of CE (Equation 4), I think this should be clarified, or alternatively the step from Equ. (3) to Equ. (4) should be justified.

line 129:

Where does the $CE = 0.25$ comes from? Is this a usual value indicating good reconstructions, also used in other studies? Or alternatively the authors could give some further insight into the simple function CE. The global maximum of this CE function, as function of a , is at the location $r = a$, which means that the maximum value of CE for any a is r^2 . This would mean that independent of the amplitude error (or a), the choice of $CE = 0.25$ requires a minimum correlation of 0.5, whereas for a perfect correlation the values of a must be between 0.13 and 1.87 (if I am not mistaken). I would find such a simple consideration of CE very helpful, otherwise I would feel a bit lost when reading about the choice of 0.25 (because initially, I had no feeling what this value means in terms of correlation or amplitude error). What I find slightly confusing about this CE-value is that for exceeding the 0.25 value at low correlations (slightly above 0.5), it must include an amplitude error ($a < 1$) in order to exceed the threshold of 0.25, so low correlations can be corrected by a reduced variance in the reconstructed fields. But the overall conclusion from this function is, that it is a well established, easy measure to evaluate the reconstruction of an original field?

lines 139-140:

But using the median would lead to a bias in the overall climatological reconstruction of this variable (sum/average over all seasons), wouldn't it? But this would be fine, because this bias is not relevant for the calculation of CE, is this correct? So what I found surprising while first reading this sentence is that there is no condition that the climatological reconstruction of this method does not necessarily need to represent the climatology. Sorry, not sure this even is a question, or if it is obvious and not worth mentioning.

line 145:

"...calculate the centre of mass of the CE...". Is there a weighting by the value of CE involved to calculate the centre of mass, or is this purely by distance in correlation-amplitude space with each value represented by a weighting value of 1 (e.g. what is the "mass" here)? I see, in line 149 the points are restricted to correlation values of above 0.5, which probably suggests that the weights are all 1. I would suggest to be clarify this in the text.

line 155:

Is uncertainty the correct term here? It does not really test the uncertainty of the calculation of CE, but checks if the resulting CE-values are statistical significant, doesn't it?

2.2 Results:

line 190, Fig. 5:

Very nice overview of the seasonal connection between different, already coherently grouped, classifications. It would be nice if the NAO- would also have the associated correlation value with the jet classification, as all the other values have. Now with the missing r-value for NAO-, one has to check always the figure caption to understand for which variable GB has a correlation of 0.64. I would also include the NAO+ correlations in the legend. As the variable without a correlation value is the one with which the correlation is calculated, there is never any confusion.

lines 190-197:

Coming back again to the choice of the d-value for the jet classifications. I think it would be helpful if the authors explicitly mention their "large" choice of $d = 0.5$. As for all cluster groups (e.g. NAO-, GB and S-jet - IWB and N-jet - etc) the jet part represents the smallest part (in terms of number of days), it seems like this is due to the restrictive choice of d and a smaller d would bring the number of days for the jet-groups closer to the rest. I would suggest the authors, as already previously mentioned, give some further support for this choice. If a smaller d than 0.5 is problematic, is this really due to the transition periods as the text of the authors seems to indicate (line 85)? If this is the case, maybe it is worth mentioning more specifically what the problem about the transition period is (are those so frequently that they can make up nearly 50 percent of the time?). If $d = 0.5$ was also chosen in the original paper and discussed in depth, I would suggest to explicitly cite it in relation to this aspect.

lines 200-201:

I see, very nice. Maybe it is worth already mentioning this while introducing the reconstruction method. However, this is for the composite of the full climatology. From what I mentioned before, my assumption was that for individual seasons with dominance of specific classifications (like N-jet), the anomaly of the unclassified cases will show a similar pattern. To estimate this effect a comparison of the variance would be helpful to get a deeper insight. Having a very small value of the anomalies for the climatological composite is nice to conclude, that the remaining unclassified cases indeed represents climatology, but it would also be nice to have a similar short comment about the variance. If it is small then the reader knows that there is absolutely nothing to worry about the procedure in this manuscript. If it is not really small, the authors could give a short explanation why they think that is (no optimal way of defining the classifications because of some cutoff problem between getting the relevant patterns and reducing the variance of the unclassified cases?).

lines 235-236, Figure S2:

Oh, it seems I did not understand the uncertainty estimation correctly. I thought that this resampling of the 35 years is done to either observation or reconstruction to get an estimate of how significant the resulting CE-value is. But in this case, I would assume that for the areas of very high correlations, some of them would also show very different values (lower) for the bootstrap mean, because the reconstructed time series was already capturing well the observations, whereas a resampling of the years will not do so. So I'm slightly puzzled about the near identity between CE and bootstrap mean, suggesting that I don't understand what was actually done. Maybe the authors can therefore explain in a bit more detail how the uncertainty estimation does or what it is for?

line 262:

That wasn't clear to me. I thought the scaling factor was always used after being introduced in the method section. Not sure I have missed this while reading through the text, but I would suggest to make this clear while showing CE for the first time (if not done and I just missed that).

lines 274-275, Fig. 10:

I would suggest to include the 0-line in the anomaly plots, which makes the reading of those plots much easier. This would also highlight the issue of the blocking classifications, that this one does for some regions seems to be only able to capture negative anomalies (10b, d and h). Further, it seems a bit confusing when talking about amplitudes while interpreting this plot. In my mind I quickly switched to my usual interpretation of amplitude, representing the strength of the anomaly. This was confusing, because e.g. Fig. 10b and d seem to indicate that for the blocking reconstruction negative amplitudes of the anomalies are being overestimated while not being able to represent positive anomalies. But here amplitude refers to a from the CE calculation, right? I would suggest to mention this again explicitly, which makes following the interpretations easier.

2.3 Discussion and Conclusion

line 286:

Although, the better reconstruction is more focused towards western Europe, whereas temperature is better reconstructed at more eastern and southeastern Europe (Fig. 9). So what does in general mean, a higher mean CE value, occurrence of very high CE values or more points showing $CE > 0.25$? I think the authors could present this a lot more convincingly, if they could attribute a quantification to it, using some sort of simple measure as mentioned above. Just by looking again at Fig. 9 this doesn't seem to be so obvious. I appreciate the authors give some reasoning behind the differences between temperature and precipitation in this paragraph, which represents some explanation beyond just showing the results of this correlation/CE test.

line 305:

So what is the actual main message of the manuscript. As long as one chooses classifications which are associated with large scale flow anomalies (NAO, blocking, jet), one will be able to capture to some extent the seasonal large scale anomalies over Europe. But this study does not come to a conclusion what metric is best and it also depends on what someone might be interested in (precipitation or temperature). So the main message is that there is no perfect metric to capture both temperature and anomalies? A possible limitation of the reconstruction could be due to extreme events, but those are not considered in this study.

lines 298-304:

I think I made this point already earlier: isn't this (necessity of including a scaling factor) also a problem of the reconstruction method? If one would only use NAO values above 2 or below -2, it would be obvious that the reconstruction would underestimate the amplitude, wouldn't it? So going down with this threshold to 1 would probably do a better job in reconstructing the seasonal anomalies, because more seasons will be represented within this class and more variability is possible. Whereas defining a class of NAO with values above -5 would do a very bad job, because it will only reproduce climatology. So there is some sort of useful cutoff, or the classification are already defined in a useful way (by EOF, clustering, etc are they already constructed in a way to capture the variability). So I assume my question is, if this shouldn't be rather discussed with more focus on the method / discussion of how necessary a scaling is to start with?

Further, I think it was not fully clear in the manuscript which definition of CE was used (I raised this point also in another comment). It wasn't clear to me that the first results were for the non-scaled CE values (sorry if I just missed it). But also the choice of CE (Equ. 4 or Equ. 6) was not fully clear to me. If I am not mistaken, Equ. (4) was taken to calculate the CE values, right? Was Equ. (6) also used somewhere? In general, what is the reason to use Equ. (4)? Since Equ. (6) doesn't really need more calculations or is too complicated, it is not obvious for me why exactly the simplification was used? Is the reason behind this the identification of a useful scaling (Fig. 8)? I think the authors could make this a bit clearer. Also, I hope I didn't miss it, but did the authors compare

the two calculations, how different are the results? This information could be useful to justify the general use and handling of Equ. (4).

lines 305-314 and lines 315-324:

Really interesting aspect in those two paragraphs. But the comparison between NAO (with large region) and jet (with smaller region) is difficult, because of different underlying variables. Further, a larger region does not necessarily help to get better correlations, as large part of the NAO region is also far way from Europe. Therefore, the argument of the NAO region has the advantage to capture also the European region is not really convincing, as the NAO, as mentioned by the authors, also only includes 2 classifications for the whole region. Apart from this, the jet-region also includes part of western Europe, and as Europe is part of the westwind regime, it is not really surprising that the upstream information has an impact on the immediate downstream weather response. Extending the jet-region eastward would be very interesting to see, if this indeed does substantially increase the reconstruction skill, but I would assume this is not clear. So this point is very interesting, but it was not investigated here. Same is true for the next interesting point (number of classifications for one variable), also very interesting but also not included in this manuscript. The question about the relevant or optimal number of classes for the reconstruction exercise does sound also really interesting. But it is not clear to me, what the example of the 30 MSL classifications tells the reader, as at some point maybe too many classifications could be reducing the skill in a similar way as extreme events could be problematic. The more classifications, the more relevant individual extremes could become, because their impact will be felt more strongly during individual seasons.

lines 325-330:

Same as for lines 305-314 and 315-324. The authors nicely present a way of how their results can be interpreted, where relevant impacts of the method on the results might occur or how specific changes in the classification setup could have an impact. I find this discussion very helpful, but I am somehow missing the part where the authors are highlighting the results of their study. Main issue with this point can be found in my general comment.