

On the occurrence of strong vertical wind shear in the tropopause region: A ten year ERA5 northern hemispheric study

Thorsten Kaluza¹, Daniel Kunkel¹, and Peter Hoor¹

¹Institute for Atmospheric Physics, Johannes-Gutenberg University Mainz, Mainz, Germany

Correspondence: Thorsten Kaluza (kaluzat@uni-mainz.de)

Abstract. A climatology of the occurrence of strong wind shear in the UTLS is presented, which gives rise to define a tropopause shear layer (TSL). Strong wind shear in the tropopause region is of interest because it can generate turbulence which can lead to cross-tropopause mixing. The analysis is based on ten years of daily northern hemispheric ECMWF ERA5 reanalysis data. The vertical extent of the region analysed is limited to the altitudes from 1.5 km above the surface up to 25 km, to exclude the planetary boundary layer as well as strong wind shear in higher atmospheric layers like the mesosphere/lower thermosphere. A threshold value of $S_t^2 = 4 \cdot 10^{-4} \text{ s}^{-2}$ is applied, which marks the top end of the spectrum of atmospheric wind shear to focus on situations which cannot be sustained by the mean static stability in the troposphere according to linear theory. This subset of the vertical wind shear spectrum is analysed for its vertical, geographical, and seasonal occurrence frequency distribution. A set of metrics is defined to narrow down the relation to planetary circulation features, as well as indicators for momentum gradient sharpening mechanisms.

The vertical distribution reveals that strong vertical wind shear above the threshold occurs almost exclusively at tropopause altitudes, within a vertically confined layer of about 1–2 km extent directly above the local lapse rate tropopause (LRT). The TSL emerges as a distinct feature in the tropopause-based 10 year temporal and zonal mean climatology, spanning from the tropics to latitudes around 70° N, with average occurrence frequencies of the order of 1 % – 10 %. The horizontal distribution of the strong vertical wind shear near to the tropopause exhibits distinctly separated regions of occurrence, which are generally associated with jet streams and their seasonality. At midlatitudes, strong wind shear values occur most frequently in regions with an elevated tropopause and at latitudes around 50° N, associated with jet streaks within northward reaching ridges of baroclinic waves. At lower latitudes in the region of the subtropical jet stream, which is mainly apparent over the East Asian continent, the occurrence frequency of strong wind shear near to the tropopause reaches maximum values of about 30 % during winter and is tightly linked to the jet stream seasonality. The interannual variability of the occurrence frequency for strong wind shear might furthermore be linked to the variability of the zonal location and strength of the jet. The east-equatorial region features a bi-annual seasonality in the occurrence frequencies of strong vertical wind shear near to the tropopause. During the summer months, large areas of the tropopause region over the Indian ocean are up to 70 % of the time exposed to strong wind shear, which can be attributed to the emergence of the tropical easterly jet. During winter, this occurrence frequency maximum shifts eastward over the maritime continent, where it is exceptionally pronounced during the DJF 2010/11 la Niña phase, as well as quite weak during the El Niño phases of 2009/10, 2014/15 and 2015/16. This agrees with the atmospheric response of the Pacific Walker circulation cell in the ENSO ocean-atmosphere coupling.

1 Introduction

30 The distribution of vertical wind shear in the atmosphere is a substantial feature of the dynamic structure because it controls the dynamic stability of the flow. Vice versa, dynamic stability limits the amount of wind shear that is sustainable by the flow. The dynamic stability of the tropopause region is ultimately of interest because turbulent mixing in consequence of dynamic instability modifies the trace gas gradients at the tropopause, which in turn can have a significant effect on the radiative budget not only locally but also at the Earth's surface (Forster and Shine, 1997; Riese et al., 2012). Turbulent mixing at the tropopause
35 is furthermore a pathway for stratosphere-troposphere exchange (STE, Holton, 1995; Stohl et al., 2003), which is important because of its impact on the chemical budget in the troposphere and the stratosphere.

According to linear theory, the dynamic stability of a medium in a stratified shear flow can be evaluated on the basis of the non-dimensional Richardson number $Ri = N^2/S^2$. It is defined as the ratio between static stability $N^2 = g/\Theta \cdot (\partial\Theta/\partial z)$ and squared vertical shear of the horizontal wind $S^2 = (\partial u/\partial z)^2 + (\partial v/\partial z)^2$, with the gravitational acceleration g and the
40 zonal and meridional wind components u and v . The static stability N^2 is determined through the vertical gradient of the potential temperature $\Theta = T(p_0/p)^{R_d/c_p}$, with the temperature T , the atmospheric air pressure p and a reference pressure p_0 , and the ratio between the specific gas constant for dry air and the specific heat capacity $R_d/c_p = 0.286$. The Richardson number describes the ratio of the suppression of turbulent kinetic energy due to buoyancy and the production of turbulent kinetic energy due to shear forces. If the flow exhibits Richardson numbers below the critical threshold value $Ri_c = 1/4$, it
45 can become dynamically unstable (Miles, 1961). A common descriptive example for this issue is the occurrence of Kelvin Helmholtz instabilities (KHI), where the shear-induced horizontal shift of a vertically displaced air parcel results in local convective instability in the elsewhere stably stratified flow, and eventually in flow overturning, turbulent breakdown, and gradient erosion.

The lapse rate tropopause (LRT) is defined as the first vertical level above the surface at which the temperature lapse rate
50 falls below 2 K km^{-1} and its mean between this level and any level up to 2 km above remains below that threshold (WMO, 1957). This increase in stratification implicates an increase in static stability, from mean upper tropospheric values on the order of $\overline{N^2}_{trop} \approx 1 \cdot 10^{-4} \text{ s}^{-2}$ to average lower stratospheric values of $\overline{N^2}_{strat} \approx 4 \cdot 10^{-4} \text{ s}^{-2}$. The vertical profile of N^2 furthermore exhibits a vertically confined maximum within the first few kilometres above the LRT which is caused by a localised temperature inversion. This feature is quasi-ubiquitous on planetary and climatological scales, and is referred to as
55 the tropopause inversion layer (TIL, Birner, 2006). The increase in static stability at the LRT in general, as well as the N^2 maximum within the TIL in particular define a background state which can sustain larger vertical wind shear S^2 compared to the troposphere. This fact is reflected in the vertical distribution of S^2 .

Early approaches towards assessing statistical vertical wind shear occurrence frequencies were performed by Dvoskin and Sissenwine (1958), in the context of preparing guidelines for missile design, using a set of radiosonde measurements mostly

60 taken during winter and spring at the East coast of the USA (41° N–45° N). The data showed distinct occurrence frequency maxima of enhanced values of S^2 within sampling windows of 0.9 km at altitudes of about 9–12 km.

The occurrence of strong S^2 in the tropopause region was described in several research studies which analyse vertical profiles of atmospheric flow properties in a vertical coordinate system relative to the LRT altitude (Birner, 2006). In the pioneering work on the existence of the TIL, Birner et al. (2002) describe a sharp peak of S^2 at the LRT in tropopause-based averaged
65 radiosonde profiles from Munich (48.1° N, 11.6° E), Germany. The peak is more pronounced during winter months (DJF) compared to summer (JJA). Another important finding is the rather large discrepancy between the radiosonde data and the ECMWF ERA-Interim reanalysis dataset, which is largely due to the limited vertical resolution of the numerical model. Zhang et al. (2015) expand these results and present a ten year averaged annual cycle of the vertical wind shear in a tropopause-
70 relative vertical coordinate, derived from radiosonde measurements taken at Boise, Idaho (43.6° N, 116.2° W). The meridional dependency of strong vertical wind shear near to the tropopause was analysed by Zhang et al. (2019), based on 13 years of radiosonde measurements at 90 stations in the northern hemisphere. The high latitude stations are located in Alaska and show no strong signal in S^2 throughout the year. The extratropics are covered by stations in the US mainland, where the long year averaged data reveal a sharp S^2 maximum within the first kilometre above the LRT between 20° N and 50° N. The tropics are represented by several stations in the Caribbean as well as in the western Pacific, and exhibit a more pronounced maximum in
75 S^2 over a larger vertical distance above the LRT compared to the extratropics. The tropical S^2 maximum near to the tropopause exhibits a bi-annual seasonality with maximum averaged values during DJF as well as during JJA.

The tropical summer maximum is associated with the emergence of the tropical easterly jet (TEJ) over the Indian Ocean, which is an inherent component of the Asian summer monsoon circulation. Koteswaram (1958) was the first to describe these upper tropospheric easterlies, on the basis of radiosonde measurements at several stations between the equator and 40° N and
80 from 20° W to 150° E, and he already noted exceptionally pronounced vertical wind shear above the wind speed maximum in individual profiles. Sunilkumar et al. (2015) investigated occurrence frequencies for vertical wind shear threshold values in the context of turbulence characteristics over India, based on a 4 year data set of GPS-radiosonde measurements at Trivandrum (8.3° N, 76.6° E) and Gadanki (13.5° N, 79.2° E). The analysis reveals a sharp peak of maximum occurrence frequencies for strong vertical wind shear close to the cold point tropopause (CPT) and particularly during summer.

85 The occurrence of strong vertical wind shear in the tropopause region can exceed the static stability and result in subcritical Richardson numbers, the emergence of dynamic instabilities, wave overturning and turbulent breakdown of the flow. This was the central subject of a variety of observational studies in the context of clear air turbulence (CAT) and STE (e.g. Shapiro, 1976, 1978). Recently, Kunkel et al. (2019) analysed the relation of the TIL and STE (e.g. Gettelman and Wang, 2015) based on a synergistic approach analysing in situ airborne measurements from the WISE field campaign along with ECMWF Integrated
90 Forecast System (IFS) forecasts and idealised numerical baroclinic life cycle experiments. They could show that strong wind shear emerged in regions of enhanced static stability above the local tropopause, i.e., the TIL. In their case this resulted in subcritical Richardson numbers, turbulent mixing and STE. Kaluza et al. (2019) describe a co-occurrence on synoptic scales of enhanced static stability N^2 and strong vertical wind shear S^2 near to the tropopause in breaking baroclinic waves as a generic feature in midlatitudes, based on a composite analysis of operational ECMWF IFS analysis data. The composites show overall

95 reduced Richardson numbers within the lowermost stratosphere (LMS, Holton, 1995) in ridge regions of breaking baroclinic waves. These results suggest that enhanced wind shear is evident close above the local tropopause, at least, in the extratropics, and might lead to the generation of turbulence and subsequent STE. Recently, the high resolution numeric simulation of a midlatitude cyclone which was associated with a large number of turbulence reports gave insight on the importance of the tropospheric jet streak and wind speed and shear enhancement within upper tropospheric outflow of deep convection on the
100 occurrence of CAT, along with the generation of gravity waves on different scales and their interaction with the background wind shear profile at critical levels as well as in regions of subcritical Richardson numbers (Trier et al., 2020).

The previous paragraphs gave an overview of observational evidence of strong vertical wind shear near to the tropopause. A majority of the shear regions described are associated with planetary circulation features, i.e. the polar front jet (PFJ) in midlatitudes, the subtropical jet stream (STJ) over east Asia and the northwest Pacific, and the summer TEJ over the Indian
105 Ocean. The following paragraphs recapitulate processes that influence the occurrence of wind shear, particularly in the upper troposphere/lower stratosphere (UTLS) region. The jet streams constitute the planetary-scale background state for the distribution of wind shear, with vertical shear zones spanning over several kilometres and lateral shear zones of several hundreds of kilometres. The PFJ or eddy-driven jet is associated with baroclinic development at midlatitudes, and can be described in the first approximation as a thermal wind. The upper level front above the level of maximum horizontal wind speed is associated
110 with exceptionally large horizontal temperature gradients, which in turn cause strong vertical wind shear according to the thermal wind relation. The analysis of Endlich and McLean (1965) showed a general agreement between the thermal wind which was calculated based on temperature measurements and a vertically confined layer of strong vertical wind shear derived from wind measurements.

The subtropical jet stream (STJ) is located closer to the equator and generally at higher altitudes compared to the polar jet, and
115 it is driven by the conservation of angular momentum during the poleward excursion of air masses, as well as the temperature gradient at the subtropical tropopause break. The winter STJ over east Asia and the west Pacific exhibits exceptionally high wind speeds (Jaeger and Sprenger, 2007) as well as large occurrence frequencies (Koch et al., 2006), and is also referred to as the east Asian jet stream (EAJS, Yang et al., 2002).

The planetary scale background wind shear which is determined by the occurrence of the jet streams is further modulated
120 by smaller scale processes. Liu (2017) confirm the relevance of processes on a large spectrum of scales through spectral decomposition of strong wind shear features in numerical model data from the NCAR Whole Atmosphere Community Climate Model (WACCM). On the mesoscale, flow deformation, convergence and differential temperature advection can result in frontal zones with the associated wind shear according to the thermal wind relation (Ellrod and Knapp, 1992). Kunkel et al. (2014) showed in numerical baroclinic life cycle simulations, how mesoscale gravity waves with intrinsic frequencies close
125 to the inertial limit (inertia gravity waves, IGW) can deform the flow, resulting in local enhancement of S^2 in the tropopause region. Gravity waves furthermore interact with the tropopause in general and the TIL in particular (Kunkel et al., 2014; Zhang et al., 2015, 2019), since the maximum in N^2 presents a maximum in the refractive index, resulting in wave refraction and a characteristic shift in the frequency spectrum, as well as partial or total wave reflection.

Bense (2019) found the transmission coefficient and the shift of the wave spectrum to exhibit a rather complex dependency on the wavelength, the angle of the wave vector, the strength and depths of the TIL, as well as the background wind profile. A case study in this work furthermore showed how critical level filtering in the tropopause region of orography-induced gravity waves resulted in a pronounced local wind shear enhancement caused by non-linear wave-mean-flow-interaction. The overall relevance of critical level filtering at the tropopause, however, depends on the wave spectrum and the background flow, and remains to be quantified. For example, Spreitzer et al. (2019) analysed the diabatic tendencies of the physical process parametrisations in the ECMWF IFS model during a baroclinic life cycle case study. The tropopause region within the ridge of the baroclinic wave featured regions of pronounced S^2 enhancement, reduced Richardson numbers, and diabatic PV modification. The contribution of the non-resolved gravity wave drag parametrisation, however, was found to be negligible.

The interrelation between gravity waves, strong vertical wind shear and the TIL is particularly of interest because of their common region of occurrence. Ridges of baroclinic waves are often associated with strongly curved jet streaks, jet exit regions, deep convection, and frontal zones. These features are sources for the emission of a large intrinsic frequency spectrum of gravity waves, reaching from the inertial limit up to small scale high frequency waves (Plougonven and Zhang, 2014). These waves propagate upwards towards the tropopause within ridges, which are associated with pronounced TIL enhancement as well as a frequent occurrence of strong vertical wind shear (e.g. Kaluza et al., 2019). IGWs can sharpen the synoptic scale background wind shear associated with the jet stream locally on the mesoscale, potentially resulting in subcritical Richardson numbers, with eventual growth of a small scale perturbations due to dynamic instabilities (Sharman et al., 2012). The evolution of dynamic instabilities, wave growth and overturning, and eventual turbulent breakdown results in the erosion of the gradients responsible for the initial instability, as well as gradient sharpening in the vertically adjacent regions, which can lead to the vertical succession of momentum gradients (Reiter, 1969).

Fritts et al. (2013) and subsequent research by the lead author (Fritts et al., 2016, 2017) analyse the *sheet and layer* structure of the atmosphere, a feature which is closely linked to the occurrence of strong vertical wind shear near to the tropopause. They use a direct numerical simulation (DNS) setup which consists of a large scale vertically propagating gravity wave in a background flow with a superimposed fine-scale structure, defined by a static small-scale wave perturbation. The phase relation between the background and the gravity wave results in regions of turbulence and gradient erosion, with adjacent gradient sharpening, resulting in layers of reduced gradients, and thin sheets of enhanced gradients up to a factor 10 of the background values.

The preceding paragraphs gave an overview of observational and theoretical research studies on the occurrence of strong vertical wind shear S^2 in the tropopause region. To the knowledge of the authors, a comprehensive statistical analysis of this atmospheric feature does not yet exist, particularly on climatological scales. This work presents an approach towards such an analysis, on the basis of ten years of northern hemispheric ECMWF ERA5 reanalysis data. The goal is to present a consistent area-wide analysis of the vertical and geographic occurrence frequency distribution for strong wind shear in a state of the art long term numerical representation of the atmosphere, i.e., the ERA5 reanalysis. Compared to observational research studies our approach has no spatial limitations to assess the occurrence frequency. However, for our analysis it is necessary to keep in

mind that the vertical wind shear features are only as well represented as the model resolution allows them to be. An important
 165 factor in this context is the vertical resolution, which has improved significantly compared to the ERA-Interim reanalysis as
 a reference dataset (e.g., Hoffmann et al., 2019). Another important aspect is the choice of the central analysis method, i.e.,
 the analysis of the occurrence frequency distribution of vertical wind shear above a certain threshold value. This approach
 conserves more information on where strong wind shear occurs in exchange for a loss of information concerning the actual
 strength of the wind shear, in contrast to the more common approach of tropopause based averaging.
 170 Thus, the analysis addresses the following central questions: Where and how frequently is the tropopause region exposed to
 strong vertical wind shear, and how do these shear regions compare to atmospheric flow features and mechanisms that modify
 momentum profiles?

The paper is structured as follows. In Sect. 2 we describe the data set as well as the data processing and general analysis
 175 methods. In Sect. 3 we present an exemplary analysis of a single day to present the issue and explain the metrics, and then
 proceed with the 10 year climatology in Sect. 4. Section 5 discusses the results, and Sect. 6 summarises the mayor outcomes
 and presents a further outlook.

2 Data and methods

For this study we use ten years of daily northern hemispheric ECMWF ERA5 reanalysis fields (Hersbach et al., 2020), from 1
 180 January 2008, 00 UTC, to 31 December 2017, 00 UTC. The reanalysis dataset is based on the Cy41r2 cycle of the ECMWF IFS
 model, and for this study the output is processed on a regular 0.25° latitude-longitude grid, as well as on the 137 native vertical
 hybrid sigma-pressure levels between the surface pressure and 1 Pa. This corresponds to an average vertical grid spacing in the
 UTLS region of about 300–400 m, depending on the elevation of the tropopause. The region analysed is restricted from level
 37 (counting towards the surface, with level 37 corresponding to about 25 km altitude) down to 1.5 km above the orography.
 185 Thus, the planetary boundary layer (PBL) is excluded, as well as regions of enhanced wind shear in the mesosphere/lower
 thermosphere (MLT) region (Liu, 2017).

Basic variables such as the temperature T and the three-dimensional wind (u, v, ω) are directly provided by the ECMWF.
 Based on the temperature profiles, the LRT altitude is determined following the definition of the WMO as the first level where
 the lapse rate falls below 2.0 K km^{-1} with the condition that it remains below this value between this level and all higher levels
 190 within 2 kilometre above. The ten year analysis of tropopause-based wind shear at midlatitudes in Sect. 4.1 furthermore makes
 use of dynamical tropopause fields provided by the ECMWF. Following the definition of Ertel (1942) the potential vorticity
 (PV) can be written as

$$Q = \frac{1}{\rho} \boldsymbol{\eta} \cdot \nabla \Theta, \quad (1)$$




Figure 1. Relative occurrence frequency histogram of static stability N^2 and vertical wind shear S^2 between 1.5 km above the surface up to 25 km altitude, on 11.09.2017 at 00:00 UTC and over the northern hemisphere. Counts are weighted with the grid box volumes. a) Static stability (N^2 , in 10^{-4} s^{-2}). Blue indicates tropospheric volume ($z < z(\text{LRT})$), red stratospheric volume ($z > z(\text{LRT})$). Black dashed lines indicate average tropospheric/stratospheric N^2 . b) Relative occurrence frequency distribution of vertical wind shear (S^2 , in 10^{-4} s^{-2}). Logarithmic occurrence frequency scale. Black dashed line indicates threshold value S_t^2 .

where ρ is the density of the medium, and $\boldsymbol{\eta} = \nabla \times \mathbf{u} + 2\boldsymbol{\Omega}$ the vector of the absolute vorticity with the angular velocity of the earth $\boldsymbol{\Omega}$. In the context of the primitive equations which are solved by the IFS this translates to

$$Q = \frac{1}{\rho} (f\mathbf{k} + \nabla \times \mathbf{u}_h) \cdot \nabla \Theta, \quad (2)$$

where $f = 2\Omega \sin(\phi)$ is the Coriolis parameter which represents the component of $\boldsymbol{\Omega}$ in $\mathbf{k}$ direction of the local rectangular coordinate system, and $\mathbf{u}_h$ is the vector of the horizontal wind. The unit for the PV is the 'potential vorticity unit' (pvu), with $1 \text{ pvu} = 1 \cdot 10^{-6} \text{ m}^2 \text{ s}^{-1} \text{ K kg}^{-1}$. The tropopause region is characterised by pronounced isentropic gradients of the potential vorticity, and the dynamic tropopause in the extratropics is commonly defined by a constant value of $Q = 2 \text{ pvu}$ (Hoskins et al., 1985).

The vertical wind shear S^2 is calculated on half levels of the native vertical hybrid sigma-pressure level of the IFS, to retain a maximum amount of information in the gradient based measure. The altitude at each model level is derived from the geopotential after vertically integrating the hydrostatic equation from the pressure and temperature profiles (for further information refer to the IFS documentation, ECMWF (2016)). The increasing vertical grid spacing with increasing altitude in the native coordinates results in a bias towards a larger resolved spectrum of vertical wind shear at lower altitudes, which should be considered. However, the analysis focusses on the UTLS region where the vertical grid spacing increases from about 300 m at 5 km altitude up to about 400 m at 20 km altitude (e.g., Hoffmann et al., 2019). Thus, the resolution bias should not have a large impact on the results presented. The central goal of this study is to quantify the occurrence of strong vertical wind shear S^2 . For this, we define a threshold value $S_t^2 = 4 \cdot 10^{-4} \text{ s}^{-2}$, marking the top end of the spectrum of




Figure 2. Potential temperature at the LRT ($\Theta(\text{LRT})$, in K) over the northern hemisphere, on 11 September 2017 at 00:00 UTC. Black lines show contours of the horizontal wind at 200 hPa, beginning at 30 m s^{-1} in steps of 15 m s^{-1} . Magenta shaded areas indicate regions $S^2 \geq S_t^2$ between 1 km below and 2 km above the local LRT. Dashed white lines indicate the location of the vertical cross sections in Fig. 3.

atmospheric vertical wind shear. The threshold value is selected based on the consideration that $S^2 \geq S_t^2$ generally cannot be sustained by the average tropospheric static stability $\overline{N^2}_{trop.}$, thus leading to low Richardson numbers and conditions favourable to turbulence. In contrast, average stratospheric static stability values of $\overline{N^2}_{strat.} = 4 \cdot 10^{-4} \text{ s}^{-2}$ lead to Richardson numbers of the order of magnitude of $\mathcal{O}(1)$. Figure 1 illustrates the previous consideration, for one exemplary time step on which we will also focus in Sect. 3. The distribution of tropospheric static stability peaks around $N^2 = 1 \cdot 10^{-4} \text{ s}^{-2}$, with an overall average of $\overline{N^2}_{trop.} = 1.33 \cdot 10^{-4} \text{ s}^{-2}$. The stratospheric static stability distribution amounts to an average value of $\overline{N^2}_{strat.} = 5.20 \cdot 10^{-4} \text{ s}^{-2}$, which is shifted towards a larger value due to the above-average static stability which defines the TIL. The average stratospheric static stability above the TIL is generally closer to the value of $\overline{N^2}_{strat.} = 4 \cdot 10^{-4} \text{ s}^{-2}$ (e.g. Birner et al., 2002). The choice of the threshold S_t^2 is furthermore motivated by previous research studies which indicate the mostly exclusive occurrence of vertical wind shear exceeding S_t^2 at tropopause altitudes (e.g. Kunkel et al., 2019).

3 Identification of a shear layer on 11 September, 2017

We will start our analysis by initially focussing on a specific time step in September 2017. This study was performed in the context of the airborne research campaign WISE that took place during SON 2017 over the North Atlantic, which motivated the choice of the date. Our intention is to introduce our metrics first and then look at longer time periods afterwards. We start with the synoptic situation to provide the large scale overview. Figure 2 shows a snapshot of the northern hemispheric potential temperature at LRT altitude on 11 September, 2017, along with maxima of the horizontal wind at 200 hPa. The primary feature standing out is the tropopause break and the associated jet streaks of the horizontal wind. Over the Asian continent and reaching to the western Pacific, the tropopause break features a sharp meridional $\Theta(\text{LRT})$ gradient with a single coherent STJ. Further west, the tropopause break is less sharp, and features a characteristic sequence of Rossby wave patterns accompanied by individual jet streaks at varying latitudes.




Figure 3. Vertical cross sections on 11 September 2017, at 00 UTC and at a) 120° W, b) 60° W, and c) 100° E. Colour contour shows zonal wind speed (u , in m s^{-1}), black dots LRT altitude, and black dotted lines isentropes (Θ , in K). Magenta lines show $S^2 = S_t^2$ isolines.

The synoptic scale wind systems are further illustrated in the vertical cross sections in Fig. 3. At 120° W (Fig. 3a), a pronounced polar jet maximum is visible at 50 – 60° N, which is also evident in the vertically integrated horizontal wind in Fig. 2. Further southward, the tropopause break at 30° N features comparatively weak westerly winds, followed by high altitude easterlies in the tropics. The northwest Atlantic (Fig. 3b) exhibits two distinct zonal westwind maxima, which can be
235 interpreted as the jet exit region of the STJ and the entry region of a jet streak of the PFJ. The cross section at 100° E (Fig. 3c) again shows a sequence of zonal winds, with high altitude easterlies in the tropics, followed by the STJ, and further northward the rotational components of a cyclonic system over Siberia.

We proceed with the analysis of the vertical and geographical distribution of strong vertical wind shear exceeding the threshold value $S_t^2 = 4 \cdot 10^{-4} \text{ s}^{-2}$. Figure 4 shows the zonally averaged relative occurrence frequency of strong vertical wind shear
240 $S^2 \geq S_t^2$ on a logarithmic color scale, for the northern hemisphere on 11 September 2017. Figure 4a reveals three meridional regions of exceptional occurrence frequencies located in the UTLS region, one in the tropics, one at 30 – 40° N, and a third one at midlatitudes. Rearranging the grid volumes in a tropopause-relative vertical coordinate system depending on their vertical distance from the LRT (Fig. 4b and c) concentrates the maxima of strong wind shear occurrence in all three regions to a vertical layer of about 1–2 km extent directly above the LRT. Secondary occurrence frequency maxima can be identified closely below
245 the tropopause in all three meridional regions.

The pronounced occurrence of $S^2 \geq S_t^2$ in a distinct layer close to the LRT motivates to proceed with the analysis focussing on the question *"Is there a global relation between vertical wind shear and the tropopause?"*. To answer this question, we choose a binary criterion where we define the tropopause region as being exposed to strong vertical wind shear if S^2 exceeds S_t^2 at least in one grid box volume between 1 km below and 2 km above the LRT. These regions are indicated in Fig. 2. The
250 wind systems described in the first paragraph of this section can now be associated with the three meridional regions of frequent occurrence of strong vertical wind shear near to the tropopause in Fig. 4. At midlatitudes, the tropopause region is exposed to strong vertical wind shear within the Rossby wave jet streaks, as well as in the cyclonically curled up ridges associated with breaking baroclinic waves (over Canada, Scandinavia and Siberia) (Kaluza et al., 2019). Further southward, the STJ also




Figure 4. Relative occurrence frequency of $S^2 \geq S_t^2$, zonally averaged over the northern hemisphere, on 11 September 2017, 00 UTC. Logarithmic frequency contour, vertically binned in $dz = 500$ m. a) Geometric altitude as the vertical coordinate. Dashed black line indicates the effect of the 1.5 km above orography cut-off. b) Relative distance from the LRT as vertical coordinate. Dashed black line shows LRT altitude. c) As in b) but with mean LRT altitude for profiles with $S^2 \geq S_t^2$ restored (dashed bold black line).

features strong vertical wind shear at the tropopause, particularly in the jet exit region over the northwest Pacific. The tropical easterlies over the Indian Ocean and the east Pacific feature large scale coherent regions of strong vertical wind shear close to the tropopause. Several smaller scale wind shear regions can be attributed to individual atmospheric wind systems, e.g. the late stage ex-hurricane Irma which is located over Florida, or the stratospheric cut-off over the Mediterranean Sea.

4 Ten year climatology of strong wind shear in the northern hemisphere UTLS

4.1 Vertical distribution and seasonality of strong vertical wind shear in the northern hemisphere

We repeat the analysis steps from Sect. 3 for the whole dataset of ten years of daily ERA5 fields. We first present the overall vertical distribution of strong vertical wind shear $S^2 \geq S_t^2$, followed by the geographic occurrence frequency and seasonality in the tropopause region. Figure 5a shows the 10 year temporal and zonal average occurrence frequency for strong vertical wind shear $S^2 \geq S_t^2$, with the geometric altitude as the vertical coordinate. The mean LRT altitude for the same time period and region is indicated by the black solid line. Distinct occurrence frequency maxima are apparent, i.e., in the midlatitudes between 40° – 60° N and mainly above the LRT, at the tropopause break at about 30° N above and below the LRT, and in the tropics in close vicinity above and below the LRT. The rearrangement of the grid boxes in the LRT-relative vertical coordinate system (Fig. 5b) concentrates the occurrence frequency maxima in a distinct layer above the LRT. This layer spans from the tropics to latitudes north of 60° N, and it exhibits occurrence frequencies of the order of 1 %–10 % over a vertical range of about 1–2 km.




Figure 5. Northern hemispheric occurrence frequency distribution of grid volumes that exhibit strong vertical wind shear $S^2 \geq S_t^2$, from 1 January 2008 to 31 December 2017. Logarithmic frequency contour, vertically binned with $dz = 500$ m. a) Geometric altitude as the vertical coordinate. Solid bold black line indicates mean LRT altitude for all 10 years and the whole northern hemisphere. Dashed thin black line indicates the effect of the 1.5 km above orography cut-off. White solid (dotted) line indicates regions where negative (positive) vertical wind shear makes for 75 % of the counts. b) As in panel a, with LRT-relative vertical coordinate and with mean LRT altitude for profiles with $S^2 \geq S_t^2$ restored (dashed bold black line). Solid bold black line as in panel a. c) As in panel b but from $0^\circ - 20^\circ$ N with the cold point tropopause (CPT) as a reference altitude, and north of 20° N with the dynamic tropopause ($Q = 2$ pvu) as a reference altitude.

270 In the following we will refer to the feature of maximum occurrence frequencies at the LRT in the zonal and/or temporal mean as a *tropopause shear layer (TSL)*, however, a comparison with the TIL should be made cautiously. Both features appear similarly in tropopause-relative zonal means (compare with Zhang et al. (2019)), however, the wind shear layer emerges less frequently as well as less area-wide. Furthermore, a different metric is applied here, which analyses the tropopause-relative occurrence frequency of a threshold value S_t^2 , instead of directly averaging S^2 . We refrain from referring to individual regions

275 of strong wind shear at individual time steps as a tropopause wind shear layer, because this would raise the question of a lower limit for the horizontal and temporal scales that mark such a layer, considering the pronounced mesoscale variability of these regions. We realise that this choice is controvertible, because on a mesoscale horizontal extent of the order of 100 km the geometric aspect ratio between horizontal and vertical extent still clearly describes a layer-like character.

280 At midlatitudes, the TSL is composed of mostly decreasing wind speeds with height. Northward of about 45° N, the profiles which exhibit $S^2 \geq S_t^2$ are on average associated with about 1 km more elevated LRT altitudes compared to the overall zonal average (dashed and solid black lines in Fig. 5b). This indicates the importance of large scale wave dynamics, particularly the polar jet streaks within ridges of baroclinic waves, which are known to exhibit exceptional wind shear in the tropopause region




Figure 6. Schematic vertical cross section of the tropopause-based averaging method. Left part shows exemplary situation at the tropopause break, right part the key measures after tropopause-based averaging. Black lines indicate the LRT, grey lines the dynamic tropopause. Blue lines show isotachs, and the yellow and blue regions indicate regions of enhanced vertical wind shear above and below the jet core.

(Kaluza et al., 2019; Kunkel et al., 2019). The above-average LRT altitudes for profiles with $S^2 \geq S_t^2$ are a unique feature for the higher midlatitudes, as they do not occur in the tropics or the subtropics, which will be further addressed in Sect. 4.2.

At the tropopause break, the TSL is more evenly composed of both decreasing and increasing wind speed with height. This is due to the contribution of enhanced vertical wind shear at the upper edge of the tropopause break and above the jet core, as well as at the lower edge and below the jet core. Furthermore, an enhanced vertical spread of $S^2 \geq S_t^2$ is apparent at the tropopause break in the LRT-relative vertical coordinates (Fig. 5b) compared to the geometric altitude (Fig. 5a). Figure 6 illustrates how this can be explained by the averaging method. Below the jet core of the STJ strong vertical wind shear occurs frequently within PV streamers (Škerlak et al., 2015), i.e., tongues of stratospheric air that reach south- and downward. They are characterised by stratospheric PV values, which is linked to the frontal character of the inherent temperature gradients. The horizontal temperature gradients are associated with a strong thermal wind forcing, and the resulting vertical wind shear can be sustained by the vertical temperature gradients. Despite the stratospheric characteristics of these air masses, the LRT criterion which requires a mean lapse rate below 2.0 K km^{-1} over a vertical distance of 2 km is no longer met at some point within the PV streamers, which is indicated by the dashed black line in Fig. 6. In these cases the LRT is located at the upper edge the tropopause break and several kilometers above the region of strong wind shear. Eventually, the 10 year temporally and zonally averaged LRT altitude exhibits a more smooth transition between the lower and the upper edge of the tropopause break (right panel in Fig. 6), which is caused by short-period as well as seasonal meandering of the tropopause break. The LRT-relative averaging method ultimately shifts the region of strong wind shear downward from the mean LRT altitude, over the distance

which was originally determined relative to the instantaneous LRT. The equivalent effect also occurs at the upper edge of the tropopause break as indicated in Fig. 6.

The tropical region in Fig. 5b features the most pronounced occurrence frequency maximum, exhibiting a larger vertical spread compared to higher latitudes, as well as a distinct secondary maximum below the LRT. The composition of positive and negative vertical wind shear in the tropics exhibits a layered structure, particularly above the LRT, where it is influenced by the quasi-biennial oscillation (QBO) phase and the elevation of the level of vanishing horizontal wind.

The significance of the processes which result in the occurrence of the TSL remains to be quantified. The clustering of grid volumes which exhibit $S^2 \geq S_t^2$ directly above the LRT in Fig. 5b agrees with the dynamic stability criterion and the thermal wind shear forcing associated with upper tropospheric fronts. However, the overall link between the tropopause definition, which is governed by the temperature profile, and the occurrence of strong vertical wind shear remains uncertain. Therefore, we repeat the analysis for the tropical cold point tropopause (CPT), i.e., the absolute temperature minimum in the tropical UTLS, and for the dynamic tropopause, i.e., the $Q = 2$ pvu isosurface in the extratropics. The tropics feature a distinct separation of up to 1 km between the LRT and the CPT (Seidel et al., 2001), which motivates the comparison at low latitudes. The PV on the other hand does not constitute a useful tropopause definition in the tropics (Holton, 1995), which is why the dynamic tropopause is only used in the extratropics in this study. From Fig. 5c it is evident that ERA5 resolves the separation of the LRT and the CPT in the tropics, along with central features like a decreasing mean distance between the two tropopauses towards the equator (Seidel et al., 2001). The clustering of strong wind shear grid volumes above the CPT is less pronounced compared to the LRT, along with a more pronounced secondary maximum below the CPT, which indicates that the occurrence of strong vertical wind shear is more closely linked to the LRT in the tropics. In the extratropics, the analysis in a vertical coordinate system relative to the dynamic tropopause (Fig. 5c and north of 20° N) compares well with the analysis in the LRT-relative vertical coordinates. At high latitudes, the profiles which exhibit strong vertical wind shear are associated with above average dynamic tropopause altitudes. However, the clustering of grid volumes with $S^2 \geq S_t^2$ above the dynamic tropopause exhibits a larger vertical spread, particularly above the tropopause break. The dynamic tropopause is identified systematically below the LRT in this region, due to the inclusion of PV streamers, as indicated in Fig. 6. Therefore, a larger amount of grid volumes which exhibit strong wind shear are shifted upwards during the averaging process. In conclusion, the averaging of the grid volumes which exhibit $S^2 \geq S_t^2$ based on the distance from the different reference altitudes reveals that the elevation of strong vertical wind shear is (not generally but in the overall mean) more closely linked to the LRT than to the CPT or the dynamic tropopause.

We proceed with the analysis of the geographical spread of wind shear regions which are located between 1 km below and 2 km above the local LRT. For that, we calculate occurrence frequencies for how often the binary criterion for strong vertical wind shear close to the LRT which was defined in Sect. 3 is met. Figure 7 shows the seasonality of these quasi-horizontal occurrence frequencies over the northern hemisphere. Throughout all seasons, a clear separation of occurrence frequency maxima is apparent, where each region can be attributed to a planetary circulation feature. The Northeast Pacific and the North Atlantic each exhibit a distinct maximum spanning over the storm track regions (Shaw et al., 2016). These maxima are most pronounced during winter and spring, exhibiting maximum occurrence frequencies of 25 – 30 %.




Figure 7. Occurrence frequency distribution of $S^2 \geq S_t^2$ between 1 km below and 2 km above the local LRT and in the northern hemisphere. Averaged over 10 years from 2008 to 2017 for a) DJF, b) MAM, c) JJA and d) SON. Red solid lines indicate isolines of the horizontal wind at 200 hPa, starting at 30 m s^{-1} in steps of 30 m s^{-1} .

Another region with maximum occurrence frequencies of up to 45 % is apparent over Asia and the Northwest Pacific during winter, which is associated with the STJ. The strength of the maximum throughout the year shows a general agreement with the seasonality of the jet stream. The tropics feature several maxima spanning around the equator, with one particular wind shear region located over the Indian Ocean which exhibits maximum occurrence frequencies up to 75 % during JJA. This maximum

340 shifts eastward during autumn and winter, where it is located centred over the maritime continent. The summer maximum can be attributed to the emergence of the TEJ during these months associated with the Asian summer monsoon circulation. The winter maximum is linked to the Pacific Walker circulation and the El Niño Southern Oscillation (ENSO) ocean-atmosphere coupling. This will be further addressed in Sect. 4.3.

The following subsections each focus on one of the regions where strong vertical wind shear occurs frequently at tropopause
 345 altitudes, beginning at high latitudes and moving towards the equator. The goal is to narrow down the relation to planetary circulation features, and to investigate formation mechanisms of the TSL in the dataset.

4.2 The TSL related to Rossby waves in the mid latitudes

At midlatitudes, the occurrence of enhanced tropopause-based vertical wind shear is linked to the jet streaks of the polar or eddy-driven jet, and therefore to the associated baroclinic wave patterns. Figure 5b and c indicated above-average lapse rate-
 350 and dynamic tropopause altitudes for profiles exposed to $S^2 \geq S_t^2$, which hints towards the role of the ridges of baroclinic waves. This behaviour agrees with the conclusions from the process study of Kunkel et al. (2019) as well as the composite study of baroclinic waves of Kaluza et al. (2019). We expect the ERA5 reanalysis to resolve vertical wind shear features in the UTLS similarly well compared to the operational IFS analysis data used in Kaluza et al. (2019). However, a direct comparison should be made carefully. The ERA5 reanalysis is based on an IFS version with a T_L639 spectral truncation (about 31 km
 355 horizontal grid spacing, Hersbach et al. (2020)) and 137 vertical level, compared to the T_L1279 spectral truncation (about 16 km horizontal grid spacing) and 91 respectively 137 vertical level in the operational analysis data used in Kaluza et al. (2019). The following subsection presents a more detailed analysis of the dependency of enhanced vertical wind shear on the tropopause altitude.

We begin by selecting two zonal regions encompassing the occurrence frequency maxima in the midlatitudes, one over the
 360 North Atlantic from 80° W to 0° W, and a second one over the Northeast Pacific from 180° W to 120° W. The latter selection is made in such a way to exclude the STJ maximum to the west. We proceed with the $Q = 2$ pvu isosurface fields provided by the ECMWF, and make use of the conservation property of the potential temperature Θ on such an isosurface under adiabatic and frictionless conditions (Hoskins, 1991). We can therefore identify Rossby waves through anomalies of $\Theta(Q = 2 \text{ pvu})$, with positive anomalies indicating a ridge of a baroclinic wave, i.e., subtropical air masses with high tropopause altitudes reaching
 365 poleward and negative anomalies indicating a trough, i.e., polar/subpolar air masses with low tropopause altitudes reaching equatorwards.

We proceed by defining a background state from which we can identify $\Theta(Q = 2 \text{ pvu})$ anomalies. Figure 8a and b show the ten year temporal and zonal average of the potential temperature $\bar{\Theta}(Q = 2 \text{ pvu})$, for both latitudinal regions and subdivided into DJF, MAM, JJA and SON. On average, the dynamic tropopause exhibits potential temperatures of about 360 K at 20° N,
 370 and decreases to values between 300 – 320 K in the polar region. The meridional $\bar{\Theta}(Q = 2 \text{ pvu})$ gradient is most pronounced during DJF and MAM in both regions. We preserve the meridional dependency and define the zonal and temporal average $\bar{\Theta}(Q = 2 \text{ pvu})$ in the meridional region from 35° N– 60° N as the background state for the following analysis. According to Fig. 7, this region encompasses the occurrence frequency maxima over the North Atlantic and the Northeast Pacific. Furthermore,




Figure 8. a) and b): Seasonal ten year temporally and zonally averaged dynamic tropopause potential temperature ($\bar{\Theta}(Q = 2 \text{ pvu})$, in K). c) and d): Seasonal ten year temporally and zonally averaged LRT potential temperature ($\bar{\Theta}(\text{LRT})$, in K). Solid lines show mean values, dashed lines interseasonal standard deviation. Left column North Atlantic region ($80^\circ \text{ W}-0^\circ \text{ W}$). Right column Northeast Pacific region ($180^\circ \text{ W}-120^\circ \text{ W}$). The black dot in panel a indicates $\bar{\Theta}(Q = 2 \text{ pvu})$ at 51° N where Fig. 9 is located. Grey dashed vertical lines in panel a and b indicate the borders of the region that is further analysed.

Fig. 8a and b show that the interseasonal standard deviation of $\bar{\Theta}(Q = 2 \text{ pvu})$ is generally small in this region, which indicates that the ten year seasonal average describes representative average potential temperatures on the dynamic tropopause for the individual years. Figure 8c and d show for the zonal mean that the LRT exhibits averaged potential temperatures close to the ones of the dynamic tropopause, particularly in the region of interest from $35^\circ \text{ N}-60^\circ \text{ N}$. We present this comparison because we make use of both tropopause definitions. On one hand, we analyse the occurrence of $S^2 \geq S_t^2$ relative to the LRT, because the analysis in Sect. 4.1. showed that strong wind shear is more closely linked to the LRT than to the dynamic tropopause. On the other hand, we identify UTLS wave features on the basis of the dynamic tropopause, because of the conservation property of the PV.

We proceed by calculating the distribution of instantaneous deviations from $\bar{\Theta}(Q = 2 \text{ pvu})$ during the ten years, for each region and season. Figure 9 exemplarily shows the result for the North Atlantic region, during DJF and at 51° N . The $\Theta(Q = 2 \text{ pvu})$ values are unimodally distributed around the ten year average background value of $\bar{\Theta}(Q = 2 \text{ pvu}) = 313.6 \text{ K}$. Vertical wind shear exceeding S_t^2 in the vicinity of the LRT however occurs almost exclusively at above-average $\Theta(Q = 2 \text{ pvu})$




Figure 9. Occurrence frequency of $S^2 \geq S_t^2$ within 1 km vertical distance from the LRT, depending on $\Theta(Q = 2 \text{ pvu})$. Blue histogram shows occurrence frequency distribution of $\Theta(Q = 2 \text{ pvu})$ at 51° N during DJF 2008–2017 and averaged over $80^\circ \text{ W} - 0^\circ \text{ W}$. $\Theta(Q = 2 \text{ pvu})$ in 1 K bins. Dashed black line shows mean value $\bar{\Theta}(Q = 2 \text{ pvu}) = 313.6 \text{ K}$ (black dot in Fig. 8a). Solid black line shows occurrence frequency for $S^2 \geq S_t^2$ between 1 km below and 2 km above the local LRT and within the $\Theta(Q = 2 \text{ pvu})$ bins.

values. The occurrence frequencies for $S^2 \geq S_t^2$ increase up to 55 % with increasing $\Theta(Q = 2 \text{ pvu})$, up to the point where the occurrence frequencies for the potential temperatures become statistically insignificant.

Figure 10 presents the extension of the previous analysis for both regions, all seasons, and over the whole selected meridional range. Figure 10a puts the result from Fig. 9 into a larger context, and reveals that $S^2 \geq S_t^2$ in the vicinity of the LRT is generally
 390 associated with positive $\Delta\Theta(Q = 2 \text{ pvu})$ anomalies, i.e. dynamic tropopause potential temperatures above the climatological mean, i.e. primarily ridge regions. The occurrence frequencies increase with increasing potential temperatures and at higher latitudes, with peak values exceeding 50 % at about 50° N and $\Delta\Theta(Q = 2 \text{ pvu}) \approx 20 \text{ K}$.

The general dependency for the occurrence frequency of $S^2 \geq S_t^2$ in the latitude- $\Delta\Theta(Q = 2 \text{ pvu})$ -coordinates holds true for both regions and all four seasons, with overall reduced values towards the summer months, accompanied by a northward
 395 shift. In the Northeast Pacific region and particularly during DJF, the influence of the STJ is visible, causing the bimodal distribution of $\Delta\Theta(Q = 2 \text{ pvu})$ at low latitudes. This is due to occasional northeastward excursions of the STJ and the subtropical tropopause break into the region of interest, which adds exceptionally high dynamic tropopause potential temperatures to the statistic and shifts the overall average towards larger $\bar{\Theta}(Q = 2 \text{ pvu})$ values.

The following consideration is made to put the results of the previous paragraphs in context. From a static point of view
 400 and according to Fig. 7a, maximum occurrence frequencies for $S^2 \geq S_t^2$ in the vicinity of the LRT over the North Atlantic and during DJF are located at about 50° N and reach up to values of 30 – 35 %. From a dynamic point of view, the occurrence can be further narrowed down, with occurrence frequencies exceeding 45 % within any elevated tropopause between $80^\circ \text{ W} - 0^\circ$




Figure 10. Occurrence frequency of $\Delta\Theta(Q = 2 \text{ pvu})$ deviations from $\bar{\Theta}(Q = 2 \text{ pvu})$ as in Fig. 8c, with meridional dependency. Dashed (solid) red lines indicate occurrence frequencies isolines, beginning at 0.5 % (1.0 %) and in steps of 1.0 %. Colour contour shows occurrence frequency for $S^2 \geq S_t^2$ between 1 km below and 2 km above the local LRT and within the $\Theta(Q = 2 \text{ pvu})$ bins. a) to d) DJF MAM JJA SON over the North Atlantic region. e) to h) Same for the Northeast Pacific region. Blue dashed line in a) indicates location of Fig. 8c

W and 45° N – 55° N that exhibits $\Delta\Theta(Q = 2 \text{ pvu}) \gtrsim 20 \text{ K}$. The northward excursion of air masses exhibiting above average tropopause altitudes within ridges of baroclinic waves and the associated jet streaks can be expected to be mainly responsible
405 for this correlation.

Overall, the analysis presents one way to quantify the link between the occurrence of strong vertical wind shear near to the tropopause and the potential temperature of the dynamic tropopause, which can be interpreted as a measure for the excursion of air masses within baroclinic waves in the midlatitudes.

4.3 The TSL around the subtropical jet stream and in the tropics

410 4.3.1 The east Asian jet stream

The tropopause region over east Asia and the northwest Pacific features a maximum in occurrence frequencies for $S^2 \geq S_t^2$, which is associated with the STJ (Fig. 7). The STJ in this region is most pronounced during DJF, and is also referred to as the east Asian jet stream (EAJS). Figure 11 shows the zonal and ten year seasonal average of the potential temperature Θ at the LRT for DJF and JJA, as well as the associated occurrence frequency of $S^2 \geq S_t^2$ between 1 km below and 2 km above
415 the local LRT. The meridional course of $\Theta(\text{LRT})$ agrees well with the climatology derived from radiosonde measurements by




Figure 11. Vertical cross sections of ten year average tropopause properties between 60° E and 180° E longitude: a) winter (DJF) and b) summer (JJA). Upper panels: Solid black line shows $\overline{\Theta}(\text{LRT})$ for the zonal region. Blue solid line shows zonally averaged occurrence frequency for $S^2 \geq S_t^2$ between 1 km below and 2 km above the local LRT, and dashed solid line depicts $\overline{\Theta}(\text{LRT})$ for such vertical profiles. Bottom panels show interannual standard deviation for every measure. The standard deviation of the occurrence frequency displays absolute percentage values.

Seidel et al. (2001). During winter, the zonally averaged occurrence frequency maximum for $S^2 \geq S_t^2$ is located right at the meridional location of the tropopause break, and exhibits values of up to 25 % at around 30° N. The strength of the occurrence frequency maximum exhibits a pronounced interseasonal standard deviation of 5 % (absolute values), which can be partly linked to variability features of the EAJS. The meridional location of the occurrence frequency maximum is rather sharply defined, in agreement with the stable meridional location of the EAJS core on interannual time scales (Yang et al., 2002). The zonal location of the jet core however exhibits a comparatively large interannual variability (Wu and Sun, 2017), which can be linked to the interseasonal variability of the occurrence frequency of $S^2 \geq S_t^2$ in Fig. 11a. In the time period analysed by Wu and Sun (2017) that intersects with our analysis, they find the EAJS core to be located comparatively far westward during the winter seasons of 2008, 2011 and 2012, accompanied by large maximum EAJS core wind speeds. According to Fig. 12a, these years feature exceptionally large occurrence frequency maxima centred within the zonal region from 60° E–180° E. In contrast, the 2009 and 2010 EAJS core is located further eastward and exhibits lower maximum wind speeds, along with reduced occurrence frequencies for $S^2 \geq S_t^2$ (Fig. 12b). Note that the zonal location of the jet core is not well reflected in the multiannual mean of the horizontal wind at 200 hPa. The link between the occurrence of strong vertical wind shear near to the tropopause and the location and strength of the jet core should not be generalised and needs further investigation, since e.g. Wu and Sun (2017) do not see a strong correlation between the zonal EAJS core location and maximum wind speeds. Furthermore, it should be kept in mind that the occurrence of $S^2 \geq S_t^2$ in the vicinity of the LRT is not necessarily linked to exceptional jet core wind speeds.




Figure 12. As in Fig. 7a for a smaller geographical section, averaged over selected DJF seasons of a) west-located EAJS core years (2008, 2011 and 2012), and b) normal located EAJS core years (2009 and 2010). For further details see text and Wu and Sun (2017).

Towards summer, the jet stream slows down and shifts to the north of the upper-tropospheric Tibetan high pressure system associated with the East Asian summer monsoon (EASM) circulation. The zonally averaged occurrence frequency maximum
 435 for $S^2 \geq S_t^2$ in the vicinity of the LRT during JJA decreases to 11 %, and is centred at the upper edge of the tropopause break (Fig. 11b).

4.3.2 The tropical easterly jet

In the tropics, the summer maximum over the Indian Ocean (Fig. 7c) is associated with the tropical easterly jet (TEJ). These easterlies generally arise from June to September as part of the EASM circulation at upper-tropospheric pressure levels around
 440 150 hPa (Krishnamurti and Bhalme, 1976). They exhibit a limited vertical extent and maximum wind speeds around 40 ms^{-1} , which is low e.g. compared to the EAJS core speeds identified by Wu and Sun (2017)). Vertical confinement and limited wind speed have an opposing effect on the vertical wind shear. Nevertheless, this results in the exceptional occurrence frequencies for $S^2 \geq S_t^2$ larger than 70 % in the vicinity of the the LRT over the Indian ocean (Fig. 7c). Figure 11b shows the zonally averaged occurrence frequency for $S^2 \geq S_t^2$ close to the LRT encompassing the TEJ summer maximum. It exhibits a significant
 445 interannual variability, a feature also seen in the vertical shear of the zonal wind found by Roja Raman et al. (2009) at different heights above the TEJ core using MST radar and GPS radiosonde measurements. Furthermore, the occurrence frequencies for strong vertical wind shear which were derived from radiosonde measurements by Sunilkumar et al. (2015) agree qualitatively with the ones at the respective geographic location of each radiosonde station in Fig. 7c. The authors analysed among others occurrence frequencies for vertical wind shear threshold values, however, not in a tropopause-relative coordinate system,
 450 and with gradients calculated for $dz = 20$ meter (implicating a larger spectrum of resolved wind shears), which should be considered when comparing the results. The radiosonde data revealed occurrence frequencies during the monsoon season (June–September) for $S^2 > 2.25 \times 10^{-4} \text{ s}^{-2}$ of almost 80 % at Trivandrum (8.3° N , 76.6° E) and 67 % at Gadanki (13.5° N , 79.2° E), and for $S^2 > 9.0 \times 10^{-4} \text{ s}^{-2}$ occurrence frequencies of 37 % at Trivandrum and 15 % at Gadanki. These peak values occur as sharp maxima above the convective tropopause (COT) and close to the cold point tropopause (CPT), where the LRT
 455 is generally located in between (Sunilkumar et al., 2013).




Figure 13. a) Comparison of the Oceanic Niño sea surface temperature anomaly Index for DJF (ONI, blue graph) with the strong shear occurrence frequency during DJF averaged over the region from 100° E to 130° E and from the equator to 10° N (red graph). The values for January–February 2008 and December 2017 are included (light red). ONI values from <https://www.noaa.gov/>. b) and c) Temporal mean of a zonal cross section at the equator, averaged in geometric height coordinates, for the two consecutive DJF seasons 2009/10 and 2010/11. Color contour shows mean zonal wind component (u , in m s^{-1}). Red solid lines show the 10 % and the 50 % isolines of occurrence frequencies for $S^2 \geq S_t^2$. Blue solid lines show the 10 % and the 50 % isolines of occurrence frequencies for $N^2 \geq 4 \cdot 10^{-4} \text{ s}^{-2}$. Black circle markers indicate LRT altitude, and black solid lines show isolines of potential temperature (Θ , in K). Light blue dashed lines indicate the isolines of 70 % and 85 % relative humidity over ice.

4.3.3 The winter Pacific Walker circulation cell

During the winter months, the tropical tropopause region features another pronounced occurrence frequency maximum for $S^2 \geq S_t^2$ eastward of the summer TEJ maximum and centred over maritime continent (Fig. 7a). It is associated with localised upper tropospheric easterlies, and exhibits a smaller zonal extent as well as about 5 – 10 % lower maximum occurrence frequencies in the ten year average compared to the summer TEJ maximum.

The pronounced interannual variability in the zonally averaged occurrence frequencies for $S^2 \geq S_t^2$ (Fig. 11a), as well as the geographic location of the maximum and the season of its occurrence indicates a link to the El Niño Southern Oscillation (ENSO) ocean–atmosphere coupling and the effect on the Pacific Walker circulation cell. To test this hypothesis, we select a

region encompassing the shear occurrence frequency maximum, from 100° E to 130° E and from the equator to 10° N, and
 465 calculate the average occurrence frequency of $S^2 \geq S_t^2$ close to the LRT in this region during DJF and for the individual years.
 The comparison of these time- and area-averaged frequencies with the Oceanic Niño sea surface temperature anomaly Index
 values for DJF (ONI) show an anticorrelation (Fig. 13a), with a Pearson correlation coefficient of $r = -0.788$. The analysis
 reveals moderate to large occurrence frequencies during neutral and La Niña conditions, and a peak during DJF 2010/11, a
 year of strong La Niña manifestation. The positive sea surface temperature (SST) anomaly in the west Pacific is accompanied
 470 by increased convection over the maritime continent, which results in enhanced upper tropospheric outflow and a strengthening
 of the adjacent Walker circulation cell (Fig. 13c). Still, the frequent occurrence of strong vertical wind shear near to the LRT
 is striking, considering the still comparatively weak easterlies responsible. The occurrence of strong vertical wind shear near
 to the tropopause could be influenced by enhanced gravity wave activity associated with deep convection (Podglajen et al.,
 2017), as convection is the major source for gravity wave generation in the tropics through several forcing mechanisms (Müller
 475 et al. (2018) and references therein). During the pronounced El Niño winter seasons 2009/10 and 2015/16 the shear occurrence
 frequency reduces down to values in the range of 30 – 40 %, which is significantly below the 10 year average value of 54.2 %.
 This is associated with the eastward shift of the less sharply defined rising branch of the Pacific Walker circulation cell (Sullivan
 et al., 2019), along with the less pronounced upper tropospheric easterlies (Fig. 13b).

We now briefly revert to Figure 5b and the secondary occurrence frequency maximum for $S^2 \geq S_t^2$ below the tropical LRT,
 480 which is strikingly separated from the one above. This secondary maximum emerges due to a comparable situation to the one
 described for the secondary maxima at the subtropical tropopause break (Fig. 6), only on smaller scales and zonally aligned.
 The following explanation is applicable to both the TEJ as well as the the winter Walker cell easterlies. We will focus on
 the latter to give an exemplary explanation. The easterlies responsible for the DJF TSL occurrence frequency maximum are
 associated with a frequently occurring lapse rate tropopause jump of 0.5 – 1.0 km vertical distance, located at the western
 485 edge of the easterlies (Fig. 13c). This is indicated by the occurrence frequency isolines for $N^2 \geq 4 \cdot 10^{-4} \text{ s}^{-2}$. These regions
 of enhanced static stability reach west- and downward, where at some point they do no longer fulfill the definition of the
 LRT criterion, causing the tropopause jump. The tropopause-based averaging method then results in a secondary occurrence
 frequency maximum for $S^2 \geq S_t^2$ below the LRT, due to enhanced wind shear within the regions of enhanced N^2 that reach
 below the local LRT.

490 5 Discussion

The TSL and particularly its relation to the TIL owe their potential importance to the fact, that both the TSL and the TIL
 majorly influence the dynamical stability and may be related to the formation of the extratropical transition layer (ExTL),
 which is derived from observations of chemical tracers (e.g. Hoor et al., 2002; Pan et al., 2004). Related to that, the potential
 implications of the frequently occurring TSL within the tropical tropopause layer (TTL, Fueglistaler et al., 2009) needs further
 495 investigation.




Figure 14. a) Relative occurrence frequency distribution of N^2 - S^2 pairs in the region between the LRT and 3 km above, for all daily northern hemispheric ERA5 fields from 2008-2017. Logarithmic occurrence frequency color scale. Red dashed line indicates $S^2 = S_t^2$. Dashed black lines indicate the Richardson numbers 0.25, 0.5, 1.0, 2.0 and 4.0. b) Histogram of the relative distribution of Richardson numbers associated with the data displayed in panel a. Orange bars show Ri for grid volumes with $S^2 \geq S_t^2$, and blue bars for the remaining grid volumes between the LRT and 3 km above. Dotted black line indicates $Ri = 1$.

It is important to analyse the relation of the TSL and the TIL in more detail, because both co-occur as climatological features in zonal and/or temporal tropopause-based averages (e.g. Birner et al., 2002; Birner, 2006; Grise et al., 2010; Zhang et al., 2019). However, we see no general co-location of enhanced N^2 and enhanced S^2 above the local lapse rate tropopause in instantaneous considerations and on smaller scales. Figure 14a shows the relative occurrence frequency of N^2 - S^2 pairs for all ten years and in the region between the LRT and 3 km above. The majority of grid volumes exhibit a static stability between the stratospheric background $\overline{N^2}_{strat.} = 4 \cdot 10^{-4} \text{ s}^{-2}$ and moderately enhanced values associated with the TIL. At the same time, comparatively “weak” vertical wind shear $S^2 < 4 \cdot 10^{-4} \text{ s}^{-2}$ is most prevalent. Vertical wind shear and static stability do not correlate, and enhanced values of S^2 can be found within a large spectrum of N^2 under the condition that dynamic stability $Ri > Ri_c$ is (for the most part) maintained. Particularly the largest values of N^2 and S^2 do not correlate.

Ultimately this has consequences for the dynamic stability of the flow, along with the potential for turbulence and mixing across and above the tropopause. In the case of baroclinic waves in the north Atlantic storm track region, Kaluza et al. (2019) describe a co-occurrence of the TIL and tropopause-based enhanced vertical wind shear in cyclone-based composites. This indicates that the processes responsible are active in the same region on a synoptic scale, i.e. ridges. These processes however have a different impact on the temperature and momentum profiles, or even an exclusive direct effect on either. The vertical divergence of the vertical wind in the anticyclonic flow sharpens the temperature profile (Wirth, 2003, 2004), as well as long- and shortwave radiative effects due to trace gas and cloud ice water gradients at the tropopause (Randel et al., 2007; Kunkel

et al., 2016), where the latter is particularly important in the context of warm conveyor belts and high reaching cloud formation in the tropospheric flow within ridges (Madonna et al., 2014).

The connection between the temperature field and the vertical wind shear for synoptic scale flow can be approximated through the thermal wind relation, i.e., the vertical gradient of the geostrophic wind under the assumption of hydrostatic balance. Figure 15a shows the geostrophic wind at 200 hPa on 11 September 2017, i.e., the date of the exemplary single day analysis in Sect. 3. Overall, the geostrophic wind approximates the synoptic scale flow realistically (compare Fig. 2). However, it overestimates the absolute zonal wind speed in cyclonic rotational systems like the one over the northwest Atlantic, in accordance with the fact that inertial forces are neglected (Holton and Hakim, 2012). The vertical structure of the geostrophic wind is shown exemplarily in the vertical cross sections at 60° W (Fig. 15b and c). The thermal wind relation results in several regions of strong vertical wind shear near to the tropopause. The comparison with the vertical wind shear derived from the full model wind reveals a certain degree of agreement, in particular on the synoptic scale, but also differences on the smaller scales. The strength of the vertical wind shear at 40° N and 15 km altitude is overestimated by the thermal wind approximation, as well as the shear region directly above that reaches north- and downward. The southward extent of the region of strong wind shear on the other hand is underestimated. The two pronounced wind shear regions below the tropopause and south of 40° N are not evident in the thermal wind shear. The geostrophic zonal wind in the upper troposphere at about 50° N deviates from the full model zonal wind, which results in a significant underestimation the vertical wind shear below the tropopause. At the same time, the thermal wind relation overestimates the shear region below the tropopause at 45° N, which is caused by strong meridional temperature gradients (not shown). The maximum of the thermal wind shear at 45° N directly above the LRT is not evident in the full model wind shear, but instead is apparent in a region that is located further to the north. Overall, the comparison indicates the significance of dynamic processes on smaller scales on which other forces than pressure gradient and Coriolis force need to be taken into account (Newton and Persson, 1962). This example already shows that many details, especially related to mesoscale dynamic features, need to be considered to fully address the differences in the vertical wind shear based on the full model winds and on the thermal wind relation. A comprehensive analysis of these differences is beyond the scope of the current study but will be pursued in future work.

On sub-synoptic scales, gravity waves can have a large influence on both N^2 and S^2 above the tropopause (e.g. Kunkel et al., 2014), however according to linear wave theory with a phase shift of $\pi/2$ between the perturbation of the horizontal wind u' in direction of the horizontal wave vector k and the perturbation of the temperature T' (Suzuki et al., 2010), and a resulting shift between N^2 enhancement and S^2 enhancement (compare e.g. Kunkel et al., 2014). Recently, Podglajen et al. (2020) compared long-duration superpressure balloon measurements with Lagrangian trajectories calculated from a set of numerical reanalysis products, and were able to show that the ERA5 reanalysis resolves central features of the gravity wave spectrum. The underlying IFS model resolves low frequency large scale gravity waves down to wavelengths which approach the effective resolution, which is generally estimated to exhibit a factor of about 10 compared to the effective grid spacing. The assimilation of high resolution observational data further enhances the gravity wave activity in the model, which likely involves generation processes and gravity wave scales that are not resolved in the IFS. Furthermore, Krisch et al. (2020) identified individual wave packets in the ERA5 data that had been observed with the Gimbal Limb Observer for Radiance Imaging of the Atmosphere




Figure 15. Comparison of the vertical wind shear based on the full model winds and on the thermal wind relation. a) Geostrophic wind at 200 hPa, for the northern hemisphere on 11 September 2017. Regions south of 20° N are left out because the validity of the assumption of geostrophic balance vanishes towards the equator. Black solid line indicates the vertical cross sections in panel b and c. b) Color contour shows vertical shear of the horizontal wind derived from the wind components in the ERA5 data (S^2 , in ms^{-2}). Red and blue lines show $u = 30 \text{ ms}^{-1}$ and $u = -10 \text{ ms}^{-1}$ isotachs of the zonal wind. Black dotted lines show isentropes. Black circle markers indicate LRT altitude. c) As in b but for the thermal wind calculated from the temperature field in the ERA5 data. Red and blue lines show $u = 30 \text{ ms}^{-1}$ and $u = -10 \text{ ms}^{-1}$ isotachs of the zonal geostrophic wind.

(GLORIA) during the GW-LCYCLE airborne measurement campaign. While these studies confirm that modern reanalysis products are capable to resolve central features of the gravity wave spectrum, the overall vertical wind variability due to gravity waves is likely still underestimated in the ERA5. Recently, Schäfler et al. (2020) reported a significant underestimation of the vertical wind shear near to the tropopause in the IFS, based on a comparison of Doppler wind lidar measurements with IFS analysis and forecast data. The analysis and forecast errors were most prominent at elevated tropopause altitudes above upper tropospheric ridges, i.e., regions that contribute significantly to the occurrence of the TSL in the extratropics. To put these results into context, the model version used by Schäfler et al. (2020) exhibits a spectral truncation of TCo1280 on 137 vertical level, thus, the same vertical grid spacing compared to the ERA5 as well as a larger horizontal resolution.

Another issue that needs further investigation is the relation between the TSL and the occurrence of turbulence and potential STE. The grid volumes with $S^2 < 4 \cdot 10^{-4} \text{ s}^{-2}$ in Fig. 14a are mostly located below the diagonal dashed black line which

indicates Richardson numbers of $Ri = 1$. The corresponding relative distribution of Richardson numbers is shown in Fig. 14b. The distribution peaks at $Ri = 3$ and spans over a large spectrum of larger Richardson numbers (only a section of the distribution is displayed). Richardson numbers of $Ri < 1$ are rarely associated with vertical wind shear $S^2 < 4 \cdot 10^{-4} \text{ s}^{-2}$. This indicates that the lowermost stratosphere up to 3 km above the LRT is dynamically stable in the absence of strong vertical wind shear. In contrast, a significant proportion of grid volumes which exceed S_t^2 in Fig. 14a are located above the $Ri = 1$ isoline. These grid volumes constitute the greater part of Richardson numbers $Ri < 1$ within the first 3 km above the LRT, which is indicated in Fig. 14b.

We define the TSL as an occurrence frequency maximum for strong vertical wind shear in tropopause-relative zonal and temporal averages. This occurrence frequency maximum is composed of individual localised shear patches that emerge above the LRT, as the exemplary cross sections in Fig. 3 show, resulting in reduced Richardson numbers of the order of 1 in the elsewhere dynamically stable lower stratosphere. Strong wind shear does not correlate well with the occurrence of turbulence (e.g. Knox, 1997) because it does not necessarily imply dynamic instability, but it is a prerequisite for dynamic instability to occur. Thus, the occurrence frequency distribution presented in Fig. 5a narrows down the region where turbulence due to dynamic shear instability can be expected above the tropopause. The vertical confinement within a few kilometres above the lapse rate tropopause indicates a link to the chemically defined ExTL or mixing layer. Airborne in situ measured carbon monoxide (CO) profiles in the tropopause region exhibit a distinct "kink" in the vertical transition from tropospheric to stratospheric values, which defines the upper edge of a chemical transition layer. Hoor et al. (2004) determined it to be located at about 25 K potential temperature above the $Q = 2$ pvu dynamic tropopause, or 2–3 km above the lapse rate tropopause at high latitudes according to Pan et al. (2004). Hegglin et al. (2009) expanded these results onto a global scale based on CO-O₃ and H₂O-O₃ tracer-tracer correlations derived from Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) satellite data. Berthet et al. (2007) used a statistical approach on Lagrangian trajectories that were subjected to TST, with their trajectory model being driven by operational ECMWF analysis data. The relative contribution of TST trajectories showed a largely tropopause-following behaviour with a limited vertical extent, indicating the amount and vertical reach of the permeability of the transport barrier that is the tropopause. Hoor et al. (2010) continued the Lagrangian approach and linked the chemical transition layer to troposphere-stratosphere transition transport timescales in the range of 0–50 days, which is short compared to the transport timescales in the LMS above the ExTL. The vertical confinement of the TSL may contribute to the separation of the LMS with a lower layer being frequently affected by shear driven turbulent STE (e.g. Spreitzer et al., 2019; Kunkel et al., 2019). The TSL properties indicate its role as a contributing mechanism for the formation and maintenance of the ExTL, along with other known processes like e.g. radiation induced PV modifications (Zierl and Wirth, 1997) or convective injection of tropospheric air (Homeyer et al., 2014).

6 Summary

The goal of this study was to investigate the occurrence of strong vertical wind shear in the UTLS region. It was motivated by the wide range of research studies that describe the occurrence of strong vertical wind shear at the tropopause, i.e. in the context

of case studies on turbulent stratosphere-troposphere exchange (e.g. Shapiro, 1976, 1978; Whiteway et al., 2004; Kunkel et al., 2019), in climatological studies based on radiosonde data and focussing on the TIL (Birner et al., 2002; Birner, 2006) and its interaction with gravity waves (Zhang et al., 2015, 2019), or in the context of planetary circulation features like the tropical easterly jet (Sunilkumar et al., 2015), and in numerical model data studies (Kunkel et al., 2014; Liu, 2017; Kaluza et al., 2019; 595 Spreitzer et al., 2019). However, there was a lack of a comprehensive statistical analysis of this atmospheric feature in a global and climatological sense. We approached this matter using ten years of data of daily northern hemispheric ECMWF ERA5 reanalysis fields.

Strong vertical wind shear was selected based on the threshold value $S_t^2 = 4 \cdot 10^{-4} \text{ s}^{-2}$. A single day analysis introduced the metrics, and exemplified that strong vertical wind shear exceeding the threshold occurs nearly exclusively in close vertical 600 vicinity above the lapse rate tropopause. It furthermore showed that the strong vertical wind shear near to the tropopause can generally be attributed to planetary circulation features, i.e. polar jet streaks within differently evolved baroclinic waves, the subtropical jet stream, and the upper tropospheric easterlies of the Walker circulation cells.

The ten year temporally and zonally averaged meridional cross section revealed a tropopause wind shear layer (TSL), which is defined by an occurrence frequency maximum of the order of 1 %–10 % for strong vertical wind shear near to the tropopause. 605 These occurrence frequencies are about an order of magnitude larger than anywhere else in the region analysed. The TSL is vertically confined to the first 1–2 kilometres above the lapse rate tropopause, and spans from the tropics down to high latitudes around 70° N. The geographical mapping of the TSL revealed distinctly separate regions of preferred occurrence, that can each be linked to individual planetary circulation features:

- The storm track regions over the north Atlantic and the northeast Pacific feature distinct occurrence frequency maxima 610 for enhanced tropopause-based vertical wind shear. These maxima are most pronounced during DJF, and are associated with jet streaks of the eddy-driven jet. The enhanced tropopause-based vertical wind shear emerges at above-average tropopause altitudes associated with ridges of baroclinic waves. We identified the northward excursion of air masses within ridges on the basis of deviations of the potential temperature at the dynamic tropopause from the ten year temporal and zonal average within each region. The occurrence frequencies for strong vertical wind shear near to the tropopause 615 maximise at latitudes around 50° N, and reach values up to 50 % for profiles associated with above-average dynamic tropopause potential temperatures of $\Delta\Theta \approx 20 \text{ K}$.
- The east Asian jet stream during winter exhibits occurrence frequencies of up to 45 % for strong vertical wind shear near to the tropopause, with a pronounced interannual variability. This appears to be linked to the variability in strength and location of the jet stream (Wu and Sun, 2017), which needs further investigation.
- 620 – The lapse rate tropopause region above the summer tropical easterly jet is up to 70 % of the time exposed to strong vertical wind shear near to the tropopause and over a large area of the Indian Ocean. These very striking occurrence frequency maxima as well as their interannual variability agree well with results from observation-based research studies (Roja Raman et al., 2009; Sunilkumar et al., 2015).

625 – The winter Walker circulation easterlies over the maritime continent exhibit an occurrence frequency maximum for strong vertical wind shear near to the tropopause which is comparable to the one associated with the tropical easterly jet. The strength and zonal location of these easterlies is closely linked to the el Niño sea surface temperature anomaly and the shift in the Walker circulation. The area-averaged occurrence frequencies for strong vertical wind shear near to the tropopause above the maritime continent show a pronounced interseasonal variability, ranging from 30 – 40 % during the el Niño phases in the winter seasons 2009/10 and 2015/2016, and up to 70 % during the strong la Niña phase 2010/11.

630 Overall, this analysis presents a step towards a better understanding of the dynamical structure of the transition region between the troposphere and the stratosphere, i.e. the ExTL in the extratropics and the TTL in the tropics. The occurrence of the TSL can have several implications:

635 – The TSL could be an indicator for the efficiency of the processes responsible for its formation. The occurrence frequency for strong vertical wind shear near to the tropopause could for example correlate with the gravity wave activity, depending on the relevance of gravity waves for the formation of the TSL, i.e. their bi-directional interaction with the thermal stratification gradients that define the LRT and the TIL. The occurrence of the TSL in regions that are frequently exposed to enhanced gravity wave activity hints towards such a link, i.e. jet-front systems in baroclinic waves in the extratropics, and the rising branch of the Pacific Walker circulation.

640 – In this context, the occurrence of the TSL could also be an indicator for nonlinear wave-mean-flow-interaction and momentum deposition (Zhang et al., 2019; Bense, 2019). Strong vertical wind shear could be a residuum of preceding turbulence in vertically adjacent layers, as turbulent homogenisation can result in adjacent gradient sharpening.

645 – The TSL should have a noticeable intersection with the large scale forcing for tropopause regions that are susceptible to turbulent STE. This however needs thorough further investigation, as e.g. the differences between this analysis (Fig. 7) and the tropopause-based turbulence indicators by Jaeger and Sprenger (2007) show. The limitations of the numerical models used are an important factor to consider.

Overall, the work presented puts previous research studies on strong vertical wind shear at tropopause altitudes into the context of the planetary circulation, and expands the results to present strong vertical wind shear near to the tropopause as a global-scale UTLS feature. Conceptually, the top end values of the spectrum of atmospheric vertical wind shear in the UTLS are to be expected to occur in the tropopause region, because it fulfills the requirement of exceptional thermal wind gradient forcing as well as a stratification that can sustain the wind shear. The strict vertical confinement of the occurrence of $S^2 \geq 4 \cdot 10^{-4} \text{ s}^{-2}$ within the first two kilometres above the LRT however is quite striking, and it remains to be evaluated to what amount the wind-forcing and the required thermal stratification are mutually dependent. The role of additional processes like gravity wave propagation, refraction, and momentum deposition at the tropopause should be investigated in this context.

655 *Code and data availability.* ECMWF ERA5 reanalysis data has been retrieved from the MARS server. Output from the data analysis steps and further information on the analysis code are available upon request (kaluzat@uni-mainz.de).

Author contributions. DK, PH, and TK designed the research project. TK developed the model code, performed the calculations, and analysed the data with the help of DK and PH. TK prepared the paper with contributions from all authors.

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. TEXT

660 *Acknowledgements.* The Authors acknowledge funding from the German Science Foundation, as this study was carried out in the context of the WISE campaign under funding from the HALO SPP 1294 (DFG grant no. KU 3524/1-1, HO 4225/7-1 as well as HO 4225/8-1). We are furthermore grateful to the ECMWF for providing the ERA5 data which has been downloaded from the ECMWF MARS system.

References

- Bense, V.: Modifikation von Schwerewellen bei Propagation durch die Tropopause – Idealisierte Modellstudien, Dissertation, Johannes Gutenberg-University Mainz, <https://doi.org/http://doi.org/10.25358/openscience-2356>, 2019.
- Berthet, G., Esler, J. G., and Haynes, P. H.: A Lagrangian perspective of the tropopause and the ventilation of the lowermost stratosphere, *Journal of Geophysical Research Atmospheres*, 112, 1–14, <https://doi.org/10.1029/2006JD008295>, 2007.
- Birner, T.: Fine-scale structure of the extratropical tropopause region, *Journal of Geophysical Research Atmospheres*, 111, 1–14, <https://doi.org/10.1029/2005JD006301>, 2006.
- Birner, T., Dörnbrack, A., and Schumann, U.: How sharp is the tropopause at midlatitudes?, *Geophysical Research Letters*, 29, 45–1–45–4, <https://doi.org/10.1029/2002GL015142>, 2002.
- Dvoskin, N. and Sissenwine, N.: Evaluation of AN/GMD-2 Wind Shear Data for Development of Missile Design Criteria, *Air Force Surveys in Geophysics*, 99, 72 pp., 1958.
- ECMWF, a.: IFS documentation – Cy41r2 Operational implementation 8 March 2016 Part III : Dynamics and numerical procedures, 2016.
- Ellrod, G. P. and Knapp, D. I.: An Objective Clear-Air Turbulence Forecasting Technique: Verification and Operational Use, *Weather and Forecasting*, 7, 15, [https://doi.org/https://doi.org/10.1175/1520-0434\(1992\)007<0150:AOCATF>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0434(1992)007<0150:AOCATF>2.0.CO;2), 1992.
- Endlich, R. M. and McLean, G. S.: Jet-Stream Structure over the Central United States Determined from Aircraft Observations, *Journal of Applied Meteorology*, 4, 83–90, [https://doi.org/10.1175/1520-0450\(1965\)004<0083:jssotc>2.0.co;2](https://doi.org/10.1175/1520-0450(1965)004<0083:jssotc>2.0.co;2), 1965.
- Ertel, H.: Ein neuer hydrodynamischer Erhaltungssatz, *Die Naturwissenschaften*, 30, 543–544, <https://doi.org/10.1007/BF01475602>, 1942.
- Forster, P. M. D. E. and Shine, K. P.: Radiative forcing and temperature trends from stratospheric ozone changes and globally averaged radiative was found from SBUV total column ozone coolings of up to 0 . 4 K decade ⁻¹ can be found at altitudes of 35 km ; this provides a, *Reading*, 102, 1997.
- Fritts, D. C., Wang, L., and Werne, J. A.: Gravity wave-fine structure interactions. Part I: Influences of fine structure form and orientation on flow evolution and instability, *Journal of the Atmospheric Sciences*, 70, 3710–3734, <https://doi.org/10.1175/JAS-D-13-055.1>, 2013.
- Fritts, D. C., Wang, L., Geller, M. A., Lawrence, D. A., Werne, J., and Balsley, B. B.: Numerical modeling of multiscale dynamics at a high reynolds number: Instabilities, turbulence, and an assessment of ozmidov and thorpe scales, *Journal of the Atmospheric Sciences*, 73, 555–578, <https://doi.org/10.1175/JAS-D-14-0343.1>, 2016.
- Fritts, D. C., Wang, L., Baumgarten, G., Miller, A. D., Geller, M. A., Jones, G., Limon, M., Chapman, D., Didier, J., Kjellstrand, C. B., Araujo, D., Hillbrand, S., Korotkov, A., Tucker, G., and Vinokurov, J.: High-resolution observations and modeling of turbulence sources, structures, and intensities in the upper mesosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, 162, 57–78, <https://doi.org/10.1016/j.jastp.2016.11.006>, 2017.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical Tropopause Layer, *Geophys.*, 47, RG1004, DOI:, pp. 1–31, <https://doi.org/10.1029/2008RG000267.1>.INTRODUCTION, 2009.
- Gettelman, A. and Wang, T.: Structural diagnostics of the tropopause inversion layer and its evolution, *Journal of Geophysical Research Atmospheres*, 120, 46–62, <https://doi.org/10.1002/2014JD021846>, 2015.
- Grise, K. M., Thompson, D. W., and Birner, T.: A global survey of static stability in the stratosphere and upper troposphere, *Journal of Climate*, 23, 2275–2292, <https://doi.org/10.1175/2009JCLI3369.1>, 2010.

Hegglin, M. I., Boone, C. D., Manney, G. L., and Walker, K. A.: A global view of the extratropical tropopause transition layer from Atmospheric Chemistry Experiment Fourier Transform Spectrometer O₃, H₂O, and CO, *Journal of Geophysical Research Atmospheres*, 114, 1–18, <https://doi.org/10.1029/2008JD009984>, 2009.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, pp. 1–51, <https://doi.org/10.1002/qj.3803>, 2020.

Hoffmann, L., Günther, G., Li, D., Stein, O., Wu, X., Griessbach, S., Heng, Y., Konopka, P., Müller, R., Vogel, B., and Wright, J. S.: From ERA-Interim to ERA5: The considerable impact of ECMWF’s next-generation reanalysis on Lagrangian transport simulations, *Atmospheric Chemistry and Physics*, 19, 3097–3214, <https://doi.org/10.5194/acp-19-3097-2019>, 2019.

Holton, J. R.: Stratosphere–Troposphere Exchange, *Stratosphere Troposphere Interactions*, pp. 331–355, https://doi.org/10.1007/978-1-4020-8217-7_8, 1995.

Holton, J. R. and Hakim, G. J.: An introduction to dynamic meteorology: Fifth edition, vol. 9780123848, <https://doi.org/10.1016/C2009-0-63394-8>, 2012.

Homeyer, C. R., Pan, L. L., Dorsi, S. W., Avallone, L. M., Weinheimer, A. J., O’Brien, A. S., Digangi, J. P., Zondlo, M. A., Ryerson, T. B., Diskin, G. S., and Campos, T. L.: Convective transport of water vapor into the lower stratosphere observed during double-tropopause events, *Journal of Geophysical Research*, 119, 10,941–10,958, <https://doi.org/10.1002/2014JD021485>, 2014.

Hoor, P., Fischer, H., Lange, L., Lelieveld, J., and Brunner, D.: Seasonal variations of a mixing layer in the lowermost stratosphere as identified by the CO–O₃ correlation from in situ measurements, *Journal of Geophysical Research: Atmospheres*, 107, ACL 1–1–ACL 1–11, <https://doi.org/10.1029/2000JD000289>, 2002.

Hoor, P., Gurk, C., Brunner, D., Hegglin, M. I., Wernli, H., and Fischer, H.: Seasonality and extent of extratropical TST derived from in-situ CO measurements during SPURT, *Atmospheric Chemistry and Physics Discussions*, 4, 1691–1726, <https://doi.org/10.5194/acpd-4-1691-2004>, 2004.

Hoor, P., Wernli, H., Hegglin, M. I., and Bönisch, H.: Transport timescales and tracer properties in the extratropical UTLS, *Atmospheric Chemistry and Physics*, 10, 7929–7944, <https://doi.org/10.5194/acp-10-7929-2010>, 2010.

Hoskins, B. J.: Towards a PV- θ view of the general circulation, *Tellus A*, 43, 27–35, <https://doi.org/10.1034/j.1600-0870.1991.t01-3-00005.x>, 1991.

Hoskins, B. J., McIntyre, M. E., and Robertson, A. W.: On the use and significance of isentropic potential vorticity maps, *The Quarterly Journal of the Royal Meteorological Society*, 111, 877–946, <https://doi.org/10.1002/qj.49711146602>, 1985.

Jaeger, E. B. and Sprenger, M.: A Northern Hemispheric climatology of indices for clear air turbulence in the tropopause region derived from ERA40 reanalysis data, *Journal of Geophysical Research Atmospheres*, 112, 1–13, <https://doi.org/10.1029/2006JD008189>, 2007.

Kaluza, T., Kunkel, D., and Hoor, P.: Composite analysis of the tropopause inversion layer in extratropical baroclinic waves, *Atmospheric Chemistry and Physics*, 19, 1–22, <https://doi.org/10.5194/acp-2018-1100>, 2019.

Knox, J. A.: Possible mechanisms of clear-air turbulence in strongly anticyclonic flows, *Monthly Weather Review*, 125, 1251–1259, [https://doi.org/10.1175/1520-0493\(1997\)125<1251:PMOCAT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<1251:PMOCAT>2.0.CO;2), 1997.

Koch, P., Wernli, H., and Davies, H. C.: An event-based jet-stream climatology and typology, *International Journal of Climatology*, 26, 283–301, <https://doi.org/10.1002/joc.1255>, 2006.

- Koteswaram, P.: The Easterly Jet Stream in the Tropics, *Tellus*, 10, 43–57, <https://doi.org/10.1111/j.2153-3490.1958.tb01984.x>, 1958.
- Krisch, I., Ern, M., Hoffmann, L., Preusse, P., Strube, C., Ungermann, J., Woiwode, W., and Riese, M.: Superposition of gravity waves with different propagation characteristics observed by airborne and space-borne infrared sounders, *Atmospheric Chemistry and Physics*, 20, 11 469–11 490, <https://doi.org/10.5194/acp-20-11469-2020>, 2020.
- Krishnamurti, T. N. and Bhalme, H.: Ocillations of a Monsoon System. Part I. Observational Aspects, *Journal of Atmospheric Sciences*, 33, 1937–1954, [https://doi.org/10.1175/1520-0469\(1976\)033<1937:OOAMSP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<1937:OOAMSP>2.0.CO;2), 1976.
- Kunkel, D., Hoor, P., and Wirth, V.: Can inertia-gravity waves persistently alter the tropopause inversion layer?, *Geophysical Research Letters*, 41, 7822–7829, <https://doi.org/10.1002/2014GL061970>, 2014.
- 745 Kunkel, D., Hoor, P., and Wirth, V.: The tropopause inversion layer in baroclinic life-cycle experiments: The role of diabatic processes, *Atmospheric Chemistry and Physics*, 16, 541–560, <https://doi.org/10.5194/acp-16-541-2016>, 2016.
- Kunkel, D., Hoor, P., Kaluza, T., Ungermann, J., Kluschat, B., Giez, A., Lachnitt, H. C., Kaufmann, M., and Riese, M.: Evidence of small-scale quasi-isentropic mixing in ridges of extratropical baroclinic waves, *Atmospheric Chemistry and Physics*, 19, 12 607–12 630, <https://doi.org/10.5194/acp-19-12607-2019>, 2019.
- 750 Liu, H. L.: Large Wind Shears and Their Implications for Diffusion in Regions With Enhanced Static Stability: The Mesopause and the Tropopause, *Journal of Geophysical Research: Atmospheres*, 122, 9579–9590, <https://doi.org/10.1002/2017JD026748>, 2017.
- Madonna, E., Wernli, H., Joos, H., and Martius, O.: Warm conveyor belts in the ERA-Interim Dataset (1979–2010). Part I: Climatology and potential vorticity evolution, *Journal of Climate*, 27, 3–26, <https://doi.org/10.1175/JCLI-D-12-00720.1>, 2014.
- Miles, J. W.: On the stability of heterogeneous shear flows, *Journal of Fluid Mechanics*, 10, 496–508, <https://doi.org/10.1017/S0022112061000305>, 1961.
- 755 Müller, S. K., Manzini, E., Giorgetta, M., Sato, K., and Nasuno, T.: Convectively Generated Gravity Waves in High Resolution Models of Tropical Dynamics, *Journal of Advances in Modeling Earth Systems*, 10, 2564–2588, <https://doi.org/10.1029/2018MS001390>, 2018.
- Newton, C. W. and Persson, A. V.: Structural characteristics of the subtropical jet stream and certain lower-stratospheric wind systems, *Tellus*, 14, 2, <https://doi.org/10.3402/tellusa.v14i2.9542>, 1962.
- 760 Pan, L. L., Randel, W. J., Gary, B. L., Mahoney, M. J., and Hints, E. J.: Definitions and sharpness of the extratropical tropopause: A trace gas perspective, *Journal of Geophysical Research D: Atmospheres*, 109, 1–11, <https://doi.org/10.1029/2004JD004982>, 2004.
- Plougonven, R. and Zhang, F.: Internal gravity waves from atmospheric jets and fronts, *American Geophysical Union*, pp. 1022–1049, <https://doi.org/10.1002/2015RG000489>.Received, 2014.
- Podglajen, A., Bui, T. P., Dean-Day, J. M., Pfister, L., Jensen, E. J., Alexander, M. J., Hertzog, A., Kärcher, B., Plougonven, R., and Randel, W. J.: Small-scale wind fluctuations in the tropical tropopause layer from aircraft measurements: Occurrence, nature, and impact on vertical mixing, *Journal of the Atmospheric Sciences*, 74, 3847–3869, <https://doi.org/10.1175/JAS-D-17-0010.1>, 2017.
- 765 Podglajen, A., Hertzog, A., Plougonven, R., and Legras, B.: Lagrangian gravity wave spectra in the lower stratosphere of current (re)analyses, *Atmospheric Chemistry and Physics*, 20, 9331–9350, <https://doi.org/10.5194/acp-20-9331-2020>, 2020.
- Randel, W. J., Wu, F., and Forster, P.: The Extratropical Tropopause Inversion Layer: Global Observations with GPS Data, and a Radiative Forcing Mechanism, *Journal of the Atmospheric Sciences*, 64, 4489–4496, <https://doi.org/10.1175/2007JAS2412.1>, 2007.
- 770 Reiter, E. R.: Clear Air Turbulence and its Detection, The Nature of Clear Air Turbulence: A Review, in: *Proceedings of a Symposium on Clear Air Turbulence and Its Detection*, pp. 19–34, Flight Sciences Laboratory, Boeing Scientific Research Laboratories, The Boeing Company, Plenum Press, New York, 1969.

- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, *Journal of Geophysical Research Atmospheres*, 117, 1–10, <https://doi.org/10.1029/2012JD017751>, 2012.
- Roja Raman, M., Jagannadha Rao, V. V., Venkat Ratnam, M., Rajeevan, M., Rao, S. V., Narayana Rao, D., and Prabhakara Rao, N.: Characteristics of the Tropical Easterly Jet: Long-term trends and their features during active and break monsoon phases, *Journal of Geophysical Research Atmospheres*, 114, 1–14, <https://doi.org/10.1029/2009JD012065>, 2009.
- Schäfler, A., Harvey, B., Methven, J., Doyle, J. D., Rahm, S., Reitebuch, O., Weiler, F., and Witschas, B.: Observation of jet stream winds during nawdex and characterization of systematic meteorological analysis errors, *Monthly Weather Review*, 148, 2889–2907, <https://doi.org/10.1175/MWR-D-19-0229.1>, 2020.
- Seidel, D. J., Ross, R. J., Angell, J. K., and Reid, G. C.: Climatological characteristics of the tropical tropopause as revealed by radiosondes, *Journal of Geophysical Research Atmospheres*, 106, 7857–7878, <https://doi.org/10.1029/2000JD900837>, 2001.
- Shapiro, M. A.: The role of turbulent heat flux in the generation of potential vorticity in the vicinity of upper-level jet stream systems, *Monthly Weather Review*, 104, 1976.
- Shapiro, M. A.: Further evidence of the mesoscale and turbulent structure of upper level jet stream–frontal zone systems, *Monthly Weather Review*, 106, 1978.
- Sharman, R. D., Trier, S. B., Lane, T. P., and Doyle, J. D.: Sources and dynamics of turbulence in the upper troposphere and lower stratosphere : A review, *Geophysical Research Letters*, 39, 1–9, <https://doi.org/10.1029/2012GL051996>, 2012.
- Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y. T., Li, C., O’Gorman, P. A., Rivière, G., Simpson, I. R., and Voigt, A.: Storm track processes and the opposing influences of climate change, *Nature Geoscience*, 9, 656–664, <https://doi.org/10.1038/ngeo2783>, 2016.
- Škerlak, B., Sprenger, M., Pfahl, S., Tyrlis, E., and Wernli, H.: Tropopause folds in ERA-interim: Global climatology and relation to extreme weather events, *Journal of Geophysical Research*, 120, 4860–4877, <https://doi.org/10.1002/2014JD022787>, 2015.
- Spreitzer, E., Attinger, R., Boettcher, M., Forbes, R., Wernli, H., and Joos, H.: Modification of potential vorticity near the tropopause by nonconservative processes in the ECMWF model, *Journal of the Atmospheric Sciences*, 76, 1709–1726, <https://doi.org/10.1175/JAS-D-18-0295.1>, 2019.
- Stohl, A., Bonasoni, P., Cristofanelli, P., Collins, W., Feichter, J., Frank, A., Forster, C., Gerasopoulos, E., Gaggeler, H., James, P., Kentarchos, T., Kromp-Kolb, H., Krüger, B., Land, C., Meloen, J., Papayannis, A., Priller, A., Seibert, P., Sprenger, M., Roelofs, G. J., Scheel, H. E., Schnabel, C., Siegmund, P., Tobler, L., Trickl, T., Wernli, H., Wirth, V., Zanis, P., and Zerefos, C.: Stratosphere-troposphere exchange: A review, and what we have learned from STACCATO, *Journal of Geophysical Research Atmospheres*, 108, <https://doi.org/10.1029/2002jd002490>, 2003.
- Sullivan, S. C., Schiro, K. A., Stubenrauch, C., and Gentine, P.: The Response of Tropical Organized Convection to El Niño Warming, *Journal of Geophysical Research: Atmospheres*, 124, 8481–8500, <https://doi.org/10.1029/2019JD031026>, 2019.
- Sunilkumar, S. V., Babu, A., and Parameswaran, K.: Mean structure of the tropical tropopause and its variability over the Indian longitude sector, *Climate Dynamics*, 40, 1125–1140, <https://doi.org/10.1007/s00382-012-1496-8>, 2013.
- Sunilkumar, S. V., Muhsin, M., Parameswaran, K., Venkat Ratnam, M., Ramkumar, G., Rajeev, K., Krishna Murthy, B. V., Sambhu Namboodiri, K. V., Subrahmanyam, K. V., Kishore Kumar, K., and Shankar Das, S.: Characteristics of turbulence in the troposphere and lower stratosphere over the Indian Peninsula, *Journal of Atmospheric and Solar-Terrestrial Physics*, 133, 36–53, <https://doi.org/10.1016/j.jastp.2015.07.015>, 2015.

- Suzuki, S., Nakamura, T., Ejiri, M. K., Tsutsumi, M., Shiokawa, K., and Kawahara, T. D.: Simultaneous airglow, lidar, and radar measurements of mesospheric gravity waves over Japan, *Journal of Geophysical Research Atmospheres*, 115, 1–12, <https://doi.org/10.1029/2010JD014674>, 2010.
- 815 Trier, S. B., Sharman, R. D., Muñoz-Esparza, D., and Lane, T. P.: Environment and Mechanisms of Severe Turbulence in a Midlatitude Cyclone, *Journal of the Atmospheric Sciences*, 77, 3869–3889, <https://doi.org/10.1175/jas-d-20-0095.1>, 2020.
- Whiteway, J. A., Klaassen, G. P., Bradshaw, N. G., and Hacker, J.: Transition to turbulence in shear above the tropopause, *Geophysical Research Letters*, 31, 2–5, <https://doi.org/10.1029/2003GL018509>, 2004.
- Wirth, V.: Static Stability in the Extratropical Tropopause Region, *Journal of the Atmospheric Sciences*, 60, 1395–1409, [https://doi.org/10.1175/1520-0469\(2003\)060<1395:SSITET>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<1395:SSITET>2.0.CO;2), 2003.
- 820 Wirth, V.: A dynamical mechanism for tropopause sharpening, *Meteorologische Zeitschrift*, 13, 477–484, <https://doi.org/10.1127/0941-2948/2004/0013-0477>, 2004.
- WMO: Meteorology — A three-dimensional science: Second session of the Commission for Aerology, *WMO Bulletin*, VI, 134–138, 1957.
- Wu, S. and Sun, J.: Variability in zonal location of winter East Asian jet stream, *International Journal of Climatology*, 37, 3753–3766, <https://doi.org/10.1002/joc.4947>, 2017.
- 825 Yang, S., Lau, K. M., and Kim, K. M.: Variations of the East Asian jet stream and Asian-Pacific-American winter climate anomalies, *Journal of Climate*, 15, 306–325, [https://doi.org/10.1175/1520-0442\(2002\)015<0306:voteaj>2.0.co;2](https://doi.org/10.1175/1520-0442(2002)015<0306:voteaj>2.0.co;2), 2002.
- Zhang, Y., Zhang, S., Huang, C., Huang, K., Gong, Y., and Gan, Q.: The interaction between the tropopause inversion layer and the inertial gravity wave activities revealed by radiosonde observations at a midlatitude station Yehui, *Journal of Geophysical Research Atmospheres*, 120, 175, 238, <https://doi.org/10.1038/175238c0>, 2015.
- 830 Zhang, Y., Zhang, S., Huang, C., Huang, K., and Gong, Y.: The Tropopause Inversion Layer Interaction With the Inertial Gravity Wave Activities and Its Latitudinal Variability, *Journal of Geophysical Research: Atmospheres*, 124, 7512–7522, <https://doi.org/10.1029/2019JD030309>, 2019.
- Zierl, B. and Wirth, V.: The influence of radiation on tropopause behavior and stratosphere-troposphere exchange in an upper tropospheric anticyclone., *Journal of Geophysical Research Atmospheres*, 102, 23 883–23 894, <https://doi.org/https://doi.org/10.1029/97JD01667>, 1997.
- 835