

# ~~Mechanisms driving MJO teleconnection changes with warming~~

## Drivers of uncertainty in CMIP6 future projections of MJO teleconnections

Andrea M. Jenney<sup>1,2</sup>, David A. Randall<sup>2</sup>, and Elizabeth A. Barnes<sup>2</sup>

<sup>1</sup>Department of Earth System Science, University of California, Irvine, CA, USA

<sup>2</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA

**Correspondence:** Andrea M. Jenney (ajenney@uci.edu)

**Abstract.** Teleconnections from the Madden-Julian Oscillation (MJO) are a key source of predictability of weather on the extended time scale of about 10-40 days. The MJO teleconnection is sensitive to a number of factors, including the mean ~~state~~ dry static stability, the mean flow, and the propagation and intensity characteristics of the MJO ~~itself~~, which are traditionally difficult to separate across models. Each of these factors may evolve in response to increasing greenhouse gas emissions, which will impact MJO teleconnections and potentially impact potential predictability on extended time scales. Current state-of-the-art climate models do not agree on how MJO teleconnections over much of North America will change in a future climate. Here, we use results from the Coupled Model Intercomparison Project Phase 6 (CMIP6) historical and SSP585 experiments in concert with a linear baroclinic model (LBM) to separate and investigate alternate mechanisms explaining why and how boreal winter (January) MJO teleconnections over the North Pacific and North America ~~will~~ may change in a future climate, and to identify key sources of inter-model uncertainty. We find that ~~decreases to the MJO~~ a weakening teleconnection due to increases in tropical dry static stability alone are robust across CMIP6 models, and that uncertainty in mean state winds are a key driver of uncertainty in future MJO teleconnections. ~~We find no systematic relationship between changes in Rossby wave excitation and the MJO teleconnection. However, we find that models that predict increases (decreases) in the stationary Rossby wave number over the gulf of Alaska also predict stronger (weaker) teleconnections over North America.~~ Uncertainty in future changes to the MJO's intensity, eastward propagation speed, zonal wavenumber, and eastward propagation extent are other important sources of uncertainty in future MJO teleconnections, ~~although to a lesser degree than uncertainty in the mean state.~~ ~~The overall outlook is~~. We find no systematic relationship between future changes in the Rossby wave source and the MJO teleconnection, or between changes to the zonal wind or stationary Rossby wave number and the MJO teleconnection over the North Pacific and North America. LBM simulations suggest a reduction of the boreal winter MJO teleconnection ~~across the vast majority of CMIP6 models, especially~~ over the North Pacific, ~~but with some nuance~~ and an uncertain change over North America ~~due to larger sensitivity to expansion of the MJO's eastward extent,~~ with large spread over both regions that lends to weak confidence in the overall outlook.

Copyright statement. TEXT

## 1 Introduction

25 As the most energetic mode of tropical intraseasonal variability, the Madden-Julian Oscillation (MJO) is one of the most important sources of global weather predictability on the extended time range of about 10-40 days (Robertson et al., 2015). ~~Its impact on remote weather is sensitive to a number of factors including the static stability of the tropics, the magnitude of the MJO's convective and radiative heating anomalies, its propagation speed, the eastward extent of MJO convective propagation, and the mean state winds, each of which may evolve in response to increased atmospheric greenhouse gas concentrations. There is a~~  
30 ~~considerable amount of uncertainty in the response of many of these features to warming. Additionally, cloud parameterizations and coarse grid spacing lead to a relatively poor simulation of the MJO in many climate simulations (Ahn et al., 2017). It is thus~~ While previous work suggests that over specific regions (such as the North American west coast) models agree on how MJO teleconnections will change in a future climate, over much of the North Pacific and over North America, it is unclear  
35 how the influence of the MJO ~~over much of the extratropics will change in a future warmer climate~~ will evolve with increasing atmospheric greenhouse gases (Zhou et al., 2020).

Whereas some consensus has been emerging about at least one of these controls that affects global MJO teleconnection strength—the increase of the tropical dry static stability with surface warming—recent studies suggest this control may be secondary (Bui and Maloney, 2018; Maloney et al., 2019; Bui and Maloney, 2019a, b; Zhou et al., 2020). To first order, the rate that air rises and sinks in the MJO's circulation is tightly constrained by, and inversely proportional to, the tropical dry static  
40 stability (Wolding et al., 2016). Latent heat release associated with MJO precipitation is balanced by the upward advection of dry static energy. Similarly, radiative cooling in the MJO's dry region is balanced by slow, adiabatic subsidence. That is, the MJO's circulation strength is directly proportional to its diabatic heating rate, and inversely proportional to the dry static stability. It is expected that the tropical static stability will increase in the future as the tropical temperature profile adjusts towards the moist adiabatic lapse rate of a warmer surface (e.g., Santer et al., 2005). Ignoring changes to the MJO's precipi-  
45 tation intensity or its cloud optical properties ~~—(i.e., its diabatic heating rate),~~ this will weaken the MJO's vertical circulation and its associated upper-level divergence. In simulation studies of a future climate forced with increasing greenhouse gases, many models predict a weakening of the MJO's circulation strength (Bui and Maloney, 2018; Maloney et al., 2019). However, there is a considerable amount of inter-model spread, much of which is tied to disagreement in future projections of the MJO's diabatic heating and precipitation (Bui and Maloney, 2018; Maloney et al., 2019; Bui and Maloney, 2019a, b). While previous  
50 work has attributed a simulated weakening of MJO teleconnections in one model to the increase in dry static stability (Wolding et al., 2017), a recent model intercomparison study has shown that in climate change simulations the MJO's circulation change is not correlated with its teleconnection changes (Zhou et al., 2020).

~~We hypothesize~~ Samarasinghe et al. (2021) suggest that the lack of an apparent relationship between modeled MJO circulation change and teleconnection change may be related to changes in the mean state winds. Mean state winds refer to the winds  
55 that vary slowly in time (i.e., on seasonal, rather than daily time scales). Henderson et al. (2017) emphasize the importance of the mean state winds in the MJO teleconnection pattern, showing that biases in modeled MJO teleconnections can be attributed, in part, to biases in the mean state winds. Mean state winds affect both Rossby wave excitation and their propagation. The mag-

60 nitude of the vorticity and its horizontal gradient on the equatorward flank of the jet play an important role in the excitation of Rossby waves by the MJO (~~Sardeshmukh and Hoskins, 1988; Mori and Watanabe, 2008; Seo and Son, 2012; Seo and Lee, 2017~~)  
60 (~~Sardeshmukh and Hoskins, 1988; Mori and Watanabe, 2008; Seo and Son, 2012; Seo and Lee, 2017; Zheng and Chang, 2020~~)  
. The jets also act as a waveguide, and their structure and intensity determine the path that Rossby waves take as they propa-  
gate, and the sizes of Rossby waves that can propagate through them (e.g., Hoskins and Ambrizzi, 1993; Karoly, 1983). Tseng  
et al. (2020b) ~~show that differences in Rossby wave propagation, due to differences in~~ and ~~Zheng and Chang (2020) show how~~  
~~variability in the~~ mean state winds over the eastern ~~Pacific, drives most of the interannual~~ North Pacific drives variability in the  
65 MJO teleconnection~~pattern over the northern Pacific and North America region.~~, with variations in the jet's zonal extension  
or meridional position playing an important role in modulating wave propagation, and hence MJO teleconnectivity, over North  
America. Finally, Zhou et al. (2020) show that modeled increases in the MJO's impact over the North American west coast  
with warming are due to an extension of the Pacific jet, which causes a shift in the MJO teleconnection pattern. However, little  
work has been done to explore the direct role that changes to the mean state winds have on changing MJO teleconnections to  
70 the broader North Pacific and North America region.

In addition to features of the mean state (its static stability and winds), the MJO teleconnection may also respond to changes  
in the MJO itself. The MJO teleconnection is known to be sensitive to its eastward propagation speed. ~~While one study found~~  
~~little sensitivity of extratropical teleconnections to the eastward propagation speed of tropical heating (Goss and Feldstein, 2017)~~  
~~, many studies have found the opposite. Comparisons between composites of observed geopotential height anomalies during~~  
75 ~~fast and slow MJO events show differences in both the teleconnection pattern and magnitude of the anomalies (Yadav and Straus, 2017)~~  
~~. Models with faster propagating MJOs have weaker MJO teleconnections (Wang et al., 2020; Zheng and Chang, 2019).~~ In a  
simulation study, Bladé and Hartmann (1995) showed that an eastward propagating heat source is associated with a weaker  
and smaller wavetrain than a stationary or westward propagating heat source. To first order, this is a linear effect: eastward  
(westward) propagation effectively embeds the heating in easterlies (westerlies) (see Sect. 2b of Bladé and Hartmann, 1995).  
80 Many climate models predict an increase in the MJO's propagation speed with warming (e.g., Rushley et al., 2019), which,  
~~considering previous work,~~ may contribute to a weakening, ~~or at least a change in the pattern~~ of the MJO teleconnection with  
warming.

In the present climate, MJO convection is generally confined to the Indian and West Pacific oceans, while the MJO circulation  
signal circumnavigates the tropics. In simulations of a future warmer climate, many models predict an increase in the eastward  
85 extent of MJO convection (Adames et al., 2017; Bui and Maloney, 2018; Chang et al., 2015; Subramanian et al., 2014). This  
may be driven by the expansion of the eastward edge of the Indo-Pacific warm pool, a region of very warm tropical sea surface  
temperatures (Maloney and Xie, 2013; Zhou et al., 2020). A more eastward propagating MJO ~~will~~ may extend the region of  
Rossby wave excitation by the MJO further eastward as well.

~~Beyond MJO's precipitation intensity, propagation speed, and propagation extent, the MJO's zonal wavenumber is projected~~  
90 ~~to decrease (Rushley et al., 2019). Previous work has also found an increase in the frequency of MJO events (e.g., Arnold et al., 2013; Chang~~  
~~. Lastly, the meridional extent of the MJO may also change, although it is unclear if this is expected with surface warming.~~

Studies exploring the MJO teleconnection change with warming in global climate models are sparse. Samarasinghe et al. (2021) find strengthened future MJO teleconnections over the Gulf of Alaska and North Atlantic in a climate model forced with high emissions. While Zhou et al. (2020) show that a multi-model mean ~~MJO teleconnection change over change in the MJO's~~ impact on both the circulation and precipitation over most of North America is near-zero, they do not quantify the uncertainty in this projection over this larger region. Their results for the North American west coast, which rely on a subset of the publicly available climate model data (they analyze only those models that produce the most realistic MJOs), hint that while the multi-model mean change over broader North America is small, there may be quite a bit of disagreement between models.

~~The importance of the MJO's teleconnection for making predictions on extended timescales, combined with the MJO teleconnection's complex dependence on a number of climate features makes a~~ A detailed study of potential MJO teleconnection changes is attractive. We are motivated to unpack how and why MJO teleconnections might change under global warming, and here seek to clearly and quantitatively rank the various controls on MJO teleconnection strength changes in a warmer climate to better understand how the MJO teleconnection might change in the future. Additionally, we desire to maximize the resources offered by the latest phase of the Coupled Model Intercomparison Project (CMIP6), by avoiding having to subset only those models producing more realistic MJOs.

In this study, we will use a linear baroclinic model (LBM, Watanabe and Kimoto, 2000) in concert with output from CMIP6 to ~~test the hypothesis that the uncertainty in projections of the boreal winter MJO teleconnection to the North Pacific and North America region is large, and to separately quantify the contribution to this uncertainty by various mechanisms~~ separately quantify the contributions to uncertainty in future projections of MJO teleconnection by various mechanisms: mean state changes and changes to the MJO's eastward propagation extent, propagation speed, heating intensity, and zonal wavenumber. The use of the LBM for studying the MJO teleconnection has precedent: ~~Previous work has shown that.~~ Previous work shows the MJO teleconnection is, to first order, ~~and the LBM is a common tool used~~ linear (Mori and Watanabe, 2008; Lin and Brunet, 2018). Many previous studies have used the LBM to untangle MJO teleconnection mechanisms (e.g., Henderson et al., 2017; Tseng et al., 2019, 2020a, b; Wang et al., 2020; Wolding et al., 2017). The linear model additionally has a few features that make it particularly attractive for the current study. First, it is very inexpensive to run, allowing us to cheaply and efficiently run hundreds of perturbation simulations, a task that is difficult and expensive using complex global climate models. Second, the linear framework is quite simple in that almost all simulated variability is ~~due to the addition of~~ from an external forcing, which makes the interpretation of the results very straightforward. Third, because the mean state is maintained by prescribing it at each time step, it is possible to run simulations with the temperature field out of balance with the wind field. This ~~final~~ point allows us to separately and cleanly quantify the uncertainty in the MJO teleconnection change from uncertainty in the thermal structure of the atmosphere versus its mean winds. Finally, because we can separately prescribe an MJO forcing, it is not necessary to have a realistic internally generated MJO when investigating the sensitivity of the MJO teleconnection to inter-model variations in the mean state, thus permitting us to avoid only analyzing those CMIP6 models which produce more reliable MJOs.




We explore the sensitivity of the MJO teleconnection to changes in the MJO’s propagation speed, eastward propagation extent, [zonal wavenumber](#), heating magnitude, and of the mean state dry static stability and winds using simulations with the LBM of Watanabe and Kimoto (2000). The time-integration version of the LBM that we use here is a spectral model that solves linearized equations for vorticity, divergence, temperature, and surface pressure. The model takes two inputs: a mean state and  
 130 a forcing (which is added to the simulated perturbation field at each time step).

The model equations are described in appendix B of Watanabe and Kimoto (2000). [We use a horizontal truncation of T42 \(roughly 2.8° horizontal resolution\) and 20 vertical levels.](#) The model is numerically damped with biharmonic horizontal diffusion, for which we use an e-folding time of 10 minutes for the smallest resolvable waves, and with linear drag, with a damping time scale of 0.5 days applied to the lowest 3 model levels, 1 day for the top two model levels, and 20 days for all levels  
 135 in between. We use this strong damping to inhibit the growth of baroclinic waves, particularly because in some simulations we intentionally prescribe mean state winds that are not in balance with the mean state temperature. ~~We use a horizontal truncation of T42 (roughly 2.8° horizontal resolution) and 20 vertical levels. As a consequence of the strong damping and linear framework, eddy convergences of heat and momentum in the extratropical storm tracks, including those associated with MJO variability (e.g., Deng and Jiang, 2011; Guo et al., 2017; Takahashi and Shirooka, 2014), are not explicitly accounted for.~~  
 140 ~~However, because the mean state wind used in the LBM simulations is taken from CMIP6, eddy-mean flow interactions are, in a way, included implicitly.~~

We will make the convention for composite analysis that “historical” and “future” mean states refer to January conditions during 1984-2014, and 2071-2100, respectively, of the historical and SSP585 simulations from models participating in CMIP6. SSP585 is a relatively high atmospheric greenhouse gas emissions scenario following shared socioeconomic pathway 5 (SSP5;  
 145 O’Neill et al., 2017) and representative concentration pathway 8.5 (Moss et al., 2010). ~~For each CMIP6 model mean state combination, we construct an ensemble of 30 separate LBM simulations, each using a different January. We use interannual ensembles in this way to isolate the climate change signal apart from noise due to internal variability.~~

We restrict our analysis to those 29 CMIP6 models for which there was monthly mean data for the variables required as input by the LBM on the data node at <https://esgf-node.llnl.gov/search/cmip6/> at the time of access. We use the r1i1p1f1 ~~variant~~  
 150 ~~member~~ for all models except for GISS-E2-1-G (r1i1p1f2), HADGEM3-GC31-LL (r1i1p1f3), UKESM1-0-LL (r1i1p1f2), and MIROC-ES2L (r1i1p1f2). ~~We acknowledge that sampling ensemble spread is desirable in order to isolate the forced response from internal variability. Thus, for each CMIP6 model mean state combination, we construct an ensemble of 30 separate LBM simulations, each using a different January. We use interannual ensembles in this way, rather than using the other ensemble members from each CMIP6 model, to minimize the data downloading burden and to keep the number of ensemble members used for each CMIP6 model mean state consistent.~~ Lastly, the SSP585 simulation of CAMS-CSM1-0 was only carried out until  
 155 2099, and so for this model we use only 29 ~~ensemble members~~ years.

We force the LBM with a propagating idealized horizontal dipole heating, which is preferable to a realistic heat source (for example, one obtained from composites of observed MJO events), since it allows ~~easily~~ ~~easy~~ manipulation of the MJO’s inten-



**Figure 1.** Propagating thermal forcing used in the LBM simulations at  $t =$  (a) 10 days, (b) 20 days, (c) 30 days, and (d) 40 days.

sity and propagation characteristics. The heating-idealized heating (as in Ting and Held, 1990; Seo and Son, 2012) is designed to be MJO-like, with the rate-of-change-of perturbation temperature given by

$$\frac{\partial T'(\lambda, \phi, z, t)}{\partial t} T'(\lambda, \phi, z, t) = A e^{-\frac{\sin^2 \phi}{2\sigma^2}} \sin(k\lambda - \omega t) e^{-\frac{(\lambda-b)^2}{2c^2}} \sin^2\left(\frac{\pi(z - z_{top})}{z_{bot} - z_{top}}\right), \quad (1)$$

where  $T'$  is the perturbation temperature of the forcing,  $\phi$  and  $\lambda$  are the latitude and longitude, respectively,  $z$  is the model vertical coordinate with subscripts  $bot$  and  $top$  referring to the lowest and highest model level, respectively,  $A$  is the heating amplitude,  $k$  is the zonal wavenumber,  $\omega$  is the temporal frequency of the heating, and  $t$  is the time. Unless otherwise noted, we set  $A = 0.1 \text{ K day}^{-1}$ ,  $\sigma = 0.1$ ,  $k = 1.9$ ,  $\omega$  corresponding to a temporal period of 40 days,  $b = 130^\circ$ , and  $c = 45^\circ$ . Figure 1 shows the heating-perturbation heating ( $T'$ ) in the mid-troposphere where it is a maximum for  $t = 10$  days, 20 days, 30 days, and 40 days. To impose a realistic MJO horizontal scale, and facilitate varying it, we constrain the propagating heat source to the longitudes over the Indian and West Pacific Oceans by multiplying the sinusoidal heat source by a Gaussian centered over the Maritime continent (see Eq. 1). We initialize the model with the forcing given by setting  $t = 1$  day, and integrate for 60 days.

The experiments are described in Table 1. When testing the sensitivity of the MJO teleconnection to perturbations of the mean state or of an MJO intensity or propagation characteristic, we perturb only the feature being tested and hold everything else constant. For each perturbed feature, we run two sets of experiments: one with a-mean-state-the mean climate given by historical values, and the other with the mean state-climate given by future values (see the “sub-experiment” column in Table 1). For example, in testing the sensitivity of the teleconnection to the mean state winds, we run two sets of experiments: one set in which the dry static stability-energy is held constant at historical values while the winds are perturbed, and the other with future-dry-static-stability set with the dry static energy held constant at future values while the winds are again perturbed. The LBM takes as input three-dimensional winds, temperature, geopotential height, and surface pressure. The set of variables

held constant when the "winds" are held constant are the zonal, meridional, and vertical wind. Similarly, the set of variables held constant when the "dry static energy" is held constant are the temperature, geopotential height, and surface pressure. The use of the many CMIP6 mean states permits the quantification of uncertainty in the future MJO teleconnection due to "model uncertainty" (e.g., Lehner et al., 2020) of the mean state.

We are guided by the results of previous studies for the experiments where the full mean state is held constant and MJO intensity or propagation characteristics are being perturbed. For these simulations, we aim to quantify uncertainty by obtaining the range of possible changes to the MJO teleconnection that result from changes to the MJO. We do this by conducting experiments with perturbations representing the lower and upper bound of changes to the MJO that are ~~expected in a future warmer climate. For climate models that simulate realistic MJOs~~ simulated by models participating in the Coupled Model Intercomparison Project Phase 5 (CMIP5) at the end of the century given high emissions. For climate models that simulate realistic MJOs in CMIP5, Rushley et al. (2019) find an increase in MJO propagation speed between 1.8 and 4.5 %  $\text{K}^{-1}$ , and a maximum decrease of the zonal wavenumber of about 0.2 (we use no change in the zonal wavenumber as a lower bound). Using the increase of the multi-model mean, global mean surface temperature between the historical and future climates of the models used in this study, we thus conduct experiments where the propagation speed of our idealized forcing is perturbed to correspond to an MJO period of about 37.5 (34) days for the lower bound (upper bound) experiment where sensitivity to MJO propagation speed is being tested. In modification of the MJO's eastward extent, we use no change as a lower bound and for the upper bound, extend by  $20^\circ$  the eastern edge of the gaussian used to confine the propagating forcing to the Indian and West Pacific Oceans. Specifically, the gaussian multiplier of the perturbation forcing is the control gaussian with the maximum value sustained for an additional  $20^\circ$  to the east before decreasing. This value is roughly informed from results in Maloney et al. (2019) (CMIP5) and in Zhou et al. (2020) (CMIP5 and CMIP6). Lastly, we are informed by Bui and Maloney (2018), who find that the subset of CMIP5 models that produce MJOs validating best against present-day observations tend to simulate changes of the MJO's precipitation amplitude between -10% to + 20% between the historical and future climates under RCP8.5 forcing. In the experiment testing sensitivity to MJO amplitude, we use perturbed MJO heating amplitudes of 0.9 and 0.12  $\text{K day}^{-1}$ , corresponding to a 10% reduction and 20% amplification of the control forcing. We use estimates of changes to the MJO from CMIP5, rather than CMIP6, due to the availability of published, detailed analyses of MJO changes for CMIP5. However, we note that despite recent improvement in the simulation of the MJO in CMIP6 (Ahn et al., 2020) much work is needed to deepen understanding of future changes to the MJO, and thus these estimates are crude, first guesses. We include a figure showing the perturbed MJO cases in Appendix A.

With the described setup, some of the simulations became unstable. We readily admit this is an inescapable trade-off of having ~~assumed~~ implemented a forced linear model framework; despite its advantages for rapidly sampling and intercomparing first-order effects, the neglect of secondary nonlinear buffering mechanisms can corrupt some integrations. Pragmatically, we thus omit all LBM simulations in which, at any location, the maximum value of geopotential height at 500 hPa during the second half of the integration is more than ten times the maximum value during the first half. Additionally, we omit all LBM simulations using a specific CMIP6 model's mean state if 10 or more of the 30 constituent ensemble simulations for any of the mean state combinations meets the instability criterion described above, leaving 24 of the 29 CMIP6 models for analysis.

**Table 1.** Experimental setup for the linear baroclinic model simulations. “Lower bound” and “upper bound” refer only to perturbations of the thermal forcing (last three rows). DSE = dry static energy.

Feature	Sub-experiment	Control	Perturbation (lower bound)	(upper bound)	Number of unique simulations	Number of unstable simulations
Mean state DSE	Historical wind	Historical DSE	Future DSE		1440	49
	Future wind				1440	63
Mean state winds	Historical DSE	Historical wind	Future wind			
	Future DSE					
Forcing heating magnitude	Historical mean state	$A = 0.10 \text{ K day}^{-1}$	$A = 0.09 \text{ K day}^{-1}$	$A = 0.12 \text{ K day}^{-1}$	60	0
	Future mean state				60	0
Forcing propagation speed	Historical mean state	$\omega = 1/40 \text{ day}^{-1}$	$\omega = 1/37.5 \text{ day}^{-1}$	$\omega = 1/34 \text{ day}^{-1}$	600	0
	Future mean state				600	0
Forcing eastward extent	Historical mean state		$0^\circ$ extension	$20^\circ$ extension	600	0
	Future mean state				600	1
Forcing wavenumber	Historical mean state	$k = 1.9$	$k = 1.9$ (no change)	$k = 1.7$	600	2
	Future mean state				600	3

Figure A2 in Appendix A shows, for each basic state combination, the fraction of ensemble members that became unstable. The last two rows of Table 1 list the number of total simulations run for this study (minus those from the five CMIP6 models we omit) and the number of simulations we omit from our analysis for each experiment (which is reassuringly less than 5% of the total simulations for each experiment). Note that the “number of unique simulations” field is blank for the mean state winds feature (second row) because the simulations are shared with that of mean state dry static stability (first row).

Lastly, we emphasize that the linear framework we are using neglects nonlinear interactions in simulating MJO teleconnections. For example, in reality, the mean state exerts a strong control on MJO propagation characteristics (Jiang et al., 2020). Rossby waves excited by the MJO are able to extract energy from the mean flow (Adames and Wallace, 2014; Zheng and Chang, 2020), and nonlinear interactions have been shown to lead to spatial shifts in extratropical teleconnection patterns (Lin and Brunet, 2018). Thus, we caution that the results of this study are limited by the exclusion of these effects.


### 3 Results

#### 3.1 Mean state

We conduct experiments with the experimental setup described by Table 1.


Reassuringly, the propagating MJO-like forcing in the model excites a plausible time-varying response in the extratropics. For the set of 30 simulations performed with the full mean state given by historical values from a single representative CMIP6 model (ACCESS-ESM1-5) As an example, Fig. 2a shows the time-varying response of 850 hPa meridional wind at 52500 hPa geopotential height at 60° N, 120115° W for each ensemble member (thin light blue lines) and their mean (thick black line) for the set of 30 simulations performed with the full mean state given by historical values from a single representative CMIP6 model (ACCESS-ESM1-5). The smallness of the meridional-wind-geopotential height anomalies relative to a typical observed midlatitude value is an expected consequence of the smallness of the amplitude used to force the model, which we kept small to try to minimize the number of unstable simulations.

We next introduce a scalar metric of MJO teleconnection strength that is appropriate to visualize in map form, recognizing that the magnitude of peaks and troughs of the ensemble mean response to the propagating forcing (for example, the thick black line in Fig. 2a) is one way to quantify the magnitude of the consistent (ensemble mean signal that stands apart from the noise due to internal variability) teleconnection strength. Given that the amplitude of an infinite sine or cosine wave is the square root of twice its variance, we calculate-define the teleconnection strength at each point as the square root of twice the temporal variance of the ensemble mean meridional-wind-at-850-geopotential height at hPa, ignoring the first 10 days of each simulation to avoid sampling conditions prior to the establishment of representative spread between interannually varying ensemble members. That is, we take the ensemble mean first and then calculate the variance. We will refer to this value as the “amplitude” of the modeled MJO teleconnection. This metric is qualitatively similar to the STRIPES index of Jenney et al. (2019), but is more simple because the forcing is identical between each simulation in an ensemble set, and because the propagating forcing is the sole source of variability in the LBM simulations. We use low-level meridional wind in this calculation (instead of other dynamically relevant variables like mid-tropospheric geopotential height, for example);



**Figure 2.** (a) The time series of meridional wind geopotential height at 850-500 hPa at 5260° N, 120115° W for each January and their mean (thick black line) for the set of LBM simulations with the historical mean state winds and dry static energy (DSE) from ACCESS-ESM1-5. (b) The multi-model mean of the square root of twice the variance of each model's ensemble mean meridional wind at 850 hPa teleconnection amplitude (see text for definition) for the set of simulations with mean states given by historical winds and DSE. (c) The difference between the multi-model mean teleconnection amplitudes for the set of simulations with the full historical mean state and the set of simulations with future winds and historical DSE dry static energy, (d) historical winds and future DSE dry static energy, and (e) future winds and DSE future dry static energy. The blue "x" in panel (b) is the location shown in panel (a). The white lines in panels (b)-(d) indicate the  $5 \times 10^{-3}$  contour. DSE = dry static energy.

because the MJO primarily drives surface temperature variability across North America via low-level temperature advection (Seo et al., 2016), and may thus be more relevant for quantifying changes to the MJO's impact on near-surface midlatitude weather in a future climate which is useful for quantifying the sensitivity of the extratropics to MJO variability across all MJO phases.



**Figure 3.** The change in the tropical mean (20°S-20°N) vertical temperature gradient ( $\partial T/\partial p$ ) between the historical and future periods for each CMIP6 model used in this study.

The baseline multi-model MJO teleconnection simulated by the LBM looks reassuringly plausible. Figure 2b shows the multi-model mean MJO teleconnection amplitude for the set of LBM simulations performed with both the mean state dry static energy (the vertical gradient of which gives the dry static stability) and winds given by historical values. Darker colors indicate that MJO teleconnectivity is larger. The MJO-like forcing excites Rossby waves that propagate in high density through the Pacific waveguide, leading to large values of teleconnectivity there. The pattern of MJO teleconnectivity over North America, which exhibits maxima over the southeastern and northwestern region, with a minimum over the central region, is consistent with the pattern of the observed boreal winter MJO influence on North American **temperature** (e.g., Jenney et al., 2019; Zhou et al., 2012) geopotential height, temperature, and sea level pressure (e.g., Jenney et al., 2019; Zhou et al., 2012).

260 The remainder of Fig. 2 reveals an overall, in general, a reduction of MJO teleconnectivity due to future mean state effects, decomposed into separate contributions from stability vs. winds. Figure 2c-e shows the multi-model difference in the amplitude of the modeled MJO teleconnection but for between the set of simulations performed by perturbing the mean state with the historical mean state and the set of simulations with future perturbations to the mean state. The forcing used in all simulations is the same. **The difference between the top and bottom rows is the sensitivity of the MJO teleconnection to the change in the thermal structure of the atmosphere (i.e., its dry static stability), alone. The pattern of the MJO teleconnection is the same between each top panel and the panel below it, but the teleconnection is weaker in the simulations with future dry static energy because the higher static stability weakens the MJO circulation and subsequent Rossby wave source (Sardeshmukh and Hoskins, 1988)**


265



Panel (c) shows the change to the multi-model mean teleconnection that results from future changes to the winds, alone. Over most of North America, the change in the prescribed thermal forcing. The difference between the left column and the right column is the sensitivity of the MJO teleconnection amplitude to the simulated change in mean winds between the historical and future climates of CMIP6. For most regions, the change to the mean state winds between the historical and future climates leads to a reduction of the MJO teleconnection amplitude. This reduction is particularly apparent over the central Pacific. teleconnection is near-zero. Over the eastern Pacific and Bering Sea, changes to the winds lead to a weaker teleconnection. Other studies have noted an increase in the MJO's boreal winter influence near California (Zhou et al., 2020), attributing this to an eastward extension of the Pacific jet. Here, we find a similar strengthening over this region. In Fig. 2c-e, we include a white line highlighting the  $5 \times 10^{-3} \text{ m s}^{-1}$  contour to help show this strengthening, which is apparent as an eastward extension of this contour line towards the California coast. This strengthening is not apparent here. Panel (d) shows the change to the multi-model mean teleconnection that results from future increases in tropical dry static stability, alone. That is, changes to the winds are not taken into account and are held constant between these two sets of simulations. The teleconnection is weaker in the simulations with future dry static energy because the higher static stability weakens the MJO circulation and subsequent Rossby wave source (Sardeshmukh and Hoskins, 1988) that results from the prescribed thermal forcing.

The difference between panels (b) and (Panel (e) of Fig. 2 is shows the sensitivity of the MJO teleconnection to a change in the full mean state, neglecting any changes to the MJO. In isolation (apart from changes to the MJO), changes to the mean state weaken the multi-model mean teleconnection amplitude almost everywhere due to both an intensification of the in places where the teleconnection is the strongest. This is mostly due to an increase in the tropical dry static stability and (panel d), but over the Eastern Pacific is also due to changes to the winds (panel c). Figure 3, which shows the change in the tropical vertical temperature gradient for individual CMIP6 models between the historical and future periods, confirms that models show a robust increase in the tropical dry static stability.

We now explore the inter-model spread of changes to the MJO teleconnection due to changes in the mean state. Figure 4a,c summarizes, for the separate mean states from each CMIP6 model, the change in the LBM-simulated teleconnection amplitude over the North Pacific (top panels;  $150^{\circ}$ - $235^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N) and North America (bottom panels;  $25^{\circ}$ - $70^{\circ}$  N and  $235^{\circ}$ - $290^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N), as a percent change per Kelvin multi-model mean warming. Here and throughout the remainder of this work, we choose to separate the larger Pacific-North America region into two smaller regions because we do not want to assume that the amplitude changes in the two regions are the same. For each region we plot the domain-mean percent change in the teleconnection amplitude, as well as the fractional area that is strengthening for perturbations to either the mean state dry static energy or the mean state wind. By fractional area, we mean the fraction of the region that shows a teleconnection amplitude change greater than zero. We include the fractional area as a crude way to communicate the spatial distribution of teleconnection changes over the region. For example, in, where a value of 0.5 indicates that 50% of the domain shows stronger teleconnections in the future period compared to the historical period. Likewise, a value of 1 indicates that the entire domain shows a larger teleconnection amplitude in the future period. In isolation, the mean teleconnection amplitude change over the region may be misleading if one small region experiences a very strong change in one direction (enough to make the domain mean change be of the same sign), despite the majority of the domain experiencing a change of the opposite direction. For



**Figure 4.** For individual CMIP6 model mean states and the multi-model mean, the change in the teleconnection amplitude (as a percent change per multi-model mean Kelvin warming) to the North Pacific ( $150^{\circ}$ - $235^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N) and North America ( $235^{\circ}$ - $290^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N) between LBM simulations with mean states given by future and historical quantities. The isolated change due to (a) stability, over the North Pacific; (b) wind, over the North Pacific; (c) stability, over North America; (d) wind, over North America. Horizontal axis is the domain mean change over the region, while the vertical axis shows the fractional area of teleconnection amplitude strengthening. DSE = dry static energy.

example, ~~there are a few models in panel b which lie to the right of the vertical zero line (indicating a mean positive change in the teleconnection amplitude over~~ we find that there are some cases in which the North Pacific due to the winds alone) and ~~to the bottom of the horizontal zero line (indicating that a majority fraction mean teleconnection amplitude change over the region is positive, while the majority area~~ of the region shows a ~~weakening~~ negative change.

The increase in the tropical dry static stability is a consequence of amplified upper-tropospheric warming that occurs as tropical temperatures adjust to warmer surface temperatures (e.g., Santer et al., 2005). Our first main finding is strong corroboration of the hypothesis that the increase of the tropical dry static stability is a robust thermodynamic response of the climate to warming that tends to reduce the MJO's teleconnections in almost all models (Fig. 4a,c). This is the first multi-model confirmation of this hypothesis, which first appears in the single-model study of Wolding et al. (2017). There is relatively good agreement in the teleconnection amplitude change due to the change in the atmosphere's thermal structure alone (i.e., neglecting mean state wind or MJO changes) between model mean states, particularly over the North Pacific. Comparison between panels a and c of Fig. 4 shows that there is greater spread downstream of the tropical heat source over North America, although reasons for this are unclear.

Despite agreeing on effects of stability, models disagree wildly on the effects of future wind changes. Figure 4b,d show the change in the teleconnection amplitude due to the change in the wind between the historical and future period. The multi-model mean response of the MJO teleconnection amplitude over the North Pacific is a modest weakening about zero, with a majority

of the region ~~also~~ experiencing a weakening. However, the inter-model spread is relatively large, with models showing changes between ~~-15-11~~ to  $7 \text{ \% K}^{-1}$  multi-model mean warming. Over North America, the multi-model mean change in the teleconnection due to the wind is a ~~small-strengthening, with most of the region showing weaker teleconnections~~ strengthening of  $4.7 \text{ \% K}^{-1}$ . But again, the inter-model spread is large (~~-9 to -12-3 to 19~~  $\text{ \% K}^{-1}$ ) ~~with the multi-model mean is indistinguishable from zero as was found in Zhou et al. (2020)~~.

In summary, three key results emerge concerning how mean state changes in a future warmer climate ~~will~~ may contribute to changing MJO teleconnections to the North Pacific and North America. First, we find robust support for the expectation that increases in tropical dry static stability will contribute to weaker MJO teleconnections. Second, we find that changes to the mean state winds can contribute to very large changes in the MJO teleconnection; that is, large enough to as much as double or negate the weakening that is expected due to increasing tropical dry static stability. Third, there is a large amount of inter-model spread in LBM-simulated changes to the MJO teleconnection that result from changes to mean state winds, which suggests that uncertainty in the response of the mean state winds to warming ~~dominates~~ is a large contributor to uncertainty in modeled changes to the MJO teleconnection.


It is thus important to better understand the cause of teleconnection changes associated with mean state wind changes in the CMIP6 models. Mean state winds can increase or decrease the strength of MJO teleconnections via an intensification of Rossby wave excitation and/or through mean state changes that permit more Rossby wave energy propagation into these regions. ~~The following analysis will argue that the latter—wind-induced changes to Rossby wave ducting—are more important than changes to wave excitation in producing the inter-model teleconnection spread.~~

~~To show this, we~~ We begin by quantifying the strength of Rossby wave excitation using the linearized equation for the “Rossby wave source” (Sardeshmukh and Hoskins, 1988),  $S'$ , a term that represents the generation of large-scale vorticity,

$$S' = - [\nabla \cdot (\overline{\mathbf{v}_\chi} \zeta') + \nabla \cdot (\mathbf{v}_\chi' \overline{\zeta})] \quad (2)$$

where  $\zeta$  is the absolute vorticity,  $\mathbf{v}_\chi$  is the horizontal divergent wind, the overbar represents mean state quantities, and the prime represents anomalies. Here, anomalies are the LBM-simulated deviations from the user-input mean state. In the subsequent analysis, we use quantities at 200 hPa to calculate  $S'$ .


We do not find a systematic relationship between the difference in  $S'$  and the difference in the mean teleconnection amplitude over the North Pacific or North America between simulations with historical and future winds. Figure 5 shows, for simulations with historical and future winds, the relationship between the ensemble mean, time mean, difference in the absolute value of the Rossby wave source averaged over the subtropical jet ( $80^\circ$ - $190^\circ$  E,  $20^\circ$ - $40^\circ$  N), and the difference in the mean teleconnection amplitude over the North Pacific and North America. Figure A3 in Appendix A verifies that this region indeed encloses the region of highest  $S'$  variability in the LBM simulations. Models with larger increases in  $|S'|$  over the jet do not systematically have stronger teleconnections. If changes to Rossby wave excitation were to play a key role in the difference in the teleconnection amplitude between simulations with historical and future mean state winds, then it would be reasonable to expect either a weaker or similar-sized link over North America than over the North Pacific, since most Rossby



**Figure 5.** For individual CMIP6 model mean states, the difference in the Rossby wave source ( $S'$ ) versus the difference in the teleconnection amplitude between simulations with historical and future mean state winds for (a) the North Pacific and (b) North America. The horizontal axis is the ensemble-mean, time-mean difference of the absolute value of  $S'$  averaged over the subtropical jet (80°-190° E, 20°-40° N).

waves excited by the MJO propagate first over the Pacific before reaching North America. However, the relationship between the difference in  $|S'|$  and the teleconnection amplitude difference appears stronger over North America ( $r = 0.43$ ) than over the North Pacific ( $r = 0.27$ ). Thus we conclude that the change in  $|S'|$  is not a primary mechanism that explains how mean state winds affect the teleconnection amplitude difference in our LBM simulations. [This is consistent with Wang et al. \(2020\) and Zheng and Chang \(2020\) who show that a stronger  \$S'\$  does not necessarily lead to stronger MJO teleconnections.](#)

In addition to playing a key role in Rossby wave excitation, mean state winds also determine the path that Rossby waves take as they propagate (e.g., Karoly, 1983; Hoskins and Ambrizzi, 1993). [Meridional shifts and variations in the zonal extent of the subtropical jet have been linked with variations in teleconnection amplitude across North America \(Henderson et al., 2017; Tseng et al., 2020\).](#) [An eastward extension of the subtropical jet was shown to be important for increases in the MJO's impact over the North American west coast in climate models \(Zhou et al., 2020\).](#) [Similarly, Samarasinghe et al. \(2021\) propose that the changes to the North Pacific jet may explain modeled increases in simulated MJO teleconnections in CESM2-WACCM.](#) Here, we search for a systematic (common across CMIP6 model mean states) and localized (specific to one region) change in the upper-tropospheric extratropical mean flow that explains variations in the teleconnection amplitude (inter-model spread in [Figure 4b,d](#)). We begin by looking for a link between the change in the extratropical zonal wind at 200 hPa and the teleconnection amplitude response to changes to the winds. [Figure 6a,c shows the correlation coefficient between the local change in the 200 hPa zonal wind \(the difference in the January mean zonal wind between the historical and future periods\) and the regional mean \(i.e., across the entire North Pacific or North America region\) difference in the MJO teleconnection amplitude between simulations with historical and future mean winds \(individual modeled changes in 200 hPa zonal wind are shown in \[Figure A4 in Appendix A\]\(#\)\).](#) Correlations are generally weak everywhere (for reference,  $r = 0.5$  corresponds to a p-value of roughly = 0.30 using a Fisher Z-transformation), which suggests that there may not be a change to the winds over a specific region that



**Figure 6.** Correlation coefficient between the local change in the 200 hPa zonal wind and the regional mean change in the teleconnection amplitude over (a) the North Pacific and (c) North America. (b) Correlation coefficient between the local change in the stationary Rossby wave number,  $K_s$ , and the regional mean change in the teleconnection amplitude over the North Pacific and (d) North America. Grey boxes outline the North Pacific and North America regions.

explains a majority of the spread in Figure 4b,d. However, given the importance of horizontal gradients of mean state vorticity for Rossby wave propagation, such a link may require more work to unveil.

Analyses of the stationary Rossby wave number have been useful for assessing how the mean state winds affect Rossby wave propagation (e.g., Henderson et al., 2017; Karoly, 1983; Tseng et al., 2020b; Wang et al., 2020) (e.g., Henderson et al., 2017; Karoly, 1983). The stationary Rossby wave number on a Mercator projection (to account for spherical geometry) is defined as

$$K_s = a \left( \frac{\beta_M}{u_M} \right)^{1/2}, \quad (3)$$

where  $\beta_M$  is the meridional gradient of absolute vorticity on a Mercator projection,

$$\beta_M = \frac{2\Omega \cos^2 \theta}{a} - \frac{\partial}{\partial y} \left[ \frac{1}{\cos^2 \theta} \frac{\partial}{\partial y} (u_M \cos^2 \theta) \right], \quad (4)$$

and  $u_M = u / \cos \theta$  is the mean-temporal mean, full (including summetric and asymmetric components) zonal wind divided by the cosine of latitude ( $\theta$ ). In Eqs. (3) and (4),  $a$  is the earth's radius and  $\Omega$  is the earth's rotation rate. Hoskins and Ambrizzi (1993) showed that  $K_s$  can be used to understand Rossby wave propagation: waves turn towards regions of higher  $K_s$ , are generally reflected away from regions where the zonal Rossby wavelength is equal to  $K_s$  or where  $\beta_M \leq 0$ , and are either

dissipated beyond or reflected from regions where  $u \leq 0$ . Regions of  $K_s$  maxima, which tend to occur in strong westerly jets, act  
385 as Rossby waveguides. ~~Figure 6a shows  $K_s$  computed from the multi-model mean January wind averaged across the historical period. Regions with  $\beta_M \leq 0$  are colored black, and regions with  $u \leq 0$  are colored white. Multiple previous studies have used the stationary Rossby wave number in this form (i.e., neglecting meridional wind) when exploring the impact of the mean state winds on MJO teleconnections (Henderson et al., 2017; Tseng et al., 2020b; Wang et al., 2020; Zheng and Chang, 2020).~~

~~(a) Stationary Rossby wave number,  $K_s$ , computed from the multi-model mean wind over the historical period. (b) The  
390 correlation coefficient between the local difference in  $K_s$  between the historical and future periods, averaged over a  $15^\circ \times 15^\circ$  box, and the average teleconnection amplitude change across the North Pacific and (c) North America. Black boxes in (b), (c) outline the North Pacific and North American regions, respectively.~~

We now inspect if localized changes to  $K_s$  at 200 hPa in the CMIP6 model January basic states can systematically explain the spread in Figure 4b,d. An increase in the local  $K_s$  indicates that Rossby waves with shorter wavelengths are able to propagate  
395 further before turning. Similarly, a decrease in the local  $K_s$  means that propagating Rossby waves will turn sooner. A change to the meridional width of the waveguide, which presents as an increase or decrease in  $K_s$  along the north and/or south flanks of the waveguide, affects the trajectory of Rossby waves traveling within the waveguide, with Rossby waves able to travel further meridionally and turn less often in a broader waveguide (Hoskins and Ambrizzi, 1993). Thus, the path that Rossby waves take as they exit the jet is sensitive to both the width of the waveguide and the value of  $K_s$ . Figure 6b,e-d show the correlation  
400 coefficient between the changes in the local  $K_s$  and the regional mean teleconnection amplitude change for the North Pacific and North America, respectively. ~~(for reference, we include the difference in  $K_s$  between the historical and future period for each model in Appendix A). We include boxes outlining the North Pacific and North America regions for reference.~~ In the calculation of the correlation, we use averages of the local  $K_s$  change over  ~~$15^\circ \times 15^\circ$~~   $10^\circ \times 10^\circ$  boxes to focus on the larger features; although using boxes of different sizes ~~(we tried boxes ranging in size from  $3^\circ$ - $20^\circ$ )~~ does not qualitatively change  
405 the results. Regions of ~~strong positive correlation suggest~~ stronger positive correlation would be suggestive of a systematic (common across models) positive relationship between the local change in  $K_s$  and the regional (i.e., across North Pacific or North America) teleconnection amplitude ~~change~~ response to the changes in the winds projected by CMIP6. Similarly, regions of strong negative correlation would suggest a systematic inverse relationship between the local change in  $K_s$  and the regional teleconnection change. ~~We include boxes outlining the North Pacific and North America regions for reference. Correlations~~  
410 ~~linking regional changes in the local  $K_s$  with the regional mean teleconnection amplitude change are generally weaker for the North Pacific than for North America. We thus focus on North America, but point out that the two maps have similar patterns, suggesting that changes to the winds leading to stronger teleconnections over North America may be in part due to changes in the winds that also support stronger teleconnections over the North Pacific. Figure 6c shows a region of high positive correlation over the Gulf of Alaska. Here, models exhibiting an increase in  $K_s$  tend to have an increase in the teleconnection over North~~  
415 ~~America, and vice versa. Additionally, there is a region of negative correlation just to the south of this area. We interpret this as a slight northward shift and broadening along the northward flank of the Pacific waveguide, and subsequent increase in the number of Rossby waves traveling northward towards North America, in the models that have stronger MJO teleconnections. This view is consistent with the work of Tseng et al. (2020b), who demonstrate how increased  $K_s$  over the Gulf of Alaska~~

and resultant increases in northward Rossby wave propagation out of the Pacific waveguide during La Niña years leads to an enhanced MJO teleconnection over North America.

We thus identify a systematic mechanism that may explain how mean state wind changes lead to changes to teleconnectivity over North the North Pacific or North America across CMIP6 models. This suggests that increased confidence in how winds over this region will change in a future climate may help reduce uncertainty in future estimates of the change in the strength of MJO teleconnections to North America. We would like to point out, however, that changes in the winds over this region alone are likely not the sole important change. That is, for some models, it is possible that changes to  $S'$  may be important, as well. More work is needed to understand how changes to the mean flow will impact future MJO teleconnections, which our results indicate are likely a large source of inter-model spread in future projections of MJO teleconnections. These results suggest it may not be sufficient to focus on local wind changes over only one or two regions.


### 3.2 MJO intensity and propagation characteristics

We now investigate how changes to MJO intensity and propagation characteristics affect the MJO teleconnection by conducting simulations in which the mean state is held constant, and either the propagation speed, eastward propagation extent, zonal wavenumber, or intensity of the idealized thermal forcing are perturbed to upper and lower bounds of expected end-of-century changes to the MJO. Instead of using mean states from all CMIP6 models, we use a subset of mean states from 10 models to minimize utilization of computational resources. For these experiments, we chose CMIP6 models that produced a minimum number of unstable ensemble members in experiments used in Sect. 3.1 (see Fig. A2).

Figure 7 summarizes the results of these simulations. As in Figure 4, the horizontal axis shows the regional mean difference in the teleconnection amplitude while the vertical axis shows the fractional area of the region that has stronger teleconnections (perturbed MJO compared to control MJO). Panels a and d-e show the change to the teleconnection amplitude over the North Pacific and North America, respectively, that result from perturbations to the magnitude of the propagating thermal forcing. The linearity of the LBM ensures that, in the absence of instability, the magnitude of the simulated response varies linearly with the magnitude of the forcing. We verified this with simulations using a 20% increase in the magnitude of the forcing for two different model basic states. For the remainder of the 8 models used in these experiments, we thus did not actually run simulations (see Table 1), because the magnitude of the extratropical response to the perturbed forcing amplitude is independent of the mean state. If full non-linearity nonlinear interactions had been considered, the variations across the models' mean state different model climates could have also been important in the response of the teleconnection to changes in heating intensity.

Unlike perturbations to the heating intensity, the responses of the MJO teleconnection to perturbations to the eastward propagation speed (Fig. 7b,e) and f), extent (Fig. 7c,f) are linearly g), and zonal wavenumber (Fig. 7d,h) are sensitive to the mean state. In general, climate of each CMIP6 model. Consistent with previous work finding decreased MJO teleconnection strength with increased MJO propagation speed (Zheng and Chang, 2019; Wang et al., 2020; Bladé and Hartmann, 1995), we find that increases to the propagation speed of our idealized heating produce modest decreases in the teleconnection strength amplitude over the North Pacific, with changes over North America centered on and straddling zero change and North America.






**Figure 7.** For individual CMIP6 model mean states, the change in the teleconnection amplitude (as a percent change per multi-model mean Kelvin warming) to the North Pacific ( $150^{\circ}$ - $235^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N) and North America ( $235^{\circ}$ - $290^{\circ}$  E,  $25^{\circ}$ - $70^{\circ}$  N) between LBM simulations with the control and perturbed thermal forcing. (a),(e) Response to perturbed heating intensity; (b),(f), eastward propagation speed; and (c),(g), eastward propagation extent; and (d),(h) zonal wavenumber. Unfilled (filled) markers represent the lower (upper) bounds of perturbations to each MJO feature expected at the end of the century given a high emissions scenario (see Table 1).

Increasing the eastward extent and decreasing the zonal wavenumber of the propagating MJO thermal forcing in the LBM simulations increases the teleconnection amplitude for all mean-states-model climates used and has the potential to produce the largest increases in the MJO teleconnection amplitude, more so over the North America region.

Figure 7 highlights two sources of uncertainty that lead to uncertainty in the response of MJO teleconnections to changes to the MJO itself: uncertainty in changes to the MJO and uncertainty in For increases in the eastward propagation extent, this is likely related to an eastward broadening of the region over which the MJO excites Rossby waves. Adames and Wallace (2014) found that for the mean state -Uncertainty-winds of the period from 1979-2011, the MJO excites the strongest extratropical response when heating maxima are located over the central Pacific. While we separately tested the sensitivity of the extratropical response to increases in the MJO's response to warming (here, between a 10% overall decrease or 20% overall increase between the present-day and end of the 21st-century), leads to a range of about  $8\% K^{-1}$  uncertainty in the change to propagation speed and decreases to the MJO teleconnection zonal wavenumber, these two changes to the MJO are linked and will likely occur together (Rushley et al., 2019). These two changes to the North Pacific and North America. For the MJO have opposite impacts on MJO teleconnection amplitude, with the net impact appearing to be a slight strengthening from the larger strengthening effect from the decreased zonal wavenumber.

For perturbations to the MJO's propagation characteristics, both uncertainty in the mean state and uncertainty in the MJO's response to warming leads to uncertainty in the teleconnection response to warming, with a range of about  $5\% K^{-1}$  from changes Fig. 7 highlights the sensitivity of the MJO teleconnection change to the mean climate of each CMIP6 model. That is, for a given perturbation to the MJO's propagation speed and up to about  $10\% K^{-1}$  from the MJO's eastward propagation extent, zonal wavenumber, or eastward extent, inter-model differences in the mean climate lead to variations in the teleconnection



**Figure 8.** The total change to the MJO teleconnection over (a) the North Pacific and (b) North America due to the change in the full mean state (markers), and the potential range that results from considering changes to MJO propagation and intensity characteristics (horizontal lines extending from each marker) for the 10 models for which we ran perturbed MJO experiments.

475 change. Thus, not only are changes in the mean climate important for understanding how the extratropical MJO teleconnection will evolve for an unchanging MJO, but also for understanding how MJO teleconnections will evolve in response to changes to the MJO. Additionally, for each individual MJO propagation or intensity characteristic, the uncertainty spread in the MJO teleconnection change resulting from a perturbed MJO is on the order of the model-inter-model uncertainty that results from perturbations to the mean state alone (Fig. 4). This further challenges These results challenge the notion that it may currently be possible to have currently have confidence in future projections of MJO teleconnections.

### 3.3 Sum of mean state and MJO

Putting it together, we finish by quantifying ~~an overall decrease~~ the overall change in the MJO teleconnection that results when  
480 considering the linear sum of individual changes to the mean state and MJO features. For each CMIP6 model, Fig. 8 shows  
how the change to the full mean state influences the change in the MJO teleconnection to the North Pacific and North America,  
as well as the the range of teleconnection changes that results from the sum of changes to the mean state and MJO propagation  
and intensity characteristics for the 10 CMIP6 mean states for which we also ran perturbed thermal forcing simulations. Over  
the North Pacific, for all CMIP6 models, changes to the mean state alone lead to ~~an overwhelmingly weaker teleconnection~~  
485 a weaker teleconnection for almost all models, with only ~~one model~~ two models (AWI-CM-1-1-MR, EC-EARTH3-VEG)  
showing a small strengthening. ~~When considering also how~~ Factoring in changes to the MJO ~~may factor in, another model~~  
~~joins in the potential for intensifying teleconnection amplitude. Nonetheless, these~~ leads to a range of potential changes for  
~~many models that straddles zero. These~~ results suggest that ~~it is likely that~~ over the North Pacific, ~~in a future warmer climate,~~  
~~MJO teleconnections will weaken~~ mean state changes may lead to weaker MJO teleconnections. However, a strengthening of  
490 the MJO teleconnection over the North Pacific, due to changes to the MJO (from potential increases in eastward propagation  
extent or heating intensity and decreases in zonal wavenumber), can not be ruled out.

Over North America, ~~the change there is no agreement on how changes~~ in the mean state alone ~~also leads to decreases in the~~  
~~teleconnection amplitude for most CMIP6 models, albeit less so than over the North Pacific, with 3 CMIP6 models showing~~  
~~increases to the MJO teleconnection will impact MJO teleconnections: while the median change is near-zero, the range of MJO~~  
495 teleconnection changes due to mean state changes spans from -11 % K<sup>-1</sup> to 15 % K<sup>-1</sup>. When changes to the teleconnection  
due to changes to the MJO are also considered, the final picture ~~is more blurry, and shows slightly~~ shows increased potential for  
strengthening ~~in some models. Over North America, it appears as though changes to the MJO are again likely to weaken MJO~~  
~~teleconnections overall, but with some added uncertainty such that one cannot rule out the possibility of stronger linkages;~~  
~~comparison with Fig. 7 suggests that this is from potential increases due to an intensification and increased,~~ again due to the  
500 potential for an increase in the MJO's heating intensity and eastward propagation ~~extent of the forcing,~~ and decrease in its  
zonal wavenumber (Fig. 7).

## 4 Conclusions

The Madden-Julian Oscillation (MJO) is a key source of predictability of extratropical weather on subseasonal-to-seasonal  
timescales. While it is expected that the MJO teleconnection to the extratropics will change in a future warmer climate, models  
505 do not agree (Zhou et al., 2020) over much of North America. Speculatively, this may be due in part to mean state biases or due  
to inter-model spread or biases in the simulation of the MJO, both of which have been shown to lead to biases in the simulated  
MJO teleconnection of the current climate (Henderson et al., 2017; Wang et al., 2020).

The MJO teleconnection is sensitive to a number of factors, including the large-scale wind and thermal structure of the  
atmosphere and MJO propagation and intensity characteristics. While a number of climate models simulate the MJO and  
510 its teleconnections, each model represents a unique combination of multiple changes to the various features of the climate

that impact MJO teleconnections. This makes the determination of causal mechanisms important for the change to the MJO teleconnection a challenge. Additionally, many climate models do not accurately reproduce observed characteristics of the MJO or the mean state, and thus may not be reliable for studying internally simulated MJO teleconnections. Previous studies analyzing MJO teleconnections in climate models often analyze a subset of models that produce better MJOs (e.g., Henderson et al., 2017; Zhou et al., 2020).

Motivated by a desire to both (1) untangle the impacts of individual changes to features of the climate on changes to the MJO teleconnection with warming, and (2) maximize the resources offered by the most recent phase of the Coupled Model Intercomparison Project (CMIP6), we conduct a set of sensitivity experiments with a linear baroclinic model (LBM) of the atmosphere in concert with output from CMIP6. In each LBM experiment, a climate feature (i.e., the mean state or intensity or propagation characteristics of a propagating thermal forcing) is perturbed by a climate change amount; that is, the change that is simulated in climate models by the end of the 21st century in simulations with strong CO<sub>2</sub> forcing. Specifically, we test sensitivity of the boreal winter MJO teleconnection ~~strength-amplitude~~ to the North Pacific and North America to the mean state static stability and winds, and MJO eastward propagation extent, propagation speed, and heating intensity.

As expected, given that the increase of the tropical dry static stability is a robust thermodynamic response to surface warming, we verify that the increase in the tropical dry static stability with warming produces robust decreases in the LBM-simulated MJO teleconnection amplitude. On the other hand, we find relatively large spread in the LBM-simulated teleconnection response to changes in the mean state wind. Nonetheless, when the mean state changes are considered together (Fig. 8), they lead to robust decreases in the MJO teleconnection to the North Pacific. Over North America, the sum of mean state changes leads to a weaker/~~stronger~~ MJO teleconnection for ~~most-about half of~~ CMIP6 model mean states, ~~although with many of these models hovering close to zero change~~.

Investigation into a potential ~~systematic~~-link between changes to the wind and changes to MJO teleconnection revealed ~~a possible no systematic~~ connection related to ~~Rossby wave changes in the Rossby wave source~~. Additionally, we find ~~no systematic and localized connection related to Rossby wave~~ propagation over the ~~eastern Pacific, where models that display increases (decreases) in the stationary wavenumber over the Gulf of Alaska also show increases (decreases) in the extratropics~~. Thus, while future changes to the mean state winds have the potential to lead to large changes in MJO teleconnection amplitude over ~~North America~~ the North Pacific and North America, there does not appear to be a specific change over a particular region that explains inter-model variations.

We find that for perturbations to MJO characteristics (i.e., its intensity, eastward extent, ~~zonal wavenumber~~, or propagation speed), the uncertainty in the MJO teleconnection that results is on the order of the uncertainty from the mean state changes alone. In particular, increases in the MJO's heating intensity ~~and eastward extent~~, ~~eastward extent~~, and ~~zonal wavenumber~~ have the potential to lead to large increases in the MJO teleconnection, with the sensitivity to ~~extent-propagation extent and zonal wavenumber~~ being especially strong for ~~teleconnection over North America for some mean states~~ the teleconnection for some model climates.

Overall, over the North Pacific, despite a relatively large range ~~in total teleconnection amplitude change~~ of ~~-21 to 0.6~~ ~~16 to 0.3~~ % K<sup>-1</sup>, which depends on the CMIP6 model used ~~as the mean state~~ in the LBM simulations, in general it appears as though

the MJO teleconnection to the North Pacific ~~will likely may~~ weaken with warming. Over North America, ~~despite a modest the~~ picture is more blurry, with a large range of potential changes (between  $-11$  to  $15 \text{ \% K}^{-1}$ ) with many models also hovering near, or just above zero.

550 In a multi-model ~~mean-weakening result from the influence of the mean state alone, including MJO changes results in many~~ study assessing internally-simulated MJO teleconnections in both CMIP5 and CMIP6 ~~models showing a range of possible~~ teleconnection changes that is either close to or near zero, Zhou et al. (2020) find no relationship between changes to the MJO's circulation strength and teleconnection amplitude change over North America. The MJO circulation strength is primarily a function of the MJO's heating intensity and the tropical dry static stability (Wolding et al., 2016). Thus, the lack of a relationship between modeled MJO circulation change and teleconnection amplitude change may be due to overwhelming  
555 and independent model spread in other features of the climate system, namely the mean state wind, or propagation features of the MJO, such as its eastward extent, propagation speed, or zonal wavenumber. Our results support the hypothesis suggested in Samarasinghe et al. (2021), that the lack of an apparent relationship between changes to MJO circulation strength and MJO teleconnection amplitude found in previous work appears to be due, in part, to large inter-model spread in future projections of the mean state winds.

560 Over the North Pacific in CESM2-WACCM, Samarasinghe et al. (2021) find large increases in the internally generated MJO teleconnection strength between the pre-industrial period and the end of the century. They hypothesize that this strengthening, which appears despite the weakening expected from increases in tropical dry static stability, may be due to changes in the mean winds that enhance wave propagation over the region. Here, we find that mean state changes alone in CESM2-WACCM lead to a modest *weakening* over the region (Fig. 8a). Thus, it is possible that the strengthening MJO teleconnection over the North  
565 Pacific simulated directly in CESM2-WACCM is due to changes in the MJO. In addition, inter-model spread in features of the MJO in CMIP6 may also be contributing to the results found in Zhou et al. (2020), which we leave for future work.

We have thus identified sources of uncertainty leading to uncertainty in projections of MJO teleconnections in a future warmer climate. To first order, model spread in projections of future winds is the biggest source of uncertainty in projecting the response of MJO teleconnections to warming. ~~For teleconnections to North America in particular, we have identified a~~  
570 ~~region over the eastern Pacific where changes to the winds appear to explain much of the inter-model spread. We speculate that Hence,~~ increased confidence in changes to the ~~mean flow over the Gulf of Alaska would extratropical mean flow may~~ help reduce uncertainty in teleconnection changes over ~~North America. Uncertainty the North Pacific and North America.~~ Additionally, uncertainty in the mean ~~state also affects climate also contributes to spread in~~ how MJO teleconnections respond to ~~increases in the eastward extent of MJO convection some changes to the MJO.~~ Lastly, uncertainty in how the MJO ~~itself~~  
575 will respond to climate change is nontrivial ~~leading to a range of about  $5\% \text{ K}^{-1}$  for any given mean state (see Fig. 8) but~~ interestingly ~~secondary to inter-model uncertainty in the wave ducting properties of the basic flow,~~ which is complicated further by the fact that many models still struggle to produce realistic MJOs (e.g., Ahn et al., 2020).

Over North America, a possible future outcome is that of the median ~~result found in Fig. 8 of a near-zero~~ ~~area-mean total~~ regional mean teleconnection change. In this case, ~~the it is possible that while the mean change over the region as a whole~~  
580 ~~is zero, that the~~ pattern of the teleconnection may change, ~~despite zero average change over the region as a whole leading to~~

increases in the teleconnection amplitude over some parts of North America and decreases elsewhere. In this case, a reduction of the uncertainty in future projections of the mean state winds will ~~still~~ also help reduce uncertainty in the MJO teleconnection change, due to the tight dependence of the stationary wave pattern on the mean winds.

~~We readily admit~~ A discussion of the caveats and limitations of our study is necessary. Firstly, we readily admit the limitations of our choice of a linear model to explore how the MJO teleconnection responds to changes in the mean state and to the MJO ~~itself~~. While this allows us to easily diagnose causal relationships, ~~and~~ span a previously sparsely sampled causative parameter space, ~~the~~ and separately diagnose the effect of changes to mean state winds versus the dry static stability on MJO teleconnections using multiple model mean states; the linear framework is missing nonlinear relationships between the mean state ~~and~~, the MJO forcing, and MJO-excited Rossby waves. For example, the mean state exerts a strong control on MJO propagation characteristics (Jiang et al., 2020). ~~The MJO may also play an active role in shaping the midlatitude mean state winds, for example through the modulation of the local Hadley cell (Lyu et al., 2019). Such feedbacks, which we have not included here. Rossby waves excited by the MJO are also able to extract energy from the mean flow (Adames and Wallace, 2014; Zheng and Chang, 2020), and nonlinear interactions can cause a spatial shift in extratropical maximum geopotential height anomalies associated with the MJO (Lin and Brunet, 2018). Both eddy-mean flow interactions, and any MJO dependence on the mean state~~ are not simulated in the LBM. ~~Nonetheless,~~ While previous work has ~~also~~ shown that the MJO teleconnection is, to first order, linear ~~(Mori and Watanabe, 2008).~~ (Mori and Watanabe, 2008; Lin and Brunet, 2018), future work involving a nonlinear framework may be needed to reduce uncertainty in future projections of the MJO teleconnection to the extratropics.

Secondly, our results are sensitive to the ideal forcing used to explore how a changing MJO may affect teleconnection changes. We have used a very idealized heating to represent the MJO which is constant across LBM simulations. Thus, despite using a range of perturbations of this heating to sample the range of possible changes to the MJO and subsequent impacts on teleconnections, we are lacking a comprehensive sampling of the inter-model variations in future MJO characteristics. Perturbations rely on projected changes from CMIP5, rather than CMIP6 (the latter of which has an improved MJO (Ahn et al., 2020)) due to the availability of detailed published analyses of modeled changes to the MJO. We also examine only a subset of the possible changes to the MJO that may occur; namely, its heating intensity, zonal wavenumber, propagation speed, and eastward propagation extent. For example, we have not investigated how changes to the MJO's frequency, meridional extent, or other changes to its spatial structure may affect future MJO teleconnections. However, given that many models still struggle to reproduce a reliable MJO in the current climate, our current understanding of how the MJO may change with warming is incomplete. Thus there is much more work that needs to be done to understand how MJO teleconnections may change in the future.


Lastly, we have considered only January mean states for this work. It is possible that results could be different for other months or seasons. For example, changes in the mean wind may be less important during the shoulder months when the East Asian subtropical jet is weaker.

For now, we suggest this work as a compelling first step towards understanding the intriguing model spread in future MJO teleconnection strength in the extratropics, and another reason to elevate the ongoing community quest to reduce uncertainty

of the mean midlatitude circulation's response to climate change. [Work comparing these results with changes to the MJO teleconnection directly simulated by CMIP6 models, while outside the scope of the present work, will additionally help deepen understanding of how MJO teleconnections may change in the future and are promising topics for future work.](#)

620 *Code and data availability.* Data to reproduce figures is published online at <https://zenodo.org/> with doi:10.5281/zenodo.4737438. The linear baroclinic model is available through request from M. Hayashi (michiyah@hawaii.edu). CMIP6 data is publicly available through the Earth System Grid Federation at <https://esgf-node.llnl.gov/search/cmip6/>.





**Figure A1.** As in Figure 1, but for the perturbed MJO cases for (a,e,i,m) the lower bound of the propagation speed experiment; (b,f,j,n) the upper bound of the propagation speed experiment; (c,g,k,o) the upper bound of the increased eastward propagation extent experiment; and (d,h,l,p) the upper bound of the perturbed wavenumber experiment.

## Appendix A


Figure [A1](#) shows the idealized forcing at days 10, 20, 30, and 40 for the cases where the forcing is perturbed.

Figure [A2](#) shows the number of simulations that became unstable (and hence omitted from our analysis; see Sect. 2), and indicates the CMIP6 models for which all simulations were excluded from our analysis for having 10 or more unstable ensemble members for a particular basic state combination.


Figure [A3](#) shows the temporal variance of the ensemble mean Rossby wave source ( $S'$ ) in the LBM for each CMIP6 model historical basic state used in the analysis. The box around the region ( $80^\circ$ - $190^\circ$  E,  $20^\circ$ - $40^\circ$  N) indicates the region used for Fig. 5 of the main text.

Figure [A4](#) shows the difference in the January mean 200 hPa zonal wind between the historical and future periods for individual CMIP6 models, with contours outlining the subtropical jet for reference.


Figure [A5](#) shows the difference in the January mean  $K_s$  at 200 hPa between the historical and future periods for the models included in the analysis. Contours outlining the subtropical jet are included for reference.




**Figure A2.** For each model basic state combination, the fraction of the 30 ensemble members that met the instability criterion described in Sect. 2. Models omitted from our analysis (those for which 10 or more of their ensemble members for any of the basic state combinations became unstable) are indicated by the black outline. Th (Ts) = basic state dry static stability from the historical (future) period; Vh (Vs) = basic state winds from the historical (future) period.



**Figure A3.** The temporal variance of the ensemble mean Rossby wave source ( $S'$ ) with the full mean states given by historical quantities. The box around the region (80°-190° E, 20°-40° N) indicates the region used for Figure 5 of the main text.



**Figure A4.** Shading is the difference in the January mean zonal wind at 200 hPa between the historical (1985-2014) and future periods (2071-2100) for models included in the analysis. Solid black contours show the January mean zonal wind at 200 hPa, starting at  $35 \text{ m s}^{-1}$  and increasing by  $10 \text{ m s}^{-1}$ .



**Figure A5.** Shading is the difference in the January mean  $K_s$  at 200 hPa between the historical (1985-2014) and future periods (2071-2100) for models included in the analysis. Solid black contours show the January mean zonal wind at 200 hPa, starting at  $35 \text{ m s}^{-1}$  and increasing by  $10 \text{ m s}^{-1}$ .

*Author contributions.* AMJ led the design of the experiments, performed the simulations, and prepared the manuscript. DAR and EAB aided  
635 with the experimental design, interpretation of the results, and writing of the manuscript.

*Competing interests.* The authors declare no competing interests

*Acknowledgements.* We thank [David Straus and two anonymous reviewers for helpful comments.](#) We thank Michiya Hayashi for sending the LBM code, and acknowledge Donald Dazlich, Stephanie Henderson, and Kai-Chih Tseng for their help ~~with setting up the model in~~  
640 ~~getting the model set up and running.~~ We also thank Michael Pritchard for help with manuscript editing and Eric Maloney and Tom Beucler for helpful suggestions. This work was supported by the National Science Foundation under ~~awards award~~ [AGS-1826643](#) ~~and AGS-1749261,~~  
[by the National Oceanic and Atmospheric Administration \(NOAA\) under award NA19OAR4590155](#), and by the NOAA Climate and Global Change Postdoctoral Fellowship Program, administered by UCAR's Cooperative Programs for the Advancement of Earth System Science (CPAESS) under award NA18NWS4620043B. [Support for EAB was provided, in part, by NSF grant AGS-1749261.](#)

This research also used computing resources of the Extreme Science and Engineering Discovery Environment (XSEDE), which is sup-  
645 ported by National Science Foundation grand number ACI-1548562 (Towns et al., 2014) and allocation number TG-ATM190002.

## References

- Adames, Á. F. and Wallace, J. M.: Three-Dimensional Structure and Evolution of the MJO and Its Relation to the Mean Flow, *J. Atmos. Sci.*, 71, 2007–2026, <https://doi.org/10.1175/JAS-D-13-0254.1>, 2014.
- Adames, Á. F., Kim, D., Sobel, A. H., Del Genio, A., and Wu, J.: Changes in the structure and propagation of the MJO with increasing CO  
650 2, *J. Adv. Model. Earth Syst.*, 9, 1251–1268, <https://doi.org/10.1002/2017MS000913>, 2017.
- Ahn, M., Kim, D., Kang, D., Lee, J., Sperber, K. R., Gleckler, P. J., Jiang, X., Ham, Y., and Kim, H.: MJO propagation across the maritime continent: Are CMIP6 models better than CMIP5 models?, *Geophys. Res. Lett.*, 47, 2020.
- Ahn, M.-S., Kim, D., Sperber, K. R., Kang, I.-S., Maloney, E., Waliser, D., Hendon, H., and on behalf of WGNE MJO Task Force: MJO simulation in CMIP5 climate models: MJO skill metrics and process-oriented diagnosis, *Clim. Dyn.*, 49, 4023–4045,  
655 <https://doi.org/10.1007/s00382-017-3558-4>, 2017.
- Arnold, N. P., Kuang, Z., and Tziperman, E.: Enhanced MJO-like Variability at High SST, *J. Clim.*, 26, 988–1001, 2013.
- Bladé, I. and Hartmann, D. L.: The Linear and Nonlinear Extratropical Response of the Atmosphere to Tropical Intraseasonal Heating, *J. Atmos. Sci.*, 52, 4448–4471, [https://doi.org/10.1175/1520-0469\(1995\)052<4448:TLANER>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<4448:TLANER>2.0.CO;2), 1995.
- Bui, H. X. and Maloney, E. D.: Changes in Madden-Julian Oscillation Precipitation and Wind Variance Under Global Warming, *Geophys. Res. Lett.*, 2018,  
660
- Bui, H. X. and Maloney, E. D.: Mechanisms for Global Warming Impacts on Madden–Julian Oscillation Precipitation Amplitude, *J. Clim.*, 32, 6961–6975, <https://doi.org/10.1175/JCLI-D-19-0051.1>, 2019a.
- Bui, H. X. and Maloney, E. D.: Transient Response of MJO Precipitation and Circulation to Greenhouse Gas Forcing, *Geophys. Res. Lett.*, 7, 847, <https://doi.org/10.1029/2019GL085328>, 2019b.
- 665 Chang, C.-W. J., Tseng, W.-L., Hsu, H.-H., Keenlyside, N., and Tsuang, B.-J.: The Madden-Julian Oscillation in a warmer world, *Geophys. Res. Lett.*, 42, 6034–6042, 2015.
- Cui, J. and Li, T.: Changes of MJO propagation characteristics under global warming, *Clim. Dyn.*, 53, 5311–5327, 2019.
- Deng, Y. and Jiang, T.: Intraseasonal Modulation of the North Pacific Storm Track by Tropical Convection in Boreal Winter, *J. Clim.*, 24, 1122–1137, 2011.
- 670 Goss, M. and Feldstein, S. B.: Testing the sensitivity of the extratropical response to the location, amplitude, and propagation speed of tropical convection, *J. Atmos. Sci.*, pp. JAS–D–17–0132.1, 2017.
- Guo, Y., Shinoda, T., Lin, J., and Chang, E. K. M.: Variations of Northern Hemisphere Storm Track and Extratropical Cyclone Activity Associated with the Madden–Julian Oscillation, *J. Clim.*, 30, 4799–4818, 2017.
- Henderson, S. A., Maloney, E. D., and Son, S. W.: Madden-Julian oscillation Pacific teleconnections: The impact of the basic state and MJO  
675 representation in general circulation models, *J. Clim.*, 30, 4567–4587, <https://doi.org/10.1175/JCLI-D-16-0789.1>, 2017.
- Hoskins, B. J. and Ambrizzi, T.: Rossby Wave Propagation on a Realistic Longitudinally Varying Flow, *J. Atmos. Sci.*, 50, 1661–1671, [https://doi.org/10.1175/1520-0469\(1993\)050<1661:RWPOAR>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<1661:RWPOAR>2.0.CO;2), 1993.
- Jenney, A. M., Randall, D. A., and Barnes, E. A.: Quantifying Regional Sensitivities to Periodic Events: Application to the MJO, *J. Geophys. Res. D: Atmos.*, 44, 7528, <https://doi.org/10.1029/2018JD029457>, 2019.
- 680 Jiang, X., Maloney, E., and Su, H.: Large-scale controls of propagation of the Madden-Julian Oscillation, *npj Climate and Atmospheric Science*, 3, 1–8, <https://doi.org/10.1038/s41612-020-00134-x>, 2020.



- Karoly, D. J.: Rossby wave propagation in a barotropic atmosphere, *Dyn. Atmos. Oceans*, 7, 111–125, [https://doi.org/10.1016/0377-0265\(83\)90013-1](https://doi.org/10.1016/0377-0265(83)90013-1), 1983.
- 685 Lehner, F., Deser, C., Maher, N., Marotzke, J., Fischer, E. M., Brunner, L., Knutti, R., and Hawkins, E.: Partitioning climate projection uncertainty with multiple large ensembles and CMIP5/6, *Earth system change: climate scenarios*, 11, 491–508, 2020.
- Lin, H. and Brunet, G.: Extratropical Response to the MJO: Nonlinearity and Sensitivity to the Initial State, *J. Atmos. Sci.*, 75, 219–234, 2018.
- Lyu, M., Wu, Z., Shi, X., and Wen, M.: Distinct impacts of the MJO and the NAO on cold wave amplitude in China, *Q.J.R. Meteorol. Soc.*, 145, 1617–1635, <https://doi.org/10.1002/qj.3516>, 2019.
- 690 Maloney, E. D. and Xie, S.-P.: Sensitivity of tropical intraseasonal variability to the pattern of climate warming, *J. Adv. Model. Earth Syst.*, 5, 32–47, <https://doi.org/10.1029/2012MS000171>, 2013.
- Maloney, E. D., Adames, Á. F., and Bui, H. X.: Madden–Julian oscillation changes under anthropogenic warming, *Nat. Clim. Chang.*, 9, 26–33, <https://doi.org/10.1038/s41558-018-0331-6>, 2019.
- Mori, M. and Watanabe, M.: The Growth and Triggering Mechanisms of the PNA: A MJO-PNA Coherence, *J. Meteorol. Soc. Japan*, 86, 213–236, <https://doi.org/10.2151/jmsj.86.213>, 2008.
- 695 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, 2010.
- O’Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., 700 Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century, *Glob. Environ. Change*, 42, 169–180, 2017.
- Robertson, A. W., Kumar, A., Peña, M., and Vitart, F.: Improving and promoting subseasonal to seasonal prediction, *Bull. Am. Meteorol. Soc.*, 96, ES49–ES53, <https://doi.org/10.1175/BAMS-D-14-00139.1>, 2015.
- Rushley, S. S., Kim, D., and Adames, Á. F.: Changes in the MJO under Greenhouse Gas–Induced Warming in CMIP5 Models, *J. Clim.*, 32, 705 803–821, <https://doi.org/10.1175/JCLI-D-18-0437.1>, 2019.
- Samarasinghe, S. M., Connolly, C., Barnes, E. A., Ebert-Uphoff, I., and Sun, L.: Strengthened causal connections between the MJO and the north Atlantic with climate warming, *Geophys. Res. Lett.*, 48, 2021.
- Santer, B. D., Wigley, T. M. L., Mears, C., Wentz, F. J., Klein, S. A., Seidel, D. J., Taylor, K. E., Thorne, P. W., Wehner, M. F., Gleckler, P. J., Boyle, J. S., Collins, W. D., Dixon, K. W., Doutriaux, C., Free, M., Fu, Q., Hansen, J. E., Jones, G. S., Ruedy, R., Karl, T. R., Lanzante, 710 J. R., Meehl, G. A., Ramaswamy, V., Russell, G., and Schmidt, G. A.: Amplification of surface temperature trends and variability in the tropical atmosphere, *Science*, 309, 1551–1556, 2005.
- Sardeshmukh, P. D. and Hoskins, B. J.: The Generation of Global Rotational Flow by Steady Idealized Tropical Divergence, [https://doi.org/10.1175/1520-0469\(1988\)045<1228:TGOGRF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<1228:TGOGRF>2.0.CO;2), 1988.
- Seo, K.-H. and Lee, H.-J.: Mechanisms for a PNA-Like Teleconnection Pattern in Response to the MJO, *J. Atmos. Sci.*, 74, 1767–1781, 715 <https://doi.org/10.1175/JAS-D-16-0343.1>, 2017.
- Seo, K.-H. and Son, S.-W.: The Global Atmospheric Circulation Response to Tropical Diabatic Heating Associated with the Madden–Julian Oscillation during Northern Winter, *J. Atmos. Sci.*, 69, 79–96, <https://doi.org/10.1175/2011JAS3686.1>, 2012.

- Seo, K.-H., Lee, H.-J., and Frierson, D. M. W.: Unraveling the Teleconnection Mechanisms that Induce Wintertime Temperature Anomalies over the Northern Hemisphere Continents in Response to the MJO, *J. Atmos. Sci.*, 73, 3557–3571, <https://doi.org/10.1175/JAS-D-16-0036.1>, 2016.
- Subramanian, A., Jochum, M., Miller, A. J., Neale, R., Seo, H., Waliser, D., and Murtugudde, R.: The MJO and global warming: A study in CCSM4, *Clim. Dyn.*, 42, 2019–2031, <https://doi.org/10.1007/s00382-013-1846-1>, 2014.
- Takahashi, C. and Shirooka, R.: Storm track activity over the North Pacific associated with the Madden-Julian Oscillation under ENSO conditions during boreal winter, *J. Geophys. Res.*, 119, 10,663–10,683, 2014.
- 725 Ting, M. and Held, I. M.: The Stationary Wave Response to a Tropical SST Anomaly in an Idealized GCM, *J. Atmos. Sci.*, 47, 2546–2566, 1990.
- Towns, J., Cockerill, T., Dahan, M., Foster, I., Gaither, K., Grimshaw, A., Hazlewood, V., Lathrop, S., Lifka, D., Peterson, G. D., Roskies, R., Scott, J. R., and Wilkins-Diehr, N.: XSEDE: Accelerating Scientific Discovery, *Computing in Science Engineering*, 16, 62–74, 2014.
- Tseng, K.-C., Maloney, E., and Barnes, E.: The Consistency of MJO Teleconnection Patterns: An Explanation Using Linear Rossby Wave Theory, *J. Clim.*, 32, 531–548, <https://doi.org/10.1175/JCLI-D-18-0211.1>, 2019.
- 730 Tseng, K.-C., Barnes, E. A., and Maloney, E.: The Importance of Past MJO Activity in Determining the Future State of the Midlatitude Circulation, *J. Clim.*, 33, 2131–2147, 2020a.
- Tseng, K.-C., Maloney, E., and Barnes, E. A.: The Consistency of MJO Teleconnection Patterns on Interannual Time Scales, *J. Clim.*, 33, 3471–3486, <https://doi.org/10.1175/JCLI-D-19-0510.1>, 2020b.
- 735 Wang, J., Kim, H., Kim, D., Henderson, S. A., Stan, C., and Maloney, E. D.: MJO Teleconnections over the PNA Region in Climate Models. Part II: Impacts of the MJO and Basic State, *J. Clim.*, 33, 5081–5101, 2020.
- Watanabe, M. and Kimoto, M.: Atmosphere-ocean thermal coupling in the North Atlantic: A positive feedback, *Q.J. Royal Met. Soc.*, 126, 3343–3369, <https://doi.org/10.1002/qj.49712657017>, 2000.
- Wolding, B. O., Maloney, E. D., and Branson, M.: Vertically resolved weak temperature gradient analysis of the Madden-Julian Oscillation in SP-CESM, *J. Adv. Model. Earth Syst.*, 8, 1586–1619, <https://doi.org/10.1002/2016MS000724>, 2016.
- 740 Wolding, B. O., Maloney, E. D., Henderson, S., and Branson, M.: Climate change and the Madden-Julian Oscillation: A vertically resolved weak temperature gradient analysis, *Journal of Advances in Modeling Earth Systems*, 9, 307–331, <https://doi.org/10.1002/2016MS000843>, 2017.
- Yadav, P. and Straus, D. M.: Circulation Response to Fast and Slow MJO Episodes, *Mon. Weather Rev.*, 145, 1577–1596, 2017.
- 745 Zheng, C. and Chang, E. K. M.: The role of MJO propagation, lifetime, and intensity on modulating the temporal evolution of the MJO extratropical response, *J. Geophys. Res.*, 124, 5352–5378, 2019.
- Zheng, C. and Chang, E. K.-M.: The Role of Extratropical Background Flow in Modulating the MJO Extratropical Response, *J. Clim.*, 33, 4513–4536, 2020.
- Zhou, S., L’Heureux, M., Weaver, S., and Kumar, A.: A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Clim. Dyn.*, 38, 1459–1471, 2012.
- 750 Zhou, W., Yang, D., Xie, S.-P., and Ma, J.: Amplified Madden-Julian oscillation impacts in the Pacific-North America region, *Nat. Clim. Chang.*, <https://doi.org/10.1038/s41558-020-0814-0>, 2020.