

# Supercell Convective Environments in Spain based on ERA5: Hail and Non-Hail Differences

Carlos Calvo-Sancho<sup>1</sup>, Javier Díaz-Fernández<sup>2</sup>, Yago Martín<sup>3</sup>, Pedro Bolgiani<sup>2</sup>, Mariano Sastre<sup>2</sup>, Juan Jesús González-Alemán<sup>4</sup>, Daniel Santos-Muñoz<sup>5</sup>, José Ignacio Farrán<sup>1</sup> and María Luisa Martín<sup>1,6</sup>,

- 5 <sup>1</sup>Department of Applied Mathematics, Faculty of Computer Engineering, University of Valladolid, Segovia, Spain.  
<sup>2</sup>Department of Earth Physics and Astrophysics, Faculty of Physics, Complutense University of Madrid, Madrid, Spain.  
<sup>3</sup>Department of Geography, Faculty of History and Philosophy, University Pablo de Olavide, Sevilla, Spain.  
<sup>4</sup>State Meteorological Agency (AEMET), Madrid, Spain.  
<sup>5</sup>Danmarks Meteorologiske Institut, Copenhagen, Denmark.  
10 <sup>6</sup>Institute of Interdisciplinary Mathematics (IMI), Complutense University of Madrid, Madrid, Spain.

*Correspondence to:* Carlos Calvo-Sancho (carlos.calvo.sancho@uva.es)

**Abstract.** Severe convective storms, in particular supercells, are occasionally responsible for a large number of losses of property and damages in Spain. This paper aims to study the synoptic configurations and pre-convective environments in a dataset of 262 supercells during 2011-2020 in Spain. The events are grouped into supercells with hail (diameter larger than 5  
15 cm) and without hail and the results are compared. ERA5 reanalysis data are used to study the synoptic configurations and **proximity atmospheric profiles** related to the supercell events at the initial time. Moreover, temperature, convective available potential energy, convective inhibition, lifting condensation level, level of free convection, height of freezing level, wind shear and storm-relative helicity are determined for each event. The results show that supercells are more frequent in the Mediterranean coast during the warm season. **Some of the variables analysed present statistically significant differences**  
20 **between hail and non-hail events, particularly, supercells with hail are characterized** by higher median values of most-unstable convective available potential energy than supercells without hail.

## 1 Introduction

Convective storms and their associated phenomena (lightning, hail, wind or flash-floods) have a great influence on human activities due to the destructive consequences they may have (Martín et al., 2020; Taszarek et al., 2020a; Rodriguez and Bech,  
25 2021). Europe is regularly threatened by severe convective storms (Dahl, 2006), causing considerable economic loss, social impact, and endangering aviation safety (Nisi et al., 2016; Mohr et al., 2017; Antonescu et al., 2017; Kunz et al., 2020; Chernokulsky et al., 2020; Gatzert et al., 2020). Thus, improving the knowledge on the genesis and **lifecycle** of convective storms is a constant endeavour in the meteorological community.

Thunderstorm cells can be formed either in a discrete and isolated form, or in large and organized systems, e.g., squall-lines.  
30 Based on their structure, organization, and size, three different thunderstorm types are defined by the US National Weather Service (NWS, 2019): ordinaries, multicell and supercells. Concerning supercells, Browning (1962) defines them as **convective**

storms, occurring in a significantly vertically-sheared environment, that contains a deep and persistent mesocyclone, representing the most organized, severe and long-lasting form of isolated deep convection phenomena. These systems are linked to hail reports -including hail diameters larger than 5 cm- and EF2 tornadoes or higher (Duda and Gallus, 2010; Quirantes et al., 2014). **Supercells are common phenomena in spring and summer (Brooks et al., 2019), and can be detected through ground-based or satellite lightning detection systems (Bedka et al., 2018; Galanaki et al., 2018).** However, the local phenomena associated with these systems, such as hail, require observational reports to be confirmed. As many of these events occur in unpopulated areas, the observed weather reports have a spatial bias toward the most populated areas (Groenemeijer et al., 2017; Edwards et al., 2018). In recent years, thunderstorm reports have increased due to the accessibility of the general population to new technologies, especially thanks to smart-phones and social networks. This has allowed an improvement in databases related to convective storms, with increasing availability of the information on these phenomena (Elmore et al., 2014; Krennert et al., 2018; Taszarek et al., 2020b). **Nevertheless, a rigorous quality and validity control, through validated observational data (radar, satellite...), should be applied to these severe weather reports to be scientifically valid, e.g., European Severe Weather Database (Dotzek et al., 2009).**

**Mainly due to orography, smaller Convective Available Potential Energy (CAPE) and wind shear (WS), supercells in Europe tend to be less severe and with return periods (hail  $\geq$  8cm, violent tornadoes) longer than those formed in the US, and therefore show reduced rotation and shorter life spans (Quirantes et al., 2014; Taszarek et al., 2020b).** In the particular case of Spain, the study of severe convective storms has been increasing, extending the knowledge about these systems in recent years. Martin et al. (2020) found more than a hundred supercells per year on average in Spain. Weather environments conducive to severe convective storms have been identified in different studies suggesting that synoptic environments (Merino et al., 2013; Mora et al., 2015), mesoscale characteristics (García-Ortega et al., 2012), orography (Romero et al., 1998) and **convective variables** (Calvo-Sancho and Martin, 2021) should be considered together in the research of supercells. Castro et al. (1992) explored the role of topography in the formation and evolution of convective storms in the Ebro Valley (Figure 1a), concluding that mountainous terrain affects the supercells trajectories and velocities. Regarding hailstorms, Merino et al. (2019) highlight that the main triggers of convection are thermal instability and low-level convergence. Tornadoes occurrence and intensity are not as severe as in other regions of the world (e.g., US) **mainly** due to the absence of wet fluxes inland (Rodríguez and Bech, 2018), **however, the large occurrence of these events in northeastern Spain, causes substantial damages for the local economy, especially in crop fields (López and Sanchez, 2009).** Moreover, Gayà (2011) performed a climatology of tornadoes and waterspouts in Spain and Rodríguez and Bech (2018, 2020) surveyed the mesoscale environments wherein tornadoes and waterspouts formed in the Iberian Peninsula. Both studies reveal that WS plays a more important role than CAPE in synoptic and mesoscale environments in the cold season.

Since severe supercells have caused substantial property damage and economic losses in recent years in Spain, with around 300 casualties due to severe convective storms from 1987 to 2020 (Consortio de Compensación de Seguros, 2020), this study aims to provide a better understanding of the supercell synoptic and mesoscale environments. To do this, a set of events are selected from the supercell database of Martin et al. (2020) which are then categorized into two distinct groups, i.e., with hail

diameter larger than 5 cm (SP-HAIL) and without hail (SP-NONHAIL). Manzato (2012) recorded hailstorms using hailpads to perform a hail climatology in northeast Italy. Merino et al. (2013) using hailpads data too to study the synoptic and mesoscale configurations for hailstorms in southwestern Europe.

70 Herein, these systems are analysed through their synoptic and mesoscale environments using ERA5 reanalysis (Hersbach et al., 2020). ERA5 is a state of the art, high-resolution reanalysis, which has shown successful results in studies related to severe local storm environments over North America (Coffer et al., 2020; Li et al., 2020; Taszarek et al., 2020a, 2020b), severe convective storms across Europe (Taszarek et al., 2020a, 2020b; Calvo-Sancho and Martín 2021), tornadic environments in the Iberian Peninsula (Rodríguez and Bech, 2020) and microbursts (Bolgiani et al., 2020). On the other hand, severe storms have been forecasted using numerical weather prediction models and alternative methods based on them, as it can be seen in  
75 Gascon et al. (2015) survey. Their study selected the Showalter Index, dew point temperature at 850 hPa, storm relative helicity between 0-3 km, wind speed at 500 hPa and wet bulb zero to develop a logistic equation that gave the severe storm develop probability, obtaining a strong capacity to forecast severe storms near the Ebro Valley.

This work is organized as follows. The database and methodology are described in Section 2. Section 3 shows the discussion of the main results related to synoptic and convective variables associated with SP-HAIL and SP-NONHAIL events. Finally,  
80 the main conclusions are summarized in Section 4.

## 2 Data and methodology

### 2.1 Datasets

The supercell sample used is selected from the Spanish Supercell Database (Martín et al., 2020) for the 2011-2020 period.  
85 This dataset is formed by confirmed (i.e., doppler radar images, hail greater than 5 cm reports, tornadoes greater than EF2 or images of the event) and medium-high confidence (detected in non-doppler radar images but without direct observation; see Figure 4 in Martín et al. 2020) supercell events through reports from volunteers and collaborators. In total, this dataset is formed by 1758 supercells, from which 262 of them correspond to confirmed supercells and 1495 to medium-high confidence supercells. It's worth noting that even if the database covers all the Spanish territories, there are no events reported in the  
90 Canary Islands. The database defines the supercell spatial lifecycle through an ellipse in a Geographical Information System. Furthermore, the Spanish Supercell Database collects additional information associated with the events, e.g., hail diameter, tornado intensity. In the current study, only the confirmed supercells are selected. These are then categorised as SP-HAIL and SP-NONHAIL according to the observation or not of hail with diameters larger than 5 cm. It should be noted that in this study the Spanish Supercell Database was cross-matched with the European Severe Weather Database (ESWD) and Notification System for Singular Atmospheric Observations (SINOBAS). The validation results show more than 80% of the SP-HAIL  
95 events from the Spanish Supercell Database are included in the ESWD and SINOBAS datasets. Finally, the initial formation time ( $t_0$ ) of each supercell is selected to characterize the development phase of the cell, since once convection is triggered, the environment might be "contaminated" and the variables may not be as representative of the supercell conditions.

The ERA5 reanalysis (Hersbach et al., 2020) is selected to study the synoptic characteristics and the convective variables involved in the supercell development. This is the 5th generation reanalysis created by the European Centre for Medium-Range Weather Forecasts (ECMWF). It is provided with a horizontal grid resolution of  $0.25^\circ \times 0.25^\circ$ , 1 hour as temporal resolution and 137 hybrid model levels for the vertical resolution, from 1000 hPa to 1 hPa, approximately. In the current work, the domain is delimited to  $60^\circ\text{N}/20^\circ\text{N} \times 30^\circ\text{W}/30^\circ\text{E}$  to study the environments related to the supercells in Spain. To analyse the mesoscale setting, a vertical profile of temperature, dew point, geopotential height, pressure, and wind components (u, v) are derived from the ERA5 grid for each supercell event.




Figure 1: Domain and orography of the study area (m). White star is referred to Maestrazgo area.

## 2.2 Compositing methodology

Following the methodology of Calvo-Sancho and Martin (2021) and Gensini et al. (2021), supercell soundings for SP-HAIL and SP-NONHAIL events are built for  $t_0$ . Each vertical profile is computed from ERA5 using the nearest grid point to the supercell location at  $t_0$ . A quality control is carried out to remove any sounding related to convective boundary propagation (Brooks et al. 2003, 2007; Gensini et al. 2021). Accordingly, each vertical profile must record a non-zero Most-Unstable Convective Available Potential Energy (MUCAPE) and Mixed-Layer Level of Free Convection (MLLFC) to be included in the study. Once the vertical profiles are obtained, composites for SP-HAIL and SP-NONHAIL are derived at  $t_0$ .

Synoptic patterns composites of both events are created to describe and compare the common large-scale features. The ERA5 atmospheric fields used to compute the composites are: 500 and 300 hPa geopotential height, mean sea level pressure, dew point, wind direction and wind speed at 10 meter above sea level, 700-400 hPa integrated mean of omega vertical velocity,

and 0-6 km WS. These atmospheric variables have been used in studies related to spatial patterns of hailstorms (Merino et al., 2013; Melcón et al., 2017), supercells (Gropp and Davenport, 2018) and thunderstorms (Mora et al., 2015).

## 120 2.3 Convective variables methodology

To characterize the convective environment, several thermodynamic and kinematic variables are calculated using the thunderR language package (Taszarek et al., 2021) for each vertical profile. The selection of these parameters (Table 1) is based on similar studies related to severe convective storms in US and Europe (Rasmunssen et al., 1998; Kaltenböck et al., 2009; Westermayer et al., 2017; Rodríguez and Bech, 2018, 2020; Taszarek et al., 2020; Davenport, 2021). The 2-meter temperature (T2M) and dew-point (DWPT) are selected. CAPE and Convective Inhibition (CIN) using Most-Unstable (MU), Mixed-Layer (ML; averaged over 0-500 m above ground level) and Surface-Based (SB) parcels are calculated using the virtual temperature correction (Doswell and Rasmussen, 1994). The deep-layer bulk wind shear over 0-6 km (WS06) and the effective bulk wind difference [EBWD; limited to the layer in which  $CAPE \geq 100 \text{ J kg}^{-1}$  and  $CIN \geq -250 \text{ J kg}^{-1}$ ; Thompson et al., 2007] are calculated. Finally, other parameters relevant to SP-HAIL are also included: ML lifting condensation level (MLLCL), ML level of free convection (MLLFC), height of freezing level (FZH) and height of wet-bulb freezing level (FZH\_W).

**Table 1. Description of the used convective variables.**

Parameter	Abbreviation	Units
<b>Thermodynamic parameters</b>		
2-meter temperature	T2M	°C
2-meter dew-point temperature	DWPT	°C
<b>Parcel parameters</b>		
Most-unstable convective available potential energy	MUCAPE	$\text{J Kg}^{-1}$
Surface-based convective available potential energy	SBCAPE	$\text{J Kg}^{-1}$
Mixed-layer convective available potential energy	MLCAPE	$\text{J Kg}^{-1}$
Most-unstable convective inhibition	MUCIN	$\text{J Kg}^{-1}$
Surface-based convective inhibition	SBCIN	$\text{J Kg}^{-1}$
Mixed-layer convective inhibition	MLCIN	$\text{J Kg}^{-1}$
Mixed-layer lifting condensation level	MLLCL	m
Mixed-layer level of free convection	MLLFC	m
Height of freezing level	FZH	m
Height of wet-bulb freezing level	FZH_W	m
<b>Kinematic parameters</b>		
Deep-layer bulk wind shear over 0-6 km	WS06	$\text{m s}^{-1}$
Effective bulk wind difference	EBWD	$\text{m s}^{-1}$

Storm-relative helicity over 0-1 km	SRH01	$\text{m}^2 \text{s}^{-2}$
Storm-relative helicity over 0-3 km	SRH03	$\text{m}^2 \text{s}^{-2}$

135 The application of the non-parametric Mann-Whitney test (Mann and Whitney, 1947) is used to establish statistical differences  
(at  $p < 0.05$ ) between the SP-HAIL and SP-NONHAIL groups for the above-mentioned parameters at  $t_0$  (Table 2). The results  
show differences for both thermodynamic variables, all the CAPE variables and the freezing level related variables. Also,  
MLCIN and EBWD are statistically different.

140 **Table 2. p-values of the Mann-Whitney test for all the variables analysed for SP-HAIL and SP-NONHAIL events at  $t_0$ . p-values  
equal or lower than 0.05 are in bold.**

SP-HAIL SP-NONHAIL		SP-HAIL SP-NONHAIL	
MUCAPE	<b>0.00</b>	MLLFC	0.43
SBCAPE	<b>0.00</b>	FZH	<b>0.00</b>
MLCAPE	<b>0.00</b>	FZH_W	<b>0.00</b>
T2M	<b>0.02</b>	WS06	0.23
DWPT	<b>0.00</b>	WS01	0.17
SBCIN	0.24	EBWD	<b>0.00</b>
MLCIN	<b>0.01</b>	SRH01	0.44
MUCIN	0.48	SRH03	0.62
MLLCL	0.16		


### 3 Results and discussion

The spatial and temporal distribution of supercells for both SP-HAIL and SP-NONHAIL formed in the Spanish mainland are  
first assessed. The main results relative to large-scale composites, and the thermodynamic and kinematic variables involved in  
145 supercell formation in the domain are presented and discussed in the following two subsections.

The spatial distribution of the reported supercell episodes (Figure 2a) shows that most of the events for both SP-HAIL and SP-  
NONHAIL took place in the eastern half of Spain. The Ebro Valley and the Mediterranean coastal area accumulate 79.9% of  
the SP-NONHAIL and 88.3% of SP-HAIL. This is consistent with lightning observations in Spain, as the eastern Iberian  
System area (white star in Figure 1a) has the highest density of lightning flash per year (Mora et al. 2019). This area favours  
150 convective initiation and supercell formation due to low-level convergence (northwesterly-southeasterly and south westerly-  
easterly winds), upper-level forcing for ascent, low-medium level moist coming from the Mediterranean Sea and strong diurnal  
heating (Mora et al. 2015).

155 The temporal distribution of supercell events (Figure 2b) matches with the warmest and stronger insolation months (July and August accumulate 53.3% of the SP-NONHAIL and 74.4% of the SP-HAIL storms) in the study area, since deep convection is a necessary condition to the formation of supercells (Markowski and Richardson, 2010; Miglietta et al., 2017; Taszarek et al., 2019). This is consistent with other studies on convective storms in Europe that assess the higher thunderstorm frequency in summertime, when the diurnal heating is stronger (Merino et al., 2013; Kotroni and Lagouvardos, 2016; Taszarek et al., 2018; Taszarek et al., 2019). The hourly distribution of the supercells (Figure 2c) shows a concentration of the events during the late afternoon (summer local time is Universal Time Coordinated plus 2 hours, UTC+2), shortly after the daily insolation maximum in the study area. However, the results also yield a large persistence of the conditions, as many events are reported well into the evening.

160




**Figure 2: (a) Location of the dataset events (SP-HAIL and SP-NONHAIL) from 2011 to 2020 in Spain. (b) Monthly supercell distribution. (c) Hourly supercell distribution ( $t_0$ ; UTC).**

Synoptic pattern composites for the most relevant atmospheric variables in SP-HAIL and SP-NONHAIL events are shown in this subsection to describe and compare the large-scale characteristics.

Non-substantial differences between SP-HAIL and SP-NONHAIL are found in the mean sea level pressure (Figure 3). However, the 500 hPa geopotential height displays a SP-NONHAIL composite with a deeper trough and weaker geopotential height gradient, in comparison with SP-HAIL. A similar situation is shown by the 300 hPa geopotential height. This atmospheric configuration promotes weak WS in upper-levels, which could be indicative of weaker convective environments (Weisman and Klemp, 1982; Brooks et al., 2003; Taszarek et al., 2017). Although there are differences between SP-HAIL and SP-NONHAIL, both geopotential configurations promote upper-level positive vorticity advection (not shown) and divergence over Spain, which favour a stronger upper-level forcing (Markowski and Richardson, 2010). Values of  $1.1 \text{ Pa}^{-1} \text{ s}^{-3}$  in SP-HAIL events at 700-400 hPa thickness of Q-vector divergence (Figure S1b, Supplementary) and statistical differences of Q-vector divergence between SP-HAIL and SP-NONHAIL events (Figure S1c, Supplementary) are found over eastern Spain. The positive Q-vector divergence values indicate forcing for ascent where supercells could have been originated by strong convection. Thermal lows (1012 hPa) can also be appreciated in the center of Spain (Figure 3b). These lows are typical of the summer months (Tullot, 2000), promoting east wind flows and ensuring humidity from the Mediterranean Sea in the supercell formation area favouring the initiation of deep convection. Thus, a more favourable environment for deep-moist convection should be expected for SP-HAIL, as the corresponding composite shows a deeper thermal low, covering a larger area and accompanied by an enhanced easterly flow.


Mora et al. (2015) studied electrically severe convective storms in the northern plateau of Spain during 2000-2010, finding that 31% of these thunderstorm episodes were linked to upper-level troughs. These episodes were characterized by strong baroclinic short waves and deep troughs at 500 hPa, which is a pattern very similar to the one shown in SP-HAIL and SP-NONHAIL composites in Figure 3, respectively. Therefore, the results are in line with Mora et al. (2015), showing that supercell episodes in Spain are associated with troughs at upper and medium levels of the troposphere. Overall, the higher convective activity is located on the eastern of Spain, corresponding to the right side of the troughs, with the thermal lows at the centre of Spain.





**Figure 3: 500 hPa geopotential height (coloured; dam), 300 hPa geopotential height (blue contours; dam) and mean sea level pressure (black lines; hPa) composites at  $t_0$  for (a) SP-NONHAIL, (b) SP-HAIL and (c) Differences between SP-HAIL and SP-NONHAIL.**

One of the main features favouring the deep-moist convection is the moisture at lower and medium levels (Taszarek et al., 2019). Figure 4c depicts statistically significant differences in the DWPT values between SP-HAIL and SP-NONHAIL, being these differences clearly higher in the Spanish Mediterranean coast. Over the land, a notable difference in DWPT is seen for SP-HAIL over the Ebro Valley (Figure 4b), along with a stronger easterly wind flow. This would be a result from the geopotential and thermal low configuration described above, which induces humid air advection from the Mediterranean Sea. According to the DWPT climatology (not shown), the DWPT in the Ebro Valley and the Mediterranean coast is higher in August (when the SP-HAIL are predominant; Figure 2b) than in July. The convective processes are then supported by the favourable environment that promotes deep convection in those zones and pushed by the south-westerly flows. This process is consistent with the results of the supercell observations for the period 2011-2020 (Figure 2a).




**Figure 4: 2-m dew point temperature (contours; °C) and 10 meters wind (arrows;  $m s^{-1}$ ) composites at  $t_0$  for (a) SP-NONHAIL, (b) SP-HAIL and (c) Differences between SP-HAIL and SP-NONHAIL. Black points in plot c) denote statistically significant differences ( $p$ -value  $< 0.05$ ) in dew point temperature.**

The omega vertical velocity composites show statistically significant differences between SP-HAIL and SP-NONHAIL at  $t_0$  (Figure 5). The omega maxima (referred to absolute values) for both supercell groups throughout the lifecycle of the systems are located in the Ebro valley axis and the Iberian System Mountains, where supercells are most common (Figure 5b). The maxima omega vertical velocity at 700-400 hPa thickness for SP-NONHAIL is larger than for SP-HAIL, excepting in the Ebro valley. These higher SP-HAIL omega vertical velocity along with the low-level wind convergence favour the convection initiation (Markowski and Richardson, 2010). Therefore, sustained omega vertical velocities (Figure 5b) and winds convergence (Figure 4b) trigger, enhance and reinforce deep-moist convection at  $t_0$ , favouring large hail formation over the Ebro Valley and the Mediterranean area (Gutierrez and Kumjian, 2021).

Vertical WS promotes storm organization and its longevity. However, excessive WS can be unfavourable to weak updrafts in environments of low instability and, furthermore, can be disadvantageous to convection initiation by increasing entrainment (Markowski and Richardson, 2010). Figure 5 shows similar strong WS06 values ( $> 20 m s^{-1}$ ) for both SP-HAIL and SP-NONHAIL events. The conjunction of upper-level forcing (Figure 3), low-level convergence (Figure 4) and strong omega vertical velocity (Figure 5) promotes organization, longevity and severity in the convective storm systems. Additionally, as

220 depicted in Figure S1, the maxima Q-vector divergence area ( $1.1 \text{ Pa}^{-1} \text{ s}^{-3}$ ; Figure S1b) matches with the convergence of Q-vectors. Figure S1 shows how the upper-level forcing for ascent are higher in SP-HAIL than SP-NONHAIL events in the Mediterranean coast.



225 **Figure 5: 700-400 hPa omega vertical velocity (contours;  $\text{Pa s}^{-1}$ ), 0-6 km WS (green lines;  $\text{m s}^{-1}$ ) composites at  $t_0$  for (a) SP-NONHAIL, (b) SP-HAIL and (c) Differences between SP-HAIL and SP-NONHAIL. Black points in plot c) denote statistically significant differences ( $p\text{-value} < 0.05$ ) in omega vertical velocity.**

### 3.2 Convective variables


As described in the convective variables methodology (Section 2.3), results of the T2M, DWPT, CAPE, CIN, MLLCL, MLLFC, FZH and WS variables from the ERA5 database are presented in this subsection. These results are shown as violin plots, where the probability density distributions of each variable can be seen, as well as the differences between SP-HAIL and SP-NONHAIL events at  $t_0$ .

230 Based on the synoptic compositing methodology, schematic SP-HAIL and SP-NONHAIL composite soundings at  $t_0$  are determined (Figure 6). In order to show the vertical profile of the supercells developing in highly unstable environments, the 90th percentile (based on MUCAPE values) of the Spanish Supercell Database is selected. The 90th percentile vertical profile for each supercell classification reveals interesting features, particularly on the surface, low-levels, and the convective energy.

235 The composite sounding for SP-HAIL (Figure 6b) displays a larger CAPE area than for SP-NONHAIL (Figure 6a). This high value of CAPE ( $1485.9 \text{ J kg}^{-1}$ ) is strongly associated with vertical accelerations (Markowski and Richardson, 2010), so hail formation would be favoured. Related to the CIN, it is higher for SP-HAIL ( $-94.2 \text{ J kg}^{-1}$ ) than for SP-NONHAIL ( $-56.2 \text{ J kg}^{-1}$ ). Differences between LCL (Figure 6, black dot in panels) and LFC are shown, being higher for SP-HAIL than for SP-NONHAIL events. According to Mulholland et al. (2021), a higher LCL is related to the deep convective updraft width. This

240 is resulting on a wider and deeper column and a faster vertical velocity due to the larger distance and residence time of the dry thermal to entrain. Thus, the ensuing moist updraft above the LCL is wider, less dilute and has a greater vertical velocity, which would be in line with the omega vertical velocity results (Figure 5b). Wind barbs reveal a moderate WS06 for both types of supercells. According to Markowski and Richardson (2010) WS tends to enhance the organization, severity, and longevity of the deep moist convection. This due to the degree to which precipitation and outflow affect with an updraft is


245 reduced as the WS over the updraft depth increases. In addition, SP-HAIL low-level WS is higher than for SP-NONHAIL, favouring hail growth (Gutierrez and Kumjian, 2021).



**Figure 6: 90th percentile (based on MUCAPE values) soundings composites at  $t_0$  for (a) SP-NONHAIL and (b) SP-HAIL. Black dot indicates the LCL value.**

250 The distribution for T2M and DWPT show differences between both types of events, which are statistically significant (Table 2) for both variables. Different distributions can be seen in Figure 7. The T2M for SP-HAIL distribution depicts a lower variability and larger median value (Table 3) than the corresponding SP-NONHAIL. The T2M maximum (minimum) for SP-HAIL is 33.0 °C (16.8 °C), showing both groups a very similar maximum value, while the minimum is significantly lower (7.9 °C) for SP-NONHAIL. The DWPT median value for SP-HAIL is greater than for SP-NONHAIL (Table 3). These differences are mainly originated in the wind flows, since in the Spanish Mediterranean area, Balearic Islands and places favourable for maritime fluxes, the main contributor to low-level moisture is advection from the warm Mediterranean Sea. However, in the Spanish inland the main contributor would be the evapotranspiration of the crop fields and vegetation (Vicente-Serrano et al. 2014; Tomas-Burguera et al. 2021), contributing considerably less humidity to the environment.

255




260 **Figure 7: T2M and DWPT distributions and boxplots for SP-HAIL and SP-NONHAIL at  $t_0$ . Median values are represented by white points.**

CAPE is a common and useful forecast tool, with the combination of vertical WS, for predicting supercells and hail (Kaltenböck et al., 2009; Merino et al., 2013). Its distributions in the parcel measurements of MUCAPE, SBCAPE and MLCAPE are shown in Figure 8a. The results show statistically significant differences between the distributions for SP-HAIL and SP-NONHAIL (Table 2). It is noteworthy that CAPE distributions follow positive skew distributions for SP-NONHAIL events, being the SP-HAIL median values notably larger than SP-NONHAIL. However, the 25th percentile and median values for SP-HAIL SBCAPE, 758 and 1231 J kg<sup>-1</sup>, respectively, are greater than those described by Kaltenböck et al. (2009) for Europe, approximately 400 and 1000 J kg<sup>-1</sup>. According to Weisman and Klemp (1982) and Markowski and Dotzek (2011), CAPE is dependent on humidity and orography, with slightly larger values in high elevations than in low terrains because potential temperature increases evenly with height. Therefore, the differences between the current study and Kaltenböck et al. (2019) lie in the high elevations and relatively low humidity in the research area. The MLCAPE median value is close to the SBCAPE value and both also yield larger results for SP-HAIL than for SP-NONHAIL events. Kahraman et al. (2017) analysed the convective storm environments for tornado and severe hail days from 1979 to 2013 in Turkey. In their study, severe storm environments are characterized by smaller CAPE in Turkey compared to US, highlighting that severe hail occurrence is associated with large CAPE and vertical wind shear. In the current analysis, the median values for MUCAPE, SBCAPE and MLCAPE (Table 3) in SP-HAIL events are a bit greater than those obtained by Kahraman et al. (2017) and Púčik et al. (2015) in their study of severe hail-thunderstorms in central Europe. This discrepancy might be partially attributed to the warmer eastern surrounding seas (Mediterranean and Black Sea; Shaltout and Omstedt, 2014) of the aforementioned studies. In the Taszarek et al. (2020b) study of severe convective storms with large hail, Europe MLCAPE median values are similar than for the current study, with values around 1000 J kg<sup>-1</sup> for both. However, the US MLCAPE median result is 1200 J kg<sup>-1</sup>. As it is discussed above, supercells in Europe tend to be smaller than the ones formed in the US, with lower rotation values and shorter

lifecycles (Quirantes et al., 2014; Taszarek et al., 2020b). According to Rodriguez and Bech (2018), the CAPE values found in our study would correspond with those for tornadic storms in eastern Spain and the Balearic Islands. These authors analyse a dataset of 907 tornadoes and waterspout events from 1980 to 2018 using atmospheric profiles from the ERA5 reanalysis and finding SBCAPE values higher than 700 J kg<sup>-1</sup> in tornadic storms (EF1 or stronger).

**Table 3. Median values for each parameter analysed for SP-HAIL and SP-NONHAIL events at t<sub>c</sub>.**

	<b>SP-HAIL t<sub>0</sub></b>	<b>SP-NONHAIL t<sub>0</sub></b>
<b>MUCAPE</b> (J Kg <sup>-1</sup> )	1231.0	691.7
<b>SBCAPE</b> (J Kg <sup>-1</sup> )	1231.0	662.1
<b>MLCAPE</b> (J Kg <sup>-1</sup> )	966.1	444.6
<b>T2M</b> (°C)	23.0	21.7
<b>DWPT</b> (°C)	14.7	11.9
<b>SBCIN</b> (J Kg <sup>-1</sup> )	-5.3	-3.7
<b>MLCIN</b> (J Kg <sup>-1</sup> )	-20.2	-12.2
<b>MUCIN</b> (J Kg <sup>-1</sup> )	-3.5	-3.6
<b>MLLCL</b> (m)	1075.0	1265.0
<b>MLLFC</b> (m)	1870.0	1815.0
<b>FZH</b> (m)	3227.5	2855.0
<b>FZH_W</b> (m)	2832.5	2590.0
<b>WS06</b> (m s <sup>-1</sup> )	19.6	18.1
<b>EBWD</b> (m s <sup>-1</sup> )	18.7	14.6
<b>SRH01</b> (m <sup>2</sup> s <sup>-2</sup> )	34.3	31.4
<b>SRH03</b> (m <sup>2</sup> s <sup>-2</sup> )	111.2	99.7



290

**Figure 8: As in Figure 7, but for (a) MUCAPE, SBCAPE and MLCAPE. (b) MUCIN, SBCIN and MLCIN.**

295


It is well known that due to limited vertical resolution reanalyses do not represent capping inversions very well (Nevius and Evans, 2018; Coffey et al., 2020; Taszarek et al., 2021). Here, CIN distribution in the parcel measures MUCIN, SBCIN and MLCIN are displayed (Figure 8b). The SP-HAIL and SP-NONHAIL differences for MLCIN distributions are statistically significant (Table 2). However, SBCIN and MUCIN differences are not statistically significant, presenting negative skew distributions for both events. The SP-HAIL events show a capping inversion layer (Figure 8b) with a MLCIN median of  $-20.2 \text{ J kg}^{-1}$  (Table 3), and of  $-12.2 \text{ J kg}^{-1}$  for SP-NONHAIL. These results are slightly lower than those obtained by Taszarek et al. (2020b), who found MLCIN values of  $-30 \text{ J kg}^{-1}$  for large hail events in Central Europe. A stronger CIN may delay the convective initiation until the CAPE is maximized; once the convection triggers, discrete convective modes, including isolated and elevated supercells, can be developed producing large hail in the plains (Rasmussen and Blanchard, 1998; Smith et al., 2012; Thompson et al., 2012; Taszarek et al., 2020b). However, the CIN values in this survey are lower, probably due to the complex orography of the domain (Markowski and Dotzek, 2011). Therefore, a mechanical trigger (e.g., air parcels lifted by orography or low-level convergence wind) is required to force initiation of convection to overcome the LCL. The conjunction of these factors favours great vertical motions and organized convection.

300

305

A comparison of MLLCL, MLLFC, FZH and FZH\_W percentiles and distributions between SP-HAIL and SP-NONHAIL is performed in Figure 9. The MLLCL has been an important discriminator between tornadic and non-tornadic supercells in the US (Rasmussen and Blanchard, 1998; Thompson et al., 2003). Taszarek et al. (2020b) compare the MLLCL in severe


convective storms with hail greater than 5 cm in US and Europe obtaining high similarities between both continents. However, in Europe, the MLLCL tends to have much less variability on supercell events, so the skill of this indicator is limited (Kahraman et al., 2017; Taszarek et al., 2020b). The results here shown for Spain are in line with the previous conclusion, since there are no statistically significant differences (Table 2) for MLLCL between SP-HAIL and SP-NONHAIL events. Rodriguez and Bech (2018) also observed this low MLLCL variability between tornadic and non-tornadic convective storms in the Iberian Peninsula. Nevertheless, Púčik et al. (2015) obtained a MLLCL median value of 1000 m for severe hail-thunderstorms in central Europe, that matches those herein described for SP-HAIL events (Table 3). On the other hand, the MLLFC does not show significant differences between SP-HAIL and SP-NONHAIL (Table 2) with similar median values and variability. The MLLFC and MLLCL median values (Table 3) are also similar to those obtained by Taszarek et al. (2020) for Europe. Another important factor for SP-HAIL and SP-NONHAIL events is the freezing level, with differences between both types of supercells being statistically significant for FZH and FZH\_W (Table 2). Both FZH and FZH\_W distributions for SP-NONHAIL events present higher variability than for SP-HAIL ones (Figure 9) with FZH and FZH\_W median values for SP-HAIL higher than for SP-NONHAIL (Table 3).



**Figure 9: As in Figure 7, but for MLLCL, MLLFC, FZH and FZH\_W.**

Several studies (Rasmussen and Blanchard, 1998; Púčik et al., 2015; Taszarek et al., 2019) suggest that the severity of the convective storms depend on the relationship between CAPE and WS. Furthermore, deep-moist convection tends to develop more organized systems as the WS intensifies (Markowski and Richardson 2010). WS between 0-6 km and EBWD distributions for SP-HAIL and SP-NONHAIL events are displayed (Figure 10). As expected, the WS06 and EBWD values are higher for SP-HAIL than for SP-NONHAIL with statistically significant differences in EBWD (Table 2). The WS06 and EBWD median values for SP-HAIL (Table 3) agree with Taszarek et al. (2020b) results for severe convective storms with large hail in Europe; nevertheless, the values for the US are higher than those presented here. Despite this, supercell events in

330 the domain of study can be explained by the presence of several mountain ranges in the study area (Figure 1a) where the WS is enhanced by the interaction of the wind field with orography, with a similar mechanism as observed in the Alps (Kunz et al., 2018; Taszarek et al., 2020). However, ERA5 is limited to reproduce this enhancement due to its horizontal resolution.



335 **Figure 10: As in Figure 7, but for WS06 and EBWD.**

Storm-relative helicity (SRH) is a frequent parameter used for forecasting supercells and tornadoes since it quantifies the cyclonic updraft rotation in right and left moving supercells (in this survey only the right-moving measure is used; Davies-Jones et al., 1990; Bunkers et al., 2002). Higher SRH values are usually related to the development of the mesocyclones and large hail formation (Rasmussen and Blanchard, 1998; Thompson et al., 2003). However, Hannesen et al. (1998); Kaltenbock (2004) and Gascón et al. (2015) suggested that complex orography such as in the current study domain can increase directional shear at low levels due to flow disturbance. Resulting large SRH03 values would produce favourable but not necessary conditions for severe storms formations. The SRH01 and SRH03 distributions (Figure S2, Supplementary) show no statistically significant differences (Table 2) between SP-HAIL and SP-NONHAIL. However, the SRH01 and SRH03 median values are slightly higher for SP-HAIL than for SP-NONHAIL (Table 3), showing similar variability. There are remarkable SRH03 90th percentile values for both groups of events, at  $225 \text{ m}^2 \text{ s}^{-2}$ , in line with Kahmaran et al. (2017), who state that large hail occurrences are associated with large values of SRH03. Rodriguez and Bech (2018, 2021) obtained similar SRH03 values for EF0 and EF1 tornadoes and waterspouts in Iberia. The SRH03 results are also consistent with Calvo-Sancho et al. (2021), where the SRH03 median spatial distribution in Spain displays values of almost  $100 \text{ m}^2 \text{ s}^{-2}$  in the eastern half of Spain, and with Taszarek et al. (2020b) in Europe for large hail reports.

340  
345



## 350 4 Summary and conclusions

The environments of SP-HAIL and SP-NONHAIL events are here characterized and compared in Spain from 2011 to 2020. Different atmospheric variables are retrieved from the ERA5 reanalysis to obtain the synoptic patterns and sounding composites **at the formation time of the supercell's**. Thermodynamic and kinematic parameters related to convective environments are also calculated and compared between SP-HAIL and SP-NONHAIL events.

355 The results yield several conclusions; the most important are listed below:

- There are notable differences in the spatial and monthly distributions of supercells in Spain. The eastern half of Spain accumulates 79.9% of the SP-NONHAIL and 88.3% of the SP-HAIL events. July and August accumulate 53.3% of the SP-NONHAIL and 74.4% of the SP-HAIL supercells. **Most events initiate between 12:00 and 19:00 UTC with a peak at 15:00.**

360 composites at 500 hPa, with the largest height gradients corresponding to SP-HAIL. Strong upper-level forcing is promoted by vorticity advection and upper-level divergence. Surface humidity is influenced by the 10-meters winds, being higher for SP-HAIL. The conjunction of these factors with wind convergences allows the convection initiation.

- The T2M and DWPT values are related to supercell monthly distributions with higher values corresponding to the warm season and minimum values to the cool season. Both variables are statistically different between SP-HAIL and SP-  
365 NONHAIL, being larger for the first group.

- **Environments of SP-HAIL events are characterized by approximately two times larger MUCAPE median values than for SP-NONHAIL events. Moreover, higher CAPE, FZH and EBWD values and lower MLCIN results are found for SP-HAIL with respect to SP-NONHAIL. The differences for these parameters between both events are statistically significant at  $p < 0.05$ .**

370 • Based on the ERA5 characterization results for SP-HAIL events in Spain, 75% of the supercells present **T2M > 21.1 °C, DWPT > 12.0 °C, MUCAPE > 781 J kg<sup>-1</sup>, MLCIN < -8.7 J kg<sup>-1</sup>, MLLFC > 1421 m, FZH > 2986.3 m, FZH\_W > 2497.5 m, WS06 > 13.7 m s<sup>-1</sup> and SRH03 > 74.7 m<sup>2</sup> s<sup>-2</sup>.**

Finally, the orography and convective environments have been revealed as important factors to supercell formation and development. Thus, although ERA5 resolution improves previous reanalyses, more research is needed with high-resolution  
375 models, allowing the study of the interactions between large-scale and convection processes in the genesis and development of hail supercell events.

### Data availability

ERA5 reanalysis is available from the Copernicus Climate Change Service Climate Data Store (<https://doi.org/10.24381/cds.bd0915c6>, Hersbach et al., 2018). Spanish Supercell Database upon request to the author.

## 380 **Author contributions**

CC-S, JD-F, YM, MLM, PB and JG-A designed the study. CC-S and JD-F performed the analysis and wrote the first manuscript. PB, JG-A, MS, and MLM supervised and review. DSM and JF provided computational and software support. All authors discussed the results and edited the manuscript.

## **Conflicts of Interest**

385 The authors declare that there are no conflicts of interest regarding the publication of this paper.

## **Acknowledgements.**

This work was partially supported by the following research projects: PID2019-105306RB-I00 (IBERCANES project), CGL2016-78702-C2-1-R and CGL2016-78702-C2-2-R (SAFEFLIGHT project), FEI-EU-17-16 and SPESMART AND SPESVALE (ECMWF Special Projects). Carlos Calvo-Sancho and Javier Díaz-Fernández acknowledge the grant supported  
390 by the Spanish Ministerio de Ciencia, Innovación y Universidades (FPI programs PRE2020-092343 and BES-2017-080025, respectively). Y. Martin acknowledges the grant supported by the European Union (Marie Skłodowska-Curie Programs 101019424). **We thank the three anonymous reviewers for their time and their detailed and constructive comments that helped to improve the presentation of our results.**

## **References**

- 395 Antonescu, B., Schultz, D. M., Holzer, A., & Groenemeijer, P. (2017). Tornadoes in Europe: An underestimated threat. *Bulletin of the American Meteorological Society*, 98(4), 713-728.
- Bech, J., Pineda, N., Rigo, T., Aran, M., Amaro, J., Gayà, M., ... & van der Velde, O. (2011). A Mediterranean nocturnal heavy rainfall and tornadic event. Part I: Overview, damage survey and radar analysis. *Atmospheric research*, 100(4), 621-637.
- 400 Bedka, K., Murillo, E. M., Homeyer, C. R., Scarino, B., & Mersiovsky, H. (2018). The above-anvil cirrus plume: An important severe weather indicator in visible and infrared satellite imagery. *Weather and Forecasting*, 33(5), 1159-1181.
- Bolgiani, P., Santos-Muñoz, D., Fernández-González, S., Sastre, M., Valero, F., & Martín, M. L. (2020). Microburst detection with the WRF model: Effective resolution and forecasting indices. *Journal of Geophysical Research: Atmospheres*, 125(14), e2020JD032883.
- 405 Brooks, H. E., Lee, J. W., & Craven, J. P. (2003). The spatial distribution of severe thunderstorm and tornado environments from global reanalysis data. *Atmospheric Research*, 67, 73-94.
- Brooks, H. E., Anderson, A. R., Riemann, K., Ebberts, I., & Flachs, H. (2007). Climatological aspects of convective parameters from the NCAR/NCEP reanalysis. *Atmospheric Research*, 83(2-4), 294-305.

- Brooks, H. E., Doswell III, C. A., Zhang, X., Chernokulsky, A. A., Tochimoto, E., Hanstrum, B., ... & Barrett, B. (2019). A century of progress in severe convective storm research and forecasting. *Meteorological Monographs*, 59, 18-1.
- 410 Calvo-Sancho, C. and Martín, Y. (2021). Supercell Pre-convective Environments in Spain: a dynamic downscaling of ERA-5 Reanalysis, EGU General Assembly 2021, online, 19–30 Apr 2021, EGU21-2967, <https://doi.org/10.5194/egusphere-egu21-2967>, 2021.
- Calvo-Sancho, C. (2021). Caracterización de los ambientes convectivos que favorecen el desarrollo de la convección organizada en España: exploración y evolución en el período 1979-2019 mediante reanálisis ERA5. Final Master Thesis. Universidad de Zaragoza. <http://dx.doi.org/10.13140/RG.2.2.25129.31841>.
- 415 **Carlson, T. N., & Ludlam, F. H. (1968). Conditions for the occurrence of severe local storms. *Tellus*, 20(2), 203-226.**
- Castro, A., Sánchez, J. L., & Fraile, R. (1992). Statistical comparison of the properties of thunderstorms in different areas around the Ebro-Valley (Spain). *Atmospheric research*, 28(3-4), 237-257.
- Chernokulsky, A., Kurgansky, M., Mokhov, I., Shikhov, A., Azhigov, I., Selezneva, E., ... & Kühne, T. (2020). Tornadoes in northern Eurasia: From the middle age to the information era. *Monthly Weather Review*, 148(8), 3081-3110.
- 420 Coffey, B.E., Taszarek, M., & Parker, M.D. (2020). Near-Ground Wind Profiles of Tornadic and Nontornadic Environments in the United States and Europe from ERA5 Reanalyses. *Weather and Forecasting* 35, 2621–2638. <https://doi.org/10.1175/WAF-D-20-0153.1>
- Consortio de Compensación de Seguros. (2020). Estadística de Riesgos Extraordinarios. Serie 1971–220. Accessed January 425 30, 2022. [http://www.conorseguros.es/web/documents/10184/44193/Estadistica\\_Riesgos\\_Extraordinarios\\_1971\\_2014/14ca6778-2081-4060-a86d-728d9a17c522](http://www.conorseguros.es/web/documents/10184/44193/Estadistica_Riesgos_Extraordinarios_1971_2014/14ca6778-2081-4060-a86d-728d9a17c522).
- Dahl, J. M. (2006). Supercells: their dynamics and prediction (Doctoral dissertation).
- Davis-Jones, R., Burgess, D.W., and Foster, M. (1990): Test of helicity as a forecast parameter. Preprints, 16th Conf. on Severe 430 Local Storms, Kananaskis Park, AB, Canada, Amer. Meteor. Soc., 588–592.
- Dotzek, N., Groenemeijer, P., Feuerstein, B., & Holzer, A. M. (2009). Overview of ESSL's severe convective storms research using the European Severe Weather Database ESWD. *Atmospheric research*, 93(1-3), 575-586.
- Duda, J. D., & Gallus, W. A. (2010). Spring and summer midwestern severe weather reports in supercells compared to other morphologies. *Weather and forecasting*, 25(1), 190-206. DOI: 10.1175/2009WAF2222338.1.
- 435 Edwards, R., Allen, J. T., & Carbin, G. W. (2018). Reliability and climatological impacts of convective wind estimations. *Journal of Applied Meteorology and Climatology*, 57(8), 1825-1845.
- Elmore, K. L., Flamig, Z. L., Lakshmanan, V., Kaney, B. T., Farmer, V., Reeves, H. D., & Rothfus, L. P. (2014). mPING: Crowd-sourcing weather reports for research. *Bulletin of the American Meteorological Society*, 95(9), 1335-1342.
- Farnell, C., Rigo, T., & Pineda, N. (2018). Exploring radar and lightning variables associated with the Lightning Jump. Can 440 we predict the size of the hail?. *Atmospheric Research*, 202, 175-186.

- Galanaki, E., Lagouvardos, K., Kotroni, V., Flaounas, E., & Argiriou, A. (2018). Thunderstorm climatology in the Mediterranean using cloud-to-ground lightning observations. *Atmospheric Research*, 207, 136-144.
- García-Ortega, E., Merino, A., López, L., & Sánchez, J. L. (2012). Role of mesoscale factors at the onset of deep convection on hailstorm days and their relation to the synoptic patterns. *Atmospheric Research*, 114, 91-106.
- 445 **Gascón, E., Merino, A., Sánchez, J. L., Fernández-González, S., García-Ortega, E., López, L., & Hermida, L. (2015). Spatial distribution of thermodynamic conditions of severe storms in southwestern Europe. *Atmospheric Research*, 164, 194-209.**
- Gatzen, C. P., Fink, A. H., Schultz, D. M., & Pinto, J. G. (2020). An 18-year climatology of derechos in Germany. *Natural Hazards and Earth System Sciences*, 20(5), 1335-1351.
- Gayà, M. (2011). Tornadoes and severe storms in Spain. *Atmospheric Research* 100 (4):334–43. doi: 10.1016/j. Atmosres.2010.10.019.
- 450 Gensini, V. A., Converse, C., Ashley, W. S., & Taszarek, M. (2021). Machine Learning Classification of Significant Tornadoes and Hail in the United States Using ERA5 Proximity Soundings. *Weather and Forecasting*, 36(6), 2143-2160.
- Groenemeijer, P., Púčik, T., Holzer, A. M., Antonescu, B., Riemann-Campe, K., Schultz, D. M., ... & Sausen, R. (2017). Severe convective storms in Europe: Ten years of research and education at the European Severe Storms Laboratory. *Bulletin of the American Meteorological Society*, 98(12), 2641-2651.
- 455 Gropp, M. E., & Davenport, C. E. (2018). The impact of the nocturnal transition on the lifetime and evolution of supercell thunderstorms in the Great Plains. *Weather and Forecasting*, 33(4), 1045-1061.
- Gutierrez, R. E., & Kumjian, M. R. (2021). Environmental and Radar Characteristics of Gargantuan Hail-Producing Storms. *Monthly Weather Review*, 149(8), 2523-2538.
- 460 Hamid, K. (2012). Investigation of the passage of a derecho in Belgium. *Atmospheric research*, 107, 86-105.
- Hannesen, R., Dotzek, N., Gysi, H., & Beheng, K. D. (1998). Case study of a tornado in the Upper Rhine valley. *Meteorologische Zeitschrift-Berlin-*, 7, 163-170.**
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., Thépaut, J. (2020). The ERA5 global reanalysis. *Q.J.R. Meteorol. Soc.* 146, 1999–2049. <https://doi.org/10.1002/qj.3803>.
- Kahraman, A., Kadioglu, M., & Markowski, P. M. (2017). Severe convective storm environments in Turkey. *Monthly Weather Review*, 145(12), 4711-4725.
- 470 **Kaltenböck, R. (2004). The outbreak of severe storms along convergence lines northeast of the Alps. Case study of the 3 August 2001 mesoscale convective system with a pronounced bow echo. *Atmospheric research*, 70(1), 55-75.**
- Kaltenböck, R., Diendorfer, G., Dotzek, N. (2009). Evaluation of thunderstorm indices from ECMWF analyses, lightning data and severe storm reports. *Atmos. Res.* 93 (1–3), 381–396.

- 475 Kotroni, V., & Lagouvardos, K. (2016). Lightning in the Mediterranean and its relation with sea-surface temperature. *Environmental Research Letters*, 11(3). <https://doi.org/10.1088/1748-9326/11/3/034006>
- Krennert, T., Pistotnik, G., Kaltenberger, R., & Csekits, C. (2018). Crowdsourcing of weather observations at national meteorological and hydrological services in Europe. *Advances in Science and Research*, 15, 71-76.
- Kuchera, E. L., & Parker, M. D. (2006). Severe convective wind environments. *Weather and forecasting*, 21(4), 595-612.
- 480 Kunz, M., Blahak, U., Handwerker, J., Schmidberger, M., Punge, H. J., Mohr, S., ... & Bedka, K. M. (2018). The severe hailstorm in southwest Germany on 28 July 2013: Characteristics, impacts and meteorological conditions. *Quarterly Journal of the Royal Meteorological Society*, 144(710), 231-250.
- Kunz, M., Wandel, J., Fluck, E., Baumstark, S., Mohr, S., & Schemm, S. (2020). Ambient conditions prevailing during hail events in central Europe. *Natural Hazards and Earth System Sciences*, 20(6), 1867-1887.
- 485 Li, F., Chavas, D. R., Reed, K. A., & Dawson II, D. T. (2020). Climatology of severe local storm environments and synoptic-scale features over North America in ERA5 reanalysis and CAM6 simulation. *Journal of Climate*, 33(19), 8339-8365.
- Lombardo, K. A., & Colle, B. A. (2011). Convective storm structures and ambient conditions associated with severe weather over the northeast United States. *Weather and forecasting*, 26(6), 940-956.
- López, L., & Sánchez, J. L. (2009). Discriminant methods for radar detection of hail. *Atmospheric Research*, 93(1-3), 358-490 368.
- Mann, H. B., & Whitney, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. *The annals of mathematical statistics*, 50-60.
- Manzato, A. (2012). Hail in northeast Italy: Climatology and bivariate analysis with the sounding-derived indices. *Journal of Applied Meteorology and Climatology*, 51(3), 449-467.
- 495 Markowski, P. M., & Dotzek, N. (2011). A numerical study of the effects of orography on supercells. *Atmospheric research*, 100(4), 457-478.
- Markowski, P., & Richardson, Y. (2011). *Mesoscale meteorology in midlatitudes (Vol. 2)*. John Wiley & Sons.
- Martín, Y., Cívica, M., & Pham, E. (2020). Constructing a Supercell Database in Spain Using Publicly Available Two-Dimensional Radar Images and Citizen Science. *Annals of the American Association of Geographers*, 0(0), 1-21. <https://doi.org/10.1080/24694452.2020.1812371>.
- 500 Melcón, P., Merino, A., Sánchez, J. L., López, L., & García-Ortega, E. (2017). Spatial patterns of thermodynamic conditions of hailstorms in southwestern France. *Atmospheric Research*, 189, 111-126.
- Merino, A., García-Ortega, E., López, L., Sánchez, J. L., & Guerrero-Higueras, A. M. (2013). Synoptic environment, mesoscale configurations and forecast parameters for hailstorms in Southwestern Europe. *Atmospheric Research*, 122, 183-505 198.
- Miglietta, M.M., Mazon, J. & Rotunno, R. (2017) Numerical simulations of a tornadic supercell over the Mediterranean. *Weather and Forecasting*, 32, 1209-1226. <https://doi.org/10.1175/WAF-D-16-0223.1>.


- Mohr, S., Kunz, M., & Geyer, B. (2015). Hail potential in Europe based on a regional climate model hindcast. *Geophysical Research Letters*, 42(24), 10-904.
- 510 Mohr, S., Kunz, M., Richter, A., & Ruck, B. (2017). Statistical characteristics of convective wind gusts in Germany. *Natural Hazards and Earth System Sciences*, 17(6), 957-969.
- Mora, M., Riesco, J., de Pablo Dávila, F., & Rivas Soriano, L. (2015). Atmospheric background associated with severe lightning thunderstorms in Central Spain. *International Journal of Climatology*, 35(4), 558-569.
- Mulholland, J. P., Peters, J. M., & Morrison, H. (2021). How does LCL height influence deep convective updraft width?.  
515 *Geophysical Research Letters*, 48(13), e2021GL093316.
- National Weather Service (NWS). (2019). Supercell structure and dynamics. Accessed 21 January 2021. <https://www.weather.gov/lmk/supercell/dynamics>.
- Nevis, D. S., & Evans, C. (2018). The influence of vertical advection discretization in the WRF-ARW Model on capping inversion representation in warm-season, thunderstorm-supporting environments. *Weather and Forecasting*, 33(6), 1639-  
520 1660.
- Nisi, L., Martius, O., Hering, A., Kunz, M., & Germann, U. (2016). Spatial and temporal distribution of hailstorms in the Alpine region: a long-term, high resolution, radar-based analysis. *Quarterly Journal of the Royal Meteorological Society*, 142(697), 1590-1604.
- Půčík, T., Groenemeijer, P., Rýva, D., & Kolář, M. (2015). Proximity soundings of severe and nonsevere thunderstorms in  
525 central Europe. *Monthly Weather Review*, 143(12), 4805-4821.
- Quirantes, J. A. (2003). Tornado F2/F3 y Supercélula de Alcañiz (Teruel) [EN: F2/F3 Tornado and Supercell in Alcañiz, Teruel]. Accessed 24 January 2022. <http://www.tiemposevero.es/ver-reportaje.php?id=35>.
- Quirantes Calvo, J. A., Riesco Martín, J., & Núñez Mora, J. Á. (2014). Características básicas de las supercélulas en España. Agencia Estatal de Meteorología (AEMET).
- 530 Rasmussen, E. N., & Blanchard, D. O. (1998). A baseline climatology of sounding-derived supercell and tornado forecast parameters. *Weather and forecasting*, 13(4), 1148-1164.
- Rodríguez, O., & Bech, J. (2018). Sounding-derived parameters associated with tornadic storms in Catalonia. *International Journal of Climatology*, 38(5), 2400-2414.
- Rodríguez, O., & Bech, J. (2021). Tornadic environments in the Iberian Peninsula and the Balearic Islands based on ERA5  
535 reanalysis. *International Journal of Climatology*, 41, E1959-E1979.
- Romero, R., Ramis, C., Alonso, S., Doswell III, C. A., & Stensrud, D. J. (1998). Mesoscale model simulations of three heavy precipitation events in the western Mediterranean region. *Monthly weather review*, 126(7), 1859-1881.
- Shaltout, M., & Omstedt, A. (2014). Recent sea surface temperature trends and future scenarios for the Mediterranean Sea. *Oceanologia*, 56(3), 411-443.

- 540 Smith, B. T., Thompson, R. L., Grams, J. S., Broyles, C., & Brooks, H. E. (2012). Convective modes for significant severe thunderstorms in the contiguous United States. Part I: Storm classification and climatology. *Weather and Forecasting*, 27(5), 1114-1135.
- Taszarek, M., Brooks, H. E., & Czernecki, B. (2017). Sounding-derived parameters associated with convective hazards in Europe. *Monthly Weather Review*, 145(4), 1511-1528.
- 545 Taszarek, M., Brooks, H. E., Czernecki, B., Szuster, P., & Fortuniak, K. (2018). Climatological aspects of convective parameters over Europe: A comparison of ERA-interim and sounding data. *Journal of Climate*, 31(11), 4281–4308. <https://doi.org/10.1175/JCLI-D-17-0596.1>
- Taszarek, M., Allen, J., Púčik, T., Groenemeijer, P., Czernecki, B., Kolendowicz, L., ... Schulz, W. (2019). A climatology of thunderstorms across Europe from a synthesis of multiple data sources. *Journal of Climate*, 32(6), 1813–1837. <https://doi.org/10.1175/JCLI-D-18-0372.1>
- 550 Taszarek, M., Allen, J. T., Groenemeijer, P., Edwards, R., Brooks, H. E., Chmielewski, V., & Enno, S. E. (2020a). Severe convective storms across Europe and the United States. Part I: Climatology of lightning, large hail, severe wind, and tornadoes. *Journal of Climate*, 33(23), 10239–10261. <https://doi.org/10.1175/JCLI-D-20-0345.1>
- Taszarek, M., Allen, J.T., Púčik, T., Hoogewind, K.A., & Brooks, H.E. (2020b). Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments Associated with Lightning, Large Hail, Severe Wind, and Tornadoes. *Journal of Climate* 33, 10263–10286. <https://doi.org/10.1175/JCLI-D-20-0346.1>.
- Taszarek, M., Pilgaj, N., Allen, J. T., Gensini, V., Brooks, H. E., & Szuster, P. (2021). Comparison of convective parameters derived from ERA5 and MERRA-2 with Rawinsonde data over Europe and North America. *Journal of Climate*, 34(8), 3211-3237.
- 560 Thompson, R. L., Edwards, R., Hart, J. A., Elmore, K. L., & Markowski, P. (2003). Close proximity soundings within supercell environments obtained from the Rapid Update Cycle. *Weather and Forecasting*, 18(6), 1243-1261.
- Thompson, R. L., Smith, B. T., Grams, J. S., Dean, A. R., & Broyles, C. (2012). Convective modes for significant severe thunderstorms in the contiguous United States. Part II: Supercell and QLCS tornado environments. *Weather and forecasting*, 27(5), 1136-1154.
- 565 Tomas-Burguera, M., Beguería, S., & Vicente-Serrano, S. M. (2021). Climatology and trends of reference evapotranspiration in Spain. *International Journal of Climatology*, 41, E1860-E1874.
- Trapp, R. J. (2013). *Mesoscale-convective processes in the atmosphere*. Cambridge University Press.
- Tullot, I. F. (2000). *Climatología de España y Portugal* (Vol. 76). Universidad de Salamanca.
- Vicente-Serrano, S. M., Azorin-Molina, C., Sanchez-Lorenzo, A., Revuelto, J., López-Moreno, J. I., González-Hidalgo, J. C., ... & Espejo, F. (2014). Reference evapotranspiration variability and trends in Spain, 1961–2011. *Global and Planetary Change*, 121, 26-40.
- 570 Weisman, M. L., & Klemp, J. B. (1982). The dependence of numerically simulated convective storms on vertical wind shear and buoyancy. *Monthly Weather Review*, 110(6), 504-520.

575 Westermayer, A., Pucik, T., Groenemeijer, P., & Tijssen, L. (2016). Comparison of sounding observations and reanalysis of  
thunderstorm environments. In Eighth European Conf. on Severe Storms.



Supplement



580

Figure S1. 700-400 hPa Q-vector divergence values (contours, Pa-1 s-3) and convergence of Q-vectors (arrows) composites at  $t_0$ , for (a) SP-NONHAIL, (b) SP-HAIL and (c) Differences between SP-HAIL and SP-NONHAIL. Black points in plot c) denote statistically significant differences (p-value < 0.05) in Q-vector divergence values.




Figure S2. SRH01 and SRH03 distributions and boxplots for SP-HAIL and SP-NONHAIL at  $t_0$ . The median is represented as a red point.