

1 **Replies document for reviews of:**

2 **The role of cyclones and PV cutoffs for the occurrence of unusually long wet spells in**
3 **Europe**

4 Matthias Röthlisberger,¹ Barbara Scherrer,¹ Andries Jan de Vries,^{1,a} Raphael Portmann¹

5
6
7 ¹ Institute for Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland

8 ^a now at Institute of Earth Surface Dynamics, University of Lausanne, Lausanne, Switzerland

9
10 *Corresponding author:* Matthias Röthlisberger, matthias.roethlisberger@env.ethz.ch

11
12 **General comments to the Reviewers**

13 We would like to thank both reviewers for their thoughtful, encouraging and constructive
14 comments. Below we reply to each comment and describe the changes to the manuscript
15 resulting from them. In particular Reviewer 2 requested a substantial number of additional
16 figures. We have therefore decided to accompany the main manuscript with a supplement,
17 which contains the additional figures. Furthermore, two additional figures are included at the
18 end of this document for reviewer reference. Moreover, substantial changes we made to the
19 revised manuscript include the following. (1) We now consider additional variables in the
20 discussion of our case studies, namely the integrated vapor transport as well as the quasi-
21 geostrophic vertical motion at 500 hPa, forced from the atmospheric layers above 550 hPa
22 (hereafter IVT and QG ω , respectively). Moreover, we currently explore whether a quantitative
23 measure of the tropospheric static stability can be meaningfully incorporated in this study. (2)
24 We discovered that the plotting routine we used to produce the original Figs. 8 and 9 (NCL)
25 only drew a hatching (to indicate statistical significance) when multiple neighbouring grid
26 points were deemed significant. This led to the impression that many **F**, **N**, **P** and **R** values
27 were not significant while in fact they should have been labelled as significant in the original
28 Figs. 8 and 9, based on our non-parametric test. We now plot the significance information by
29 masking out insignificant grid points in the revised Figs. 8 and 9 as well as in Figs. S6–9.

30 All other comments have also been addressed and were particularly useful to more clearly
31 present our results. Reviewer comments are included below in blue font colour and our replies
32 in black.

33 **Reviewer 1**

34 This paper presents an assessment of long wet spells across Europe and their association with
35 PV cutoffs and extratropical cyclones. I find this an interesting study that contributes to the
36 breadth of knowledge of extreme event drivers, although provides only small amounts of new
37 scientific insight that has not been documented in the already published literature. Overall, I
38 have very few comments as this study presents four case studies and the dynamics surrounding
39 them. My main query is regarding to the anomalies presented in Figs 8 and 9. I find the method
40 of calculating the climatology unusual and expand more upon this below. Furthermore, I
41 question the use of ERA-Interim reanalysis when the newer and higher resolution ERA5 has
42 been readily available for some time now. Once the authors address my comments I recommend
43 this manuscript for publication as I believe it will suit the journal well. My main points to be
44 addressed can be found below.

45

46 **Comments**

47 1. L50 – the reference needs re-formatting. The comma should not be there.

48

49 Ok, changed as requested, thank you for spotting this typo.

50

51 2. L55-65 – I would re-phrase this paragraph. The way it is introduced suggests that features
52 such as WCB, fronts, cyclones are individual features, when this is rarely the case and they are
53 often all part of one synoptic system. I appreciate the authors do mention this toward the end
54 of the paragraph, however I think this could be phrased better.

55

56 This paragraph has been rephrased to more clearly emphasize that these features are
57 dynamically related and often occur in association with one another.

58

59 3. L135-143 – The choice of ERA-Interim as an analysis dataset is a confusing one. Newer
60 reanalysis products such as ERA5 have been readily available for several years now and using
61 a more up-to-date product, with higher resolution would surely be beneficial for a study such
62 as this. The specific dynamics and features that would be resolved would increase and also the
63 issues with precipitation mentioned by the authors may be reduced. Have the authors tested
64 their selection of the wet spells to the different precipitation products? Would there be different
65 climatologies in Figs. 1, 7, 8, 9 as a result?

66

67 We see the rationale of this comment but we still believe that we have good reasons for the
68 choice of the used data set. We very intentionally worked with precipitation data from a
69 reanalysis data set as opposed to observational precipitation data, because the purpose of the
70 study is to examine the role of cyclones and cutoffs for unusually long wet spells, and thus the
71 consistency between the precipitation data and the SLP, wind, PV fields etc. is crucial.
72 However, we agree with the reviewer that using ERA5 data in this study would, in principle,
73 be desirable. The main reason for choosing ERA-Interim instead, though, is that the Portmann
74 et al. (2021) PV cutoff climatology is only available for ERA-Interim and cannot easily be
75 adapted to ERA5, due to very large computational costs arising from the sophisticated PV cutoff
76 tracking routine which is part of the Portmann et al. (2021) algorithm. In the following three
77 arguments we motivate our choice for using ERA-Interim data and elaborate on why using
78 ERA-Interim as opposed to ERA5 is not expected to affect the qualitative findings of this study.

79
80 (1) The Portmann et al. (2021) PV cutoff identification algorithm involves a sophisticated
81 three dimensional Lagrangian tracking routine, which is based on kinematic air parcel
82 trajectories. This tracking scheme is clearly distinct and, in our opinion, superior to other
83 tracking routines for cutoffs (e.g., Bell and Bosart 1989; Nieto et al. 2005; Pinheiro et
84 al. 2017; Muñoz et al. 2020), for three reasons. First, it is the only one that uses the PV
85 framework, while others are based on relative vorticity and/or geopotential height.
86 Second, the tracking uses kinematic air parcel trajectories and quasi-conservation of PV
87 on isentropic surfaces. These two reasons render this approach particularly consistent
88 with fundamental principles in atmospheric dynamics. In addition, this trajectory-based
89 approach also works in regions where cutoffs move rapidly, for example near the jet
90 stream where consecutive features do not always overlap spatially. And third, it allows
91 for three dimensional feature tracking and therefore circumvents any dependence on the
92 choice of a vertical level. This is important because cutoffs often strongly evolve in their
93 vertical structure (e.g., Portmann et al. 2018) and can therefore only be meaningfully
94 tracked with a three-dimensional tracking scheme. However, this tracking scheme is
95 computationally very expensive already for ERA-Interim, and applying it to ERA5
96 would further increase the computational costs by a factor of 24 (6 times higher
97 temporal resolution, four times more grid points per model level), which we do not
98 consider feasible at the current stage.

100 (2) We expect that synoptic to large-scale flow structures such as cyclones and upper-level
101 PV cutoffs are well resolved already in ERA-Interim, as the major improvement from
102 ERA-Interim to ERA5 lies in the resolution of smaller-scale processes and weather
103 features. Note that throughout the manuscript we investigate these synoptic to large-
104 scale flow structures and not their smaller-scale substructure. Therefore, we do not
105 expect that discrepancies in ERA-Interim and ERA5 cyclones and cutoffs would
106 question our conclusions on a qualitative level.

107
108 (3) We only rely on the ERA-Interim precipitation field for identifying the wet spells,
109 whose characteristics are displayed in Fig. 1. Reproducing Fig. 1 with ERA5 data (Fig.
110 A1, at the end of this document) reveals no drastic differences between the wet spell
111 duration, accumulated precipitation, mean precipitation rate or seasonality compared to
112 ERA-Interim. Especially for this reason, we do not expect a major benefit from
113 reproducing the analysis using ERA5.

114
115 In summary, we share the reviewers view that the use of ERA5 in principle would be preferable
116 for the current study, in particular if the PV cutoff climatology of Portmann et al. (2021) were
117 available for ERA5. However, for the three reasons above we believe that using ERA-Interim
118 is justified given the purpose of this study.

119
120 4. L175 – How sensitive are the results to the choice of mask radius/distance from gridpoint?
121 Why did the authors choose 400km?

122
123 We tested radii of 200 to 600 km for both weather features and now included the respective
124 results as Figs S6–9 in the Supplemental Material. There is little qualitative sensitivity to the
125 exact choice of the radius r for this range of values. A smaller radius than 200 km or larger
126 radius than 600 km seems unjustified based on synoptic experience. Both cyclones and cutoffs
127 can surely induce precipitation further away than 200 km from the identified mask, e.g., along
128 trailing cold fronts (of cyclones) or downstream of propagating cutoffs, where the quasi-
129 geostrophic forcing for ascent can be expected to be largest. Furthermore, if r is increased
130 beyond 600 km then during some time steps most of the study domain is “allegedly” under the
131 influence of cyclones and cutoffs and almost all precipitation would be attributed to either of
132 these systems. This also does not seem justified, as, e.g., in summer over complex topography
133 the majority of precipitation falls due to diurnal convection (Rüdisühli et al. 2020).

134

135 In the revised manuscript we mention our sensitivity analysis for r and the results for $r = 200$
136 km and $r = 600$ km.

137

138 5. L194-201 – I find the choice of how the climatologies created confusing. From my
139 interpretation you take all the days of the year that the wet spells occur (from start to end) and
140 create the climatology based on those days of the year? Firstly, how many days of the year are
141 in the climatology of each grid point – surely this varies depending on the average length of the
142 spell and how likely the spells are to overlap/be in the same season. Secondly, would it make
143 sense to have the climatology for all wet days and then the anomalies would be for how the
144 unusually wet days differ from just wet days? On this, the wet spells in summer are also likely
145 averaging some significantly warm (and cyclone-less) days as well, do these skew the
146 anomalies significantly? Is the question of the paper how do unusually long wet spells differ
147 from wet periods, or from all other days in general? This needs to be made clearer in the
148 introduction.

149

150 Both reviewer raised concerns with regard to the computation of climatological values for F ,
151 N , P and R for the two weather systems. We therefore adopted a suggestion of Reviewer 2,
152 which was to compute the climatological values simply as the mean values over the respective
153 Monte Carlo sample. These new climatological values do not differ substantially from the
154 original ones, but given that both reviewers found our original approach somewhat confusing
155 we chose to adapt this simpler definition of climatological F , N , P and R in order to increase
156 the clarity of our approach. The number of days contained in this new climatological values
157 still vary in space depending on how long the S_{20} are at each grid point (as correctly noticed by
158 the reviewer), but this seems justified given the strongly differing seasonality and duration of
159 the S_{20} . In the revised manuscript (last paragraph of Section 2.5) we now more explicitly
160 mention the variable number of days in the climatology. Moreover, the new climatological
161 values reflect climatological values for wet and dry days and we now also specifically mention
162 this fact in the revised manuscript (last paragraph of Section 2.5). The anomalies in the original
163 and revised Figs. 8 and 9 therefore inform about differences in cyclone/cutoff characteristics
164 during the S_{20} and average conditions during the respective days of the year. Constructing a
165 climatology solely of wet days would be a valid alternative, which, however, would help to
166 address a slightly different research question, namely how cyclone/cutoff characteristics differ

167 during the S_{20} from average wet day conditions. We believe both research questions are
168 worthwhile, but it is the first of these two research questions that we would like to address here.

169

170 6. Fig. 1 – it would be good to also show the variation in the length of the extreme wet spells.
171 How much does this variation skew the averages shown in this figure? Would the median be a
172 better choice for some of the panels? L294-295 (and throughout) – are the numbers quotes for
173 $N_{cyclone}$ and $F_{cyclone}$ statistically significant? If not then this does not suggest that these
174 wet spells feature unusual synoptic conditions.

175

176 To visualize this variability amongst the S_{20} panels (a-d) of the new Supplemental Figure 1
177 now show the duration and accumulated precipitation of the longest spell per grid point (S_1)
178 and the twentieth longest wet spell (S_{20}). Moreover, we follow the advice of the reviewer and
179 show in the revised Fig. 1a,b the median duration of the S_{20} (Fig. 1a) and the median
180 accumulated precipitation (Fig. 1b).

181

182 Furthermore, the reviewer is right in noticing that statistically not significant $N_{cyclone}$ or
183 $F_{cyclone}$ do not indicate unusual cyclone characteristics. However, the purpose of quantifying
184 the four cyclone and cutoff characteristics is not just to identify significant departures from the
185 local climatology. Rather these quantities also inform about regional differences in the roles of
186 cutoffs and cyclones. For the latter reason the four quantities are included in the case study
187 discussions, irrespective of whether or not they statistically significantly differ from respective
188 local climatological values.

189

190 7. Fig. 5 – please define the Streamers in the figure caption and the text. These are not
191 introduced prior to this in the text and therefore should be explained.

192

193 Ok, we now introduce the term PV streamer more clearly in the revised introduction, following
194 Appenzeller and Davies (1992).

195

196 8. L463-464 – I would argue that the residence times are somewhat similar for the UK and the
197 Italian seas. I'd rephrase this paragraph to reflect the lack of differences in this field.

198

199 Agreed, in the revised manuscript the sentence reads: “A further region with particularly
200 noteworthy cyclone characteristics (and their anomalies) during the \mathcal{S}_{20} are found in the seas
201 around Italy.”

202

203 **Reviewer 2**

204 In this study, Röthlisberger et al. examine, through illustrative case studies and systematic
205 climatological analysis, the role of cyclones and PV cutoffs for the occurrence of unusually
206 long wet spells in Europe, defined as the 20 longest wet spells at each grid point in the ERA
207 Interim reanalysis during 1979–2018. Overall, I found the manuscript to be well-written, and I
208 believe that the topic has substantial scientific merit. In addition, the results may help to inform
209 future work on predictability and climate change impacts for these events. Increased
210 understanding of the synoptic-scale dynamical processes and weather systems that result in very
211 long wet spells is clearly needed. The motivation for the study, the data and methods, and the
212 results are described in a clear and concise manner. The figures are, for the most part,
213 straightforward to interpret and support the conclusions drawn in the text.

214 In my review, I came up with a number of minor comments, suggestions, and questions for the
215 authors to consider. Once these have been satisfactorily addressed, I believe that this manuscript
216 will be acceptable for publication in *WCD*.

217 **Comments**

218 1. Line 88–89: A brief discussion of the dynamical link between PV streamers and cutoffs and
219 the process of Rossby wave breaking is needed here to provide a basis for later discussions of
220 wave breaking and the formation of PV streamers and cutoffs throughout section 3.
221 Accordingly, a basic definition of Rossby wave breaking in the text would also be helpful.

222 Ok, we added a brief discussion in the introduction of the revised manuscript.

223 2. Line 146: “(large-scale and convective)” It would be better to explicitly state that the
224 precipitation amounts analyzed in this study are the sum of the large-scale and convective
225 precipitation in the ERA-Interim data.

226 Ok, changed as requested.

227 3. Line 147: Is there a particular reason why you limited the analysis to the 20 longest wet
228 spells? Would not the statistics be more robust if you were to include, e.g., the 50 longest wet
229 spells instead?

230 The reviewer is right in noticing that a larger sample of events would increase the robustness
231 of our statistical analyses. However, the purpose of this study is to examine *unusually long* wet
232 spells and this speaks against substantially increasing the sample size for two reasons: (1) A
233 long wet spell can be unusual in the sense of being a rare event (hereafter “rareness criterion”),
234 i.e., an event for which the average waiting time until a comparable event occurs (i.e., the return
235 period) is comparatively long. This rareness criterion is important for the motivation of this
236 study, because an event that occurs multiple times per year is much less likely to cause
237 significant impacts than an event with a return period of several years. With the 20 longest
238 spells in a 40-year period, we sample events whose return period typically exceeds two years.
239 Substantially increasing the number of spells would violate the rareness criterion for an
240 “unusually long wet spell”. (2) The analyzed spells should also be exceptional with regard to
241 their duration in comparison to all other spells at the same grid point (hereafter “exceptionality
242 criterion”). However, in some regions in Europe the total number of multiday wet spells (based
243 on our 5mm/day criterion) is simply too small for substantially increasing the number of
244 analyzed spells without violating the exceptionality criterion. To make this second aspect
245 explicit, the new Supplementary Figure 1e shows the total number of identified spells
246 (minimum duration of two days) and reveals that this number is highly variable in space. Along
247 the Norwegian west coast, the spell count exceeds 900, however, e.g., around Crimea or the
248 westernmost part of the Mediterranean fewer than 200 multi-day wet spells occurred. Over most
249 regions, though, the total spell count exceeds 200, which means that fewer than 10% of all
250 multi-day wet spells are considered in this study. Based on Fig. S1e and the arguments made
251 above we believe that a sample size of 20 spells is a reasonable compromise between the
252 statistical robustness of the results and the unusualness of the spells we analyze.

253 Note further that we tested the sensitivity of our results to the sample size by considering the
254 top five and top ten longest spells at each grid point. These sensitivity tests reveals no qualitative
255 difference to the results in Figs. 8 and 9, however, as anticipated by the reviewer, the statistical
256 significance of the results were much reduced (not shown).

257 4. Line 175: How was this radius determined to be suitable for this analysis? How sensitive are
258 the results to this radius? I suspect there are situations in which cyclones or cut-offs play an

259 important dynamical role in a wet spell at a given location but are located farther than 400-km
260 from the location. Of course in this type of analysis you need to choose discrete thresholds to
261 define events/ features and to examine relationships between them. I am not arguing that you
262 should change this radius, but I do think some discussion regarding why it was chosen would
263 be helpful here.

264 We tested radii of 200, 400 and 600 km and found little qualitative differences, even though
265 quantitatively, the value of r of course strongly affects F , N , P and R . Analogous figures to the
266 revised Figs. 8 and 9 but for radii of 200 km and 600 km are now included in the supplement
267 (Figs S6–9). Importantly, none of our key findings in this analysis (spatial variability of F , N ,
268 P and R ; sign- and significance/non-significance of anomalies of F , N , P and R) are altered
269 qualitatively when varying the radius from 200 to 600 km. The exact choice of 400 km is based
270 on synoptic expertise, which suggests that both cyclones and cutoffs can cause precipitation
271 more than 200 km away from the identified mask, e.g., along trailing cold fronts for cyclones.
272 However, considering areas beyond 600 km around the identified masks as directly affected by
273 the cyclones/cutoffs appears inappropriate too, as with such a radius almost all precipitation
274 might be attributed to either of the two systems. For instance, it is well known that summer
275 precipitation over complex topography is often due to diurnal convection that is not directly
276 due to cyclones or cutoffs (e.g., Rüdüsühli et al. 2020).

277 We now explicitly mention the robustness of these results to variations in the radius in the
278 Section 2.5 and, as mentioned above, include analogous figures to the revised Figs. 8 and 9 but
279 for radii of 200 and 600 km as Figs. S6–9 in the supplement.

280 5. Line 194–201: I find this explanation a bit confusing. It is not clear to me how climatological
281 values for the number of distinct cyclones per spell are computed in this manner if all days of
282 the year in any year corresponding to the $S20(x,y)$ are grouped. Perhaps I am misunderstanding
283 the explanation of the methodology. It might be better to use the mean from the 1000-sample
284 Monte Carlo distribution at each grid point as the "climatological value" as the Monte Carlo
285 approach that you apply retains information about the consecutive days comprising each
286 individual spell in the $S20$ sample.

287 Both reviewers raised concerns with regard to how climatological values of F , N , P and R were
288 originally computed. We therefore decided to adopt this reviewer suggestion and now simply
289 compute the climatological F , N , P and R for both weather systems as the mean over the
290 respective Monte Carlo samples. The anomalies in the revised Figs. 8 and 9 and Figs S6–9 were

291 computed with respect to these new climatologies. Moreover, we rephrased large parts of
292 Section 2.5 (former Section 2.4) to better explain the Monte Carlo simulations as well as the
293 computation of anomalies of F , N , P and R .

294 6. Line 216: How much variability is there in the duration of the S20 cases at each location?
295 Are there some locations where the duration is highly variable between the S20 cases?

296 Yes, the variability amongst the S20 is substantial, as we are sampling wet spells that are in the
297 upper tail of the wet spell duration distribution. To illustrate this variability Supplementary
298 Figure 1a-d now shows the duration (a,c) and accumulated precipitation (b,d) of the spells with
299 rank 1 and 20, respectively. Moreover, in the revised manuscript we explicitly mention this
300 large variability when discussing Fig.1.

301 7. Line 217: A map of the terrain elevation over the domain in Fig. 1 could aid the reader in
302 interpreting the results.

303 Supplementary Fig 1f now shows the ERA-Interim topography in the study region.

304 8. Line 235: Are the climatological percentiles computed for all wet days in all months of the
305 year, or do they vary seasonally based on when the wet spell occurred?

306 All wet days of the year. This is now mentioned explicitly in the revised Section 3.1.

307 9. Line 258: An explanation is needed here regarding why these four particular cases and
308 locations were selected.

309 In the last sentence of Section 3.1. we explicitly state that we selected these cases subjectively
310 due to their archetypal nature, out of a much larger set of cases we analyzed.

311 10. Line 260: I really appreciate the concise yet information-dense synoptic analysis and
312 discussion for the four selected wet spells. A main criticism I have for this study is that the
313 synoptic analysis does not include quantitative diagnostics of processes and ingredients by
314 which the cutoffs and cyclones support the persistent precipitation. While these processes are
315 at times inferred or surmised in the text, no diagnostics for moisture transport, forcing for
316 ascent, convective instability are provided. Inclusion of additional fields and diagnostics for
317 each case would, of course, result in an increase in the number of figures and in the complexity

318 of the manuscript, so perhaps it is outside the scope of this study. Could additional analyses and
319 diagnostics be provided as online supplemental materials instead?

320 The reviewer correctly notices that excessive analyses of additional variables would push the
321 length of the paper beyond reasonable limits. Moreover, our synoptic discussion of the case
322 studies is based on well-known effects of cyclones and cutoffs on static stability and
323 precipitation formation, which are also not contested by either of the reviewers. Therefore, we
324 think it would be excessive to add another four figures to the main manuscript (one per case
325 study) to quantify what is qualitatively apparent already in the current Figs. 3–6. Nevertheless
326 we have compiled ERA-Interim climatologies of IVT and $QG\omega$ at 500 hPa, forced from the
327 atmospheric layers above 550 hPa as in (Graf et al. 2017) and now show these additional
328 (quantitative) diagnostics for all four case studies at the same time steps as in Figs 3–6 as Figs.
329 S2–5. Moreover, we have rephrased parts of Section 3.2 to also discuss these new figures. As
330 expected, the new figures generally support quantitatively what was qualitatively evident
331 already from the original Figs. 3–6. Furthermore we currently explore whether we can
332 meaningfully incorporate a quantitative measure of tropospheric static stability in Figs S2–5.

333 11. Line 260: Perhaps this is outside the scope of your study, but I wonder if it is possible to
334 include analysis and/or discussion of the large-scale/planetary-scale conditions that contributed
335 to the occurrence of the four selected wet spells. Were persistent weather regimes in place over
336 the Atlantic/Europe region that favored the synoptic-scale dynamical processes operating in
337 each archetypal synoptic story line?

338 Interesting comment. We specifically and very much intentionally narrowed the scope of this
339 study to cyclones and PV cutoffs, but of course the reviewer is right in mentioning that large-
340 scale/planetary-scale conditions would be interesting too. However, in particular with the
341 rather numerous other additional analyses that were requested by the reviewers, we feel that
342 including further analyses of weather regimes and/or large-scale modes of variability would
343 go beyond the scope of this study.

344 12. Line 280: It could be useful to include more fields in the composite analyses for the four
345 locations. Possible additional fields include sea level pressure anomalies, PV anomalies, and
346 frequency anomalies of cyclones and cutoffs. Such additional fields could provide a more
347 detailed, complete picture of the synoptic-scale conditions conducive to the S20 cases at each
348 location.

349 For the revised Fig. 7 we removed the cyclone/cutoff tracks and show instead composites of
350 SLP anomalies, IVT anomalies and anomalies of $QG\omega$. We feel that these additional variables
351 further clarify the composite structure of the S_{20} and therefore significantly add to the value
352 of Fig. 7. Many thanks for this comment!

353 13. Line 288: The information density in Figs. 3–6 is high. Overall, I think this is fine; I am
354 able to read and interpret the figures fairly well. I do, however, recommend drawing the
355 geographical boundaries and the SLP in different colors. This could help the reader
356 distinguish the SLP field, especially when contours for several fields are superimposed.

357 Ok, we changed the color of the geographical boundaries to a lighter gray.

358 14. Line 297: You clearly and convincingly describe how recurrence and/or persistence of
359 weather systems is key to the long durations of the four wet spells analyzed. I propose that
360 Hovmoller diagrams of, say, upper-level PV anomalies or upper-level meridional wind
361 anomalies overlaid by the cyclone and cutoff masks averaged in some latitude band would
362 help to more clearly illustrate the recurrence and persistence during the spells. These diagrams
363 would nicely complement the plan-view analyses in Figs. 3–6.

364 Interesting comment, thank you. We produced the respective Hovmöller diagrams of 250 hPa
365 meridional wind, with feature tracks overlaid (Fig. A2, at the end of this document). For the
366 Norway case study the Hovmöller diagram together with the cyclone tracks is indeed
367 underlining the discussion of this case in the manuscript. There is clearly recurrent ridge
368 formation over the North Atlantic, which is associated with the recurrent passage of (fast
369 moving) cyclones. For the other cases, though, the meridional wind signal induced by the
370 cutoffs/PV streamers does not come out very clearly, presumably due to the complex shape of
371 these features (see Figs. 3, 5 and 6). Also, the cyclone/cutoff tracks do reveal to some extent
372 the stationarity of the respective features, but due to the intricacies of feature tracking (e.g.,
373 merger and splitting events) they are difficult to interpret. We therefore think that Fig. A2
374 creates more confusion than clarity and refrain from including it in the main manuscript.

375 15. Line 343: Physically this makes sense because PV cut-offs often form in association with
376 Rossby wave breaking that results in meridional elongation of PV streamers equatorward of
377 the midlatitude jet/waveguide. This location is too far north to be frequently impacted by
378 cutoffs.

379 Yes, we agree mostly, although not all cutoffs in the Portmann et al. (2021) climatology
380 necessarily need to form from anticyclonic wave breaking. In fact, Portmann et al. (2021)
381 found that, aside from the aforementioned classical storyline of cutoff formation, they also
382 frequently form from cyclonic Rossby wave breaking associated with extratropical cyclones
383 in the storm track regions (i.e. poleward of the jet). According to Portmann et al. (2021), the
384 PV cutoff frequency in DJF over Norway is between 7-9%, which is not substantially less
385 than the 9-11% in the Mediterranean region. We therefore chose to keep our original,
386 somewhat more general wording.

387 16. Line 367: What processes were conducive to recurrent wave breaking/cutoff formation in
388 this case? It seems that the recurrence was associated with temporal clustering of cyclone
389 developments and associated ridge building upstream along the waveguide over the North
390 Atlantic. Was this flow evolution related to an anomalous configuration of the North Atlantic
391 waveguide? It may be worthwhile to briefly speak to the upstream processes that result in
392 recurrent wave breaking.

393 Interesting and certainly valid comment. However, we do not see a very straight forward way
394 to determine what exactly was conducive to the recurrent wave breaking. To thoroughly
395 address this question, simulations and/or statistical analyses of a large-enough sample of
396 similar episodes would be required. We feel that such analyses would clearly go beyond the
397 scope of this study. Moreover, we would like to refrain from simply hypothesizing about
398 these causes, as without the aforementioned statistical/model-based analyses, hypothesizing is
399 really all we could do.

400 17. Line 430: I understand the justification for plotting all of the PV cutoff and cyclone tracks
401 in this figure. However, I find it very difficult to make sense of the messy bundles of tracks in
402 the maps, with the exception being Fig. 7b in which the tracks are mostly zonal. Is there a
403 way to more clearly illustrate the track information? Alternatively, could the tracks be
404 removed from these figures without compromising the discussion?

405 For the revised Fig. 7 we removed the cyclone/cutoff tracks and now show instead composites
406 of SLP anomalies, IVT anomalies and anomalies of $QG\omega$.

407 18. Line 461: Can you briefly explain why the Pcyclone quantity is anomalously low nearly
408 everywhere on the map?

409 This simply means that cyclones occur at a higher rate during the S20 compared to
410 climatology, because the rate at which distinct cyclones occur is the inverse of the cyclone
411 period P .

412 **Technical corrections:**

413 Line 9: Define “PV” acronym here?

414 Ok

415 Line 67: “For unusually long-lasting wet spells, it is much less clear how and in association
416 with which weather systems they form.” This sentence is a bit clunky. Here is a possible
417 alternative: “The mechanisms and weather systems contributing to the occurrence of
418 unusually long-lasting wet spells are less clear.”

419 Ok

420 Line 67–68: Change “Only few” to “few”

421 Ok

422 Line 109: Remove “responsible”

423 Ok

424 Line 163: Insert “the method of” before “Portmann”

425 Ok

426 Line 223: Insert “daily” before “precipitation rate”

427 Ok

428 Line 379: Start a new sentence with “The fifth”

429 Ok

430 Line 446: Insert “tend to” after “do not”

431 We prefer our original formulation.

432 Line 522: Replace the colon with a period.

433 Ok

434 Line 525: Check whether "e.g.," is needed here.

435 We think it is ok as it is.

436 Line 528: Replace "behaviour" to "characteristics" ?


437 Ok

438 **References**

- 439 Appenzeller, C., and H. C. Davies, 1992: Structure of stratospheric intrusions into the
440 troposphere. *Nature*, **358**, 570–572, doi:10.1038/358570a.
- 441 Bell, G. D., and L. F. Bosart, 1989: A 15-year climatology of Northern Hemisphere 500 mb
442 closed cyclone and anticyclone centers. *Mon. Weather Rev.*, **117**, 2142–2163,
443 doi:https://doi.org/10.1175/1520-0493(1989)117<2142:AYCONH>2.0.CO;2.
- 444 Graf, M. A., H. Wernli, and M. Sprenger, 2017: objective classification of extratropical
445 cyclogenesis. *Q. J. R. Meteorol. Soc.*, **143**, 1047–1061, doi:10.1002/qj.2989.
- 446 Muñoz, C., D. Schultz, and G. Vaughan, 2020: A midlatitude climatology and interannual
447 variability of 200- and 500-hPa cut-off lows. *J. Clim.*, **33**, 2201–2222, doi:10.1175/JCLI-
448 D-19-0497.1.
- 449 Nieto, R., and Coauthors, 2005: Climatological features of cutoff low systems in the Northern
450 Hemisphere. *J. Clim.*, **18**, 3085–3103, doi:10.1175/JCLI3386.1.
- 451 Pinheiro, H. R., K. I. Hodges, M. A. Gan, and N. J. Ferreira, 2017: A new perspective of the
452 climatological features of upper-level cut-off lows in the Southern Hemisphere. *Clim.*
453 *Dyn.*, **48**, 541–559, doi:10.1007/S00382-016-3093-8/TABLES/1.
- 454 Portmann, R., B. Crezee, J. Quinting, and H. Wernli, 2018: The complex life-cycles of two
455 long-lived potential vorticity cutoffs over Europe. *Q. J. R. Meteorol. Soc.*, **144**, 701–719,
456 doi:10.1002/qj.3239.
- 457 Portmann, R., M. Sprenger, and H. Wernli, 2021: The three-dimensional life cycles of potential
458 vorticity cutoffs: A global and selected regional climatologies in ERA-Interim (1979–
459 2018). *Weather Clim. Dyn.*, **2**, 507–534, doi:10.5194/WCD-2-507-2021.
- 460 Rüdüsühli, S., M. Sprenger, D. Leutwyler, C. Schär, and H. Wernli, 2020: Attribution of
461 precipitation to cyclones and fronts over Europe in a kilometer-scale regional climate
462 simulation. *Weather Clim. Dyn.*, **1**, 675–699, doi:https://doi.org/10.5194/wcd-1-675-2020.
- 463 Ventura, V., C. J. Paciorek, and J. S. Risbey, 2004: Controlling the proportion of falsely rejected
464 hypotheses when conducting multiple tests with climatological data. *J. Clim.*, **17**, 4343–
465 4356, doi:10.1175/3199.1.
- 466 Wilks, D. S., 2016: “The stippling shows statistically significant grid points”: How research
467 results are routinely overstated and overinterpreted, and what to do about it. *Bull. Am.*
468 *Meteorol. Soc.*, **97**, 2263–2273, doi:10.1175/BAMS-D-15-00267.1.
- 469


470 **Additional figures for reviewer reference**

471



472

473 Fig A1: As (the revised) Fig. 1 but for ERA5. Note that contrary to the original Fig. 1 panels
474 (a) and (b) show the median duration and median accumulated precipitation of the ERA5 S₂₀.



475

476

477

478

479

480

481

482

483

484

Fig A2: Hovmöller plots of 250 hPa meridional wind (v) for the four case studies (a) $S_1(37^\circ E, 55^\circ N)$, (b) $S_1(14^\circ E, 66^\circ N)$, (c) $S_1(12^\circ E, 43^\circ N)$ and (d) $S_1(25^\circ E, 48^\circ N)$. The meridional wind v has been averaged in a latitude band of $\pm 15^\circ$ latitude around the latitude of the respective spell, i.e., in (a) $40^\circ N$ – $70^\circ N$, in (b) $51^\circ N$ – $81^\circ N$, in (c) $28^\circ N$ – $58^\circ N$ and in (d) $33^\circ N$ – $63^\circ N$. Horizontal lines depict the start and end dates of the respective spell, while the vertical line in each panel denote the longitude of the respective spell. Purple and yellow lines indicate the longitude–time tracks of all cyclones and cutoffs whose masks overlapped with a circle of radius 400 km around the location of the respective wet spell. In panel (b) no cutoff tracks are shown due to the insignificance of cutoffs for the synoptic storyline of this spell.